

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
1 April 2004 (01.04.2004)

PCT

(10) International Publication Number
WO 2004/026846 A1

(51) International Patent Classification⁷: **C07D 253/10**,
401/12, 403/12, A61K 31/53, A61P 35/00

(21) International Application Number:
PCT/NZ2003/000210

(22) International Filing Date:
17 September 2003 (17.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
521436 17 September 2002 (17.09.2002) NZ

(71) Applicants (for all designated States except US): **AUCKLAND UNISERVICES LIMITED** [NZ/NZ]; Level 10, 70 Symonds Street, Auckland (NZ). **THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY** [US/US]; 900 Welch Road, Suite 350, Palo Alto, CA 94304 1850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BROWN, Martin, J.** [US/US]; 2010 Arbryan Way, Redwood City, CA 94061 (US). **DENNY, William, Alexander** [NZ/NZ]; 165 Gossamer Drive, Pakuranga, Auckland, 1706 (NZ). **HAY, Michael, Patrick** [NZ/NZ]; 52 St Leonard's Road, Epsom, Auckland 1003 (NZ). **HICKS, Kevin, Owen** [NZ/NZ]; 26 Duke Street, Mount Roskill, Auckland 1004 (NZ). **GAM-AGE, Swarnalatha, Akuratiya** [NZ/NZ]; Cancer Society

Research Laboratory, 14B Oakdale Road, Hillsborough, Auckland (NZ). **PRULJIN, Frederik, Bastiaan** [NL/NZ]; 1052 Scenic Drive North, Swanson, Auckland 1008 (NZ). **WILSON, William, Robert** [NZ/NZ]; 199 Smith Road, RD2, Waiuku (NZ).

(74) Agent: **BALDWIN SHELSTON WATERS**; P.O. Box 852, Wellington (NZ).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DNA-TARGETED BENZOTRIAZINE 1,4-DIOXIDES AND THEIR USE IN CANCER THERAPY

(57) Abstract: The present invention relates to DNA-targeted 1,2,4-benzotriazine- 1,4-dioxides and related analogues, to their preparation, and to their use as hypoxia-selective drugs and radiosensitizers for cancer therapy, both alone or in combination with radiation and/or other anticancer drugs.



WO 2004/026846 A1

DNA-TARGETED BENZOTRIAZINE 1,4-DIOXIDES AND THEIR USE IN CANCER THERAPY

REFERENCE TO GOVERNMENT CONTRACT

- 5 The invention described herein was made in the course of work under grant or contract from the United States Department of Health and Human Services. The United States Government has certain rights to this invention.

TECHNICAL FIELD

- 10 The present invention relates to DNA-targeted 1,2,4-benzotriazine-1,4-dioxides and related analogues, to their preparation, and to their use as hypoxia-selective drugs and radiosensitizers for cancer therapy, both alone or in combination with radiation and/or other anticancer drugs.

15 BACKGROUND TO THE INVENTION


- It has been established that many human tumors contain a significant hypoxic fraction of cells (Kennedy et al., *Int. J. Radiat. Oncol. Biol. Phys.*, **1997**, 37, 897-905; Movsas et al., *Urology*, **1999**, 53, 11-18). The presence of hypoxic cells arises because of chaotic growth and an inefficient microvasculature system within the tumor, which leads to
- 20 large intercapillary distances and variable blood flow. Reduction of oxygen tension in tumors leads to radioresistance. This reduction of oxygen tension causes up to a three-fold increase in radiation dose being required to kill anoxic tumor cells. A link has been identified between the presence of tumor hypoxia and failure of local control by radiation therapy (Brizel et al., *Radiother. & Oncol.*, **1999**, 53, 113-117). This
- 25 phenomenon of tumor hypoxia has been exploited in the development of a class of anticancer agents termed 'bioreductive drugs' (Brown et al., *Semin. Radiat. Oncol.*, 1966, 6, 22-36; Denny et al., *Br. J. Cancer*, **1996**, 74 (Suppl. XXVII) 32-38; Stratford & Workman, *Anti-Cancer Drug Des.*, **1998**, 13, 519-528). These agents are selectively active against hypoxic cells in tumors by targeting the DNA of these cells. The agents
- 30 cause irreversible damage to the DNA of the tumor cells, thereby causing the destruction and breakdown of the tumor.

Tirapazamine (TPZ, 3-amino-1,2,4-benzotriazine 1,4-dioxide) is a bioreductive agent (Kelson et al., *Anti-Cancer Drug Des.*, **1998**, 13, 575-592; Lee et al., WO 9104028,

April 1991) and is undergoing clinical trials in combination with radiotherapy and various chemotherapeutics, notably cisplatin (Denny & Wilson, *Exp. Opin. Invest. Drugs*, **2000**, 9, 2889-2901).

- 5 TPZ is activated by one electron reductases (Patterson et al., *Anti-Cancer Drug Des.* **1998** 13, 541-573; Denny & Wilson, *Exp. Opin. Invest. Drugs*, **2000**, 9, 2889-2901) to form a radical anion (Scheme A). This TPZ radical anion may be oxidized back to TPZ by molecular oxygen under aerobic conditions.

Scheme A.



10

Under hypoxic conditions the radical or species ultimately derived from TPZ can interact with DNA, although the exact mechanism is unclear (Jones et al., *Cancer Res.*, **1996**, 56, 1584-1590; Daniels et al., *Chem. Res. Toxicol.*, **1998**, 11, 1254-1257; Hwang et al., *Biochem.*, **1999**, 38, 14248-14255). TPZ causes DNA double-strand breaks under anoxic conditions (Jones et al., *Cancer Res.*, **1996**, 56, 1584-1590) and these results correlate with cytotoxicity (Dorie et al., *Neoplasia*, **1999**, 1, 461-467). Reversible one-electron reduction of TPZ that gives rise to a reactive radical species that is thought to be the basis for selective toxicity to hypoxic cells. Two electron reduction of TPZ or further reduction of the TPZ radical produces the metabolite 1-oxide (SR 4317) and further reduction gives the nor-oxide (SR 4330) (Baker et al., *Cancer Res.*, **1988**, 48, 5947-5952; Laderoute & Rauth, *Biochem Pharmacol.*, **1986**, 35, 3417-3420) (Scheme A). The metabolites (SR 4317) and (SR 4330) are both inactive under aerobic or hypoxic conditions.


20

It is also known that reactive species can be effectively targeted to DNA by attachment to DNA-affinic carriers. Thus, the intrinsic cytotoxicities and *in vivo* potencies of aniline mustards can be significantly increased (up to 100-fold), and the usual dependence of cytotoxicity on mustard reactivity lowered, by targeting to DNA via a 9-aminoacridine carrier (Gourdie et al., *J. Med. Chem.*, **1990**, 33, 1177-1185). DNA alkylation patterns can also be significantly altered (Prakash et al., *Biochem.*, **1990**, 29, 9799-9807; Boritzki et al., *Chem. Res. Toxicol.*, **1994**, 7, 41-46). Alkylation of DNA by DNA-targeted compounds is more rapid than with the corresponding untargeted compounds (O'Connor et al., *Chem.-Biol. Int.*, **1992**, 85, 1-14). However, the extent of DNA binding needs to be carefully adjusted to achieve effective targeting without significantly compromising the transport/diffusion properties (Hicks et al., *J. Pharmacol. Exp. Therapeut.* **2001**, 297, 1088-1098; Hicks et. al, *Brit. J. Cancer.* **1997**, 76, 894-903). Binding ability can be varied by alteration of both the chromophore and substituents on the DNA targeted compound (Palmer et al., *J. Med. Chem.*, **1988**, 31, 707-712).

It is an object of the present invention to utilize DNA-affinic carriers in combination with benzotriazine 1,4-dioxides to target DNA for cancer therapy purposes, or to at least provide the public with a useful choice.

DISCLOSURE OF THE INVENTION

In a first aspect, the present invention provides a compound of Formula I,



wherein

Y₁ and Y₂ at one or more of the available carbons 5-8 on the benzo ring: are each independently selected from the following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

wherein each R is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the said optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and

wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted or extended by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and

wherein the DNA-targeting unit is any moiety of a molecular weight below 700 Daltons that has an association constant (K) for binding to double-stranded random-sequence DNA of >10³ M⁻¹ at an ionic strength of 0.01 M at 20 °C,

or a pharmacologically acceptable salt thereof.

5 The definition of the DNA targeting unit above refers to double-stranded random-sequence DNA. An example of such double-stranded random-sequence DNA is DNA extracted from calf thymus.

A preferred compound of Formula I is one in which X is NH or CH₂.


10 A further preferred compound of Formula I is one in which Y₁ and Y₂ each represent H.

A further preferred compound of Formula I is one in which Y₁ represents OMe

15 A preferred embodiment of Formula I are compounds wherein A is selected from -(CH₂)₆NH-, -(CH₂)₃NH(CH₂)₃NHCO-, -(CH₂)₃NMe(CH₂)₃NHCO-, -(CH₂)₃NH-, -(CH₂)₂NH(CH₂)₂NHCO- or -(CH₂)₂NMe(CH₂)₂NHCO-.

A further preferred embodiment of Formula I are compounds wherein the DNA-targeting unit is selected from one of formulae **II- XVI**,

20



wherein in structures **XI-XVI** R^6 is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NO_2 , NHR^7 , NR^7R^7 , SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$;

R^6 can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NH_2 , NHR^7 , NR^7R^7 , SH, SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

- wherein each R^7 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^8 , NH_2 , NHR^8 , NR^8_2 or $N(OH)R^8$ wherein each R^8 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN , CO_2H or SH ;
- 5 D represents up to four of the following groups as substituents at any available ring carbon position; H, R^9 , hydroxy, alkoxy, halogen, NO_2 , NH_2 , NHR^9 , NR^9_2 , SH , SR^9 , SO_2R^9 , CF_3 , CN , CO_2H , CO_2R^9 , CHO , COR^9 , $CONH_2$, $CONHR^9$ or $CONR^9R^9$, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino, wherein each R^9 is independently selected
- 10 from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^{10} , NH_2 , NHR^{10} , NR^{10}_2 or $N(OH)R^{10}$ wherein each R^{10} is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN , CO_2H or SH ;
- 15 and wherein any available ring carbon position of formulae **II** - **XVI** can also be optionally replaced by -N- when the valency and configuration of the formula allows, the point of attachment of formulae **II** - **XVI** to the A group defined above is represented by \blacklozenge ; and wherein in formulae **XI**, **XII**, , m is selected from 2, 3 or 4, and wherein in formulae **XI**, **XII**, **XV** and **XVI**, J is selected from CH or N;
- 20 and wherein in formulae **XIII** and **XIV** n is selected from 0, 1 or 2; and wherein in formulae **XV** and **XVI** o is selected from 1 and 2.

A preferred embodiment of formula I is one in which the DNA targeting unit is selected from one of formulae IV, V, VI, VII, VIII, or IX.

25 A preferred embodiment of formula I is one in which D of the DNA targeting unit of Formulae **II** - **X** is H or Me.

Further preferred compounds of formula I include the following

30 wherein X is $NH-$, Y_1 is H, Y_2 is H, A is $-(CH_2)_6NH-$, the DNA targeting unit represents formula VII and D is H;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

5 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is H;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

10 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula IV and D is H;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VI and D is H;

15 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is Me;

20 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula IX and D is Me;

wherein X is NH-, Y₁ is 7-MeOCH₂CH₂O-, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

25 wherein X is CH₂-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula XI and D is H;

30 wherein X is NH-, Y₁ is 7-Me, Y₂ is H, A is $-(CH_2)_3NMeH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

wherein X is NH-, Y₁ is 7-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA

targeting unit represents formula VI and D is H;

wherein X is NH-, Y₁ is 6-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

5

wherein X is NH-, Y₁ is 6-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VI and D is H;

10

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is H;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H;

15

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula XI and D is Me;

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is Me;


20

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H; and

wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is Me.

25

In a second aspect, the present invention provides a compound of Formula I',



wherein

Y₁ represents at one or more of the available carbons 5-8 on the benzo ring the following groups:

halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO,
5 COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

Y₃ is selected from the following groups halo, H, R, OR, NH₂, NHR, NR₂, SO₂R, CF₃, CN,
CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl,
10 alkylpiperazinyl and morpholino;

wherein each R of groups Y₁ and Y₃ is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂,
15 NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each
20 independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted
25 C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and

30 wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂ or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN,

CO₂H or SH; and wherein the optionally substituted C₂₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and
5 wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and

wherein the DNA-targeting unit is any moiety of a molecular weight below 700 Daltons
10 that has an association constant (K) for binding to double-stranded random-sequence DNA of $>10^3 \text{ M}^{-1}$ at an ionic strength of 0.01 M at 20 °C,

or a pharmacologically acceptable salt thereof.

15 The definition of the DNA targeting unit above refers to double-stranded random-sequence DNA. An example of such double-stranded random-sequence DNA is DNA extracted from calf thymus.


A preferred compound of Formula I' is one in which X is O, NH or CH₂.

20

A further preferred compound of Formula I' is one in which Y₁ represents H.

A preferred embodiment of Formula I' are compounds wherein A is selected from
-(CH₂)₆NH-, -(CH₂)₃NH(CH₂)₃NHCO-, -(CH₂)₃NMe(CH₂)₃NHCO-, -(CH₂)₃NH-,
25 (CH₂)₂NH(CH₂)₂NHCO- or -(CH₂)₂NMe(CH₂)₂NHCO-.

A further preferred embodiment of Formula I' are compounds wherein the DNA-targeting unit is selected from one of formulae II- XVI,



wherein in structures **XI - XVI** R^6 is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NO_2 , NH_2 , NHR^7 , NR^7R^7 , SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$;

R^6 can also be represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NH_2 , NHR^7 , NR^7R^7 , SH, SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S; wherein each R^7 is independently selected from an optionally substituted C_{1-4} alkyl

or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR⁸, NH₂, NHR⁸, NR⁸₂ or N(OH)R⁹⁸ wherein each R⁸ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH;

- 5 D represents up to four of the following groups as substituents at any available ring carbon position; H, R⁹, hydroxy, alkoxy, halogen, NO₂, NH₂, NHR⁹, NR⁹₂, SH, SR⁹, SO₂R⁹, CF₃, CN, CO₂H, CO₂R⁹, CHO, COR⁹, CONH₂, CONHR⁹ or CONR⁹R⁹, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino, wherein each R⁹ independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and
- 10 wherein the optional substituents are each independently selected from OH, OR¹⁰, NH₂, NHR¹⁰, NR¹⁰₂ or N(OH)R¹⁰ wherein each R¹⁰ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH;

- and wherein any available ring carbon position of formulae **II- XVI** can also be optionally
- 15 replaced by -N- when the valency and configuration of the formula allows, the point of attachment of formulae **II- XVI** to the A group defined above is represented by ♦; and wherein in formulae **XI** and **XII**, m is selected from 2, 3 or 4, and wherein in formulae **XI, XII, XV or XVI** J is selected from CH or N; and wherein in formulae **XIII** and **XIV** n is selected from 0, 1 or 2, and
- 20 wherein in formulae **XV** and **XVI** o is selected from 1 or 2.

A preferred embodiment of formula I' is one in which the DNA targeting unit is selected from one of formulae **III - IX**.

- 25 A preferred embodiment of formula I' is one in which D of the DNA targeting unit of Formulae **II - X** is H or Me.

Preferred compounds of formula I' include the following

- 30 wherein X is O-, Y is H, A is-(CH₂)₃NH(CH₂)₃NHCO-, the DNA targeting unit represents formula VI and D is H;

wherein X is O-, Y is H, A is-(CH₂)₃NMe(CH₂)₃NHCO-, the DNA targeting unit

represents formula VI and D is H;

wherein X is O-, Y is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H;

5

wherein X is O-, Y is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H;

wherein X is O-, Y is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

10

wherein X is O-, Y is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

wherein X is O-, Y is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is H;

15

wherein X is O-, Y is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is H;

20

wherein X is O-, Y is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is Me;

wherein X is O-, Y is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is Me;

25

wherein X is O-, Y is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is Me;

wherein X is O-, Y is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII and D is Me;

30

wherein X is O-, Y is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula IX and D is Me;

wherein X is O-, Y is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula IX and D is Me; and wherein X is O-, Y is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula IX and D is Me;

5

wherein X is O-, Y is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula IX and D is Me;

10 In a third aspect the invention provides for the use in a method of therapy for treating cancers including the step of administering a compound of Formula I as defined above or a compound of Formula I' as defined above or a mixture thereof in a therapeutically effective amount to tumour cells in a subject.

Preferably the tumour cells are in a hypoxic environment.

15

It is preferred that the method of therapy further includes the step of administering radiotherapy to the tumor cells before, during or after the administration of the compound of Formula I as defined above or a compound of Formula I' as defined above or a mixture thereof to the tumour cells.

20

It is preferred that the method of therapy further includes the step of administering one or more chemotherapeutic agents to the tumor cells before, during or after the administration of the compound of Formula I as defined above or a compound of Formula I' as defined above or a mixture thereof to the tumour cells.

25

While these compounds will typically be used in cancer therapy of human subjects, they can be used to target tumor cells in other warm blooded animal subjects such as other primates, farm animals such as cattle, and sports animals and pets such as horses, dogs, and cats.

30

A "therapeutically effective amount", is to be understood as an amount of a compound of Formula I as defined above or a compound of Formula I' as defined above or a mixture thereof that is sufficient to show benefit to a patient. The actual amount, rate and time-

course of administration, will depend on the nature and severity of the disease being treated. Prescription of treatment is within the responsibility of general practitioners and other medical doctors.

- 5 A hypoxic environment is to be understood as either an *in vitro* or *in vivo* environment having a poorer blood supply and lower oxygen tension than normal tissues.

It is to be understood that the compound of Formula I or Formula I' can be administered alone or in combination with other chemotherapeutic agents or treatments, especially
10 radiotherapy, either simultaneously or sequentially dependent upon the condition to be treated.

Preferred chemotherapeutic agents can be selected from:

- Cisplatin or other platinum-based derivatives,
15 Temozolomide or other DNA methylating agents,
Cyclophosphamide or other DNA alkylating agents,
Doxorubicin, mitoxantrone, camptothecin or other topoisomerase inhibitors,
Methotrexate, gemcitabine or other antimetabolites.

20 In a fourth aspect of the present invention there is provided a pharmaceutical composition including a therapeutically effective amount of a compound of formula I or compound of formula I' or a mixture thereof, a pharmaceutically acceptable excipient, adjuvant, carrier, buffer or stabiliser.

25 The pharmaceutically acceptable excipient, adjuvant, carrier, buffer or stabiliser should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which can be oral, or by injection, such as cutaneous, subcutaneous, or intravenous injection.

30 Pharmaceutical compositions for oral administration can be in tablet, capsule, powder or liquid form. A tablet may comprise a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water,


petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included. A capsule may comprise a solid carrier such as gelatin.

5

For intravenous, cutaneous or subcutaneous injection, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has a suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride injection, Ringer's injection, Lactated Ringer's injection. Preservatives, stabilisers, buffers antioxidants and/or other additives may be included as required.

10

In a fifth aspect of the present invention there is provided a method of making a compound of formula XVII



15

wherein

Y_1 and Y_2 at one or more of the available carbons 5-8 on the benzo ring; are each independently selected from the following groups: halo, H, R, OH, OR, NO_2 , NH_2 , NHR, NR_2 , SH, SR, SO_2R , CF_3 , CN, CO_2H , CO_2R , CHO, COR, CONH_2 , CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

20

wherein each R is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR^1 , NO_2 , NH_2 , NHR^1 , NR^1R^1 , SH, SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN, CO_2H , CO_2R^1 , CHO, COR^1 , CONH_2 , CONHR¹, CONR¹R¹;

25

R can also represent an optionally substituted aryl or an optionally substituted

heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more

5 heteroatoms in its ring system which are each independently selected from O, N or S; wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or

10 SH, and


A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN,

15 CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂,

20 NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH;

or a pharmacologically acceptable salt thereof,

including the step of coupling a compound (a) using a palladium reagent to form compound (b) which can then be converted into a compound of XVII as defined above;



25

wherein in compound (a)


V is halogen which is selected from Cl, Br or I and Y₁, Y₂ are as defined above; and wherein in compound (b) Y₁, Y₂ are as defined above, W is selected from an optionally substituted

C₁₋₁₂alkyl, optionally substituted C₂₋₁₂alkenyl, and optionally substituted C₂₋₁₂alkynyl group, wherein the optional substituents is selected from halo, OH, OR⁶, NO₂, NH₂, NHR⁶, NR⁶R⁶, SH, SR⁶, imidazolyl, R⁶-piperazinyl, morpholino, SO₂R⁶, CF₃, CN, CO₂H, CO₂R⁶, CHO, COR⁶, CONH₂, CONHR⁶, CONR⁶R⁶, wherein each R⁶ is

5 independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR⁷, NR⁷₂ or N(OH)R⁷ wherein each R⁷ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH.

10

In a sixth aspect of the present invention there is provided a method of making a compound of formula XVII'



15

wherein Y₁ represents at one or more of the available carbons 5-8 on the benzo ring the following groups:

halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

20

Y₃ is selected from the following groups H, R, OR, NH₂, NHR, NR₂, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino

25

wherein each R of groups Y₁ and Y₃ is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂,

NHR^1 , NR^1R^1 , SH , SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN , CO_2H , CO_2R^1 , CHO , COR^1 , CONH_2 , CONHR^1 , CONR^1R^1 ;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR^1 , NH_2 , NHR^1 , NR^1R^1 , SH , SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN , CO_2H , CO_2R^1 , CHO , COR^1 , CONH_2 , CONHR^1 , CONR^1R^1 , and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R^1 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH_2 , NHR^2 , NR^2 or $\text{N}(\text{OH})\text{R}^2$ wherein each R^2 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN , CO_2H or SH , and


wherein X represents NH, NMe, CH_2 , SO, SO_2 , or O;

A represents an optionally substituted C_{1-12} alkyl group wherein the optional substituents are each independently selected from OH, OR, NH_2 , NHR^3 , NR^3 or $\text{N}(\text{OH})\text{R}^3$ wherein each R^3 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN , CO_2H or SH ; and wherein the optionally substituted C_{1-12} alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR^4 , CONH , CONR^4 , NHCO , NR^4CO , wherein each R^4 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional R^4 substituents are each independently selected from OH, OR, NH_2 , NHR^5 , NR^5 or $\text{N}(\text{OH})\text{R}^5$ wherein each R^5 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN , CO_2H or SH ; and

or a pharmacologically acceptable salt thereof;

including the steps of coupling a compound (a) using a palladium reagent to form

compound (b) which can then be converted into a compound of XVII' as defined above;



wherein in compound (a)

V is halogen which is selected from Cl, Br or I; Y₁, X and A are as defined above;

and wherein in compound (b) Y₁, X and A are as defined above,


5 W is selected from an optionally substituted

C₁₋₁₂alkyl, optionally substituted C₂₋₁₂alkenyl, and optionally substituted C₂₋₁₂alkynyl group, wherein the optional substituents is selected from halo, OH, OR⁶, NO₂, NH₂, NHR⁶, NR⁶R⁶, SH, SR⁶, imidazolyl, R⁶-piperazinyl, morpholino, SO₂R⁶, CF₃, CN, CO₂H, CO₂R⁶, CHO, COR⁶, CONH₂, CONHR⁶, CONR⁶R⁶, wherein each R⁶ is

10 independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR⁷, NR⁷₂ or N(OH)R⁷ wherein each R⁷ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH.

15

In a seventh aspect of the present invention there is provided a compound of formula XVIII



20 wherein

Y₁ and Y₂ at one or more of the available carbons 5-8 on the benzo ring: are each independently selected from the following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

wherein each R is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹,
 5 imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹,
 10 imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;


wherein each R¹ is independently selected from an optionally substituted
 15 C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and

20 wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN,
 25 CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂,
 30 NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH;
 or a pharmacologically acceptable salt thereof.

In an eighth aspect of the present invention there is provided a compound of formula

XVII'



wherein

- 5 Y₁ represents at one or more of the available carbons 5-8 on the benzo ring the following groups:

halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

10

Y₃ is selected from the following groups H, R, OR, NH₂, NHR, NR₂, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino

- 15 wherein each R of groups Y₁ and Y₃ is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

- 20 R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more
- 25 heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R²

wherein each R^2 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH, and

wherein X can represent NH, NMe, CH_2 , SO, SO_2 , or O;

5

A can represent an optionally substituted C_{1-12} alkyl group wherein the optional substituents are each independently selected from OH, OR, NH_2 , NHR^3 , NR^3_2 or $N(OH)R^3$ wherein each R^3 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH; and wherein the optionally substituted C_{1-12} alkyl chain is

10 optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR^4 , CONH, $CONR^4$, NHCO, NR^4CO , where each R^4 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional R^4 substituents are each independently selected from OH, OR, NH_2 , NHR^5 , NR^5_2 or $N(OH)R^5$ wherein each R^5 is independently selected from C_{1-4}
15 alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH; and

wherein X represents NH, NMe, CH_2 , SO, SO_2 , or O;

or a pharmacologically acceptable salt thereof.

20

In a ninth aspect of the present invention there is provided a method of making a compound of Formula I defined above including the steps of

- 1 preparing a compound of Formula XVIII as defined above
- 25 2 coupling the compound of Formula XVIII with a DNA targeting agent as defined above to provide a compound of Formula I.

In a tenth aspect of the present invention there is provided a method of making a compound of Formula I' defined above including the steps of

30

- 1 preparing a compound of Formula XVII' as defined above
- 2 coupling the compound of Formula XVII' with a DNA targeting agent as defined above to provide a compound of Formula I'.

It is to be recognised that certain compounds of the present invention may exist in one or more different enantiomeric or diastereomeric forms. It is to be understood that the enantiomeric or diastereomeric forms are included in the above aspects of the invention.

5

The term halo or halogen group used throughout the specification is to be taken as meaning a fluoro, chloro, bromo or iodo group.

10

The term pharmaceutically acceptable salt used throughout the specification is to be taken as meaning any acid or base derived salts formed from hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic, isoethionic acids and the like and potassium carbonate sodium or potassium hydroxide ammonia, triethylamine, triethanolamine and the like.

15

Further aspects of the present invention will become apparent from the following description given by way of example only and with reference to the accompanying synthetic schemes.

20


DETAILED DESCRIPTION OF THE INVENTION

Methods for preparing compounds of Formula I of the invention.

25

3-Chloro-1,2,4-benzotriazine 1-oxide (**3**) was readily synthesised from 2-nitroaniline in 3 steps (50% yield) (Scheme 1). Preparation of the diamine **4** can be achieved as shown in Scheme 2. Coupling of chloride **3** with the monoprotected diamine **4**, readily prepared in 85% yield from the 6-aminohexan-1-ol, gave carbamate **5** as illustrated in Scheme 3. Reaction of **5** with MCPBA in DCM gives 1,4-dioxide **6** in 39% yield and recovered starting material **5** (50%). This represents a departure from known methods (Lee et al, US Patent 5616584, April, 1997) that use trifluoroperacetic acid as the oxidant. Cleavage of the 1,4-dioxide carbamate **6** with HCl in MeOH gave 1,4-dioxide **7** in good yield.


30

Scheme 1

Reagents: (yield %)


a) NH_2CN , HOAc , HCl ;5 b) NaOH ;c) HCl , NaNO_2 , 49% from nitroaniline;d) POCl_3 , PhNMe_2 , 59%**Scheme 2**

10



Reagents:


e) BOC_2O , DCM ;f) MsCl , Et_3N , DCM ;15 g) NaN_3 , DMF .

Scheme 3

Reagents: (yield %)

- 5 i) Et_3N , DCM, 65%;
 j) MCPBA, DCM, 37% + 50% SM;
 k) HCl, MeOH, 85%

An alternative approach to using trifluoroacetic anhydride to provide protection for the primary amine and to generate trifluoroperacetic acid *in situ* was also used (Scheme 4). Deprotection of carbamate 5 gave the amine 8. Reaction of 8 with trifluoroacetic anhydride followed by 30% H_2O_2 gave a mixture of the 1-oxide 9 (22% yield) and 1,4-dioxide 10 (51% yield). 1-Oxide 9 was oxidised with trifluoroperacetic acid to give 10 (29% yield) as well as starting material 9 (61% yield). Deprotection of the trifluoroacetamide 10 provided 1,4-dioxide 7 in good yield.


Scheme 4

Reagents: (yield %)

- a) HCl, MeOH, 87%;
- 5 b) $(\text{CF}_3\text{CO})_2\text{O}$, 35% H_2O_2 , DCM, 51% + 10 (22%);
- c) $\text{CF}_3\text{CO}_3\text{H}$, DCM, 29% + SM (61%);
- d) NaOH, MeOH, 83%.

Coupling of 1,4 dioxide 7 with 9-methoxyacridine (Albert, "The Acridines" 2nd ed.


- 10 1966, Edward Arnold, London, p. 281) provided a compound of Formula I: the aminoacridine derivative 11 (Scheme 5). Similarly, reaction of 7 with 4-(1*H*-imidazol-1-ylcarbonyl)acridine (Spicer et al., *Anti-Cancer Drug Des.*, **1999**, 14, 281-289) gave 12, a compound of Formula I. Similarly, reaction of the imidazolidine of quinoline 4-acetic acid gave 13, a compound of Formula I.

Scheme 5

Reagents: (yield %)

- 5 a) 9-methoxyacridine, MeOH, 60%;
- b) acridine 4-carboxylic acid, CDI, DMF; 7, THF, 91%;
- c) quinoline 4-carboxylic acid, CDI, DMF, 80%; 7, DMF/THF.

Reaction of chloride 3 with tert-butyl 3-aminopropylcarbamate gives 14, which was
 10 oxidised to 1,4-dioxide 15 with MCPBA (Scheme 6). Deprotection of 15 under acid
 conditions gave amine 16 which was reacted with 4-(1*H*-imidazol-1-ylcarbonyl)acridine
 to give 17, a compound of Formula I.

Scheme 6

5 Reagents: (yield %)

a) Et₃N, DCM, 74%;

b) MCPBA, DCM, 24% + 45% SM ;


c) HCl, MeOH, 80%;

d) acridine 4-carboxylic acid, CDI, DMF; **16**, DCM, 80%.

10

Coupling of chloride **3** with 2-(aminoethoxy)ethanol gave alcohol **18** in 63% yield which was converted to the mesylate and displaced with sodium azide to give azide **19** in 89% yield (Scheme 7). Selective reduction of the azide group rather than 1-oxide of **19** could not be effected by hydrogenation using palladium on charcoal or Lindlar catalyst (Rolla et al., *J. Org. Chem.*, **1965**, 47, 4322-432). Other methods for reducing azides such as NaBH₄ under PTC (Corey et al., *Synth.*, **1975**, 590-591), BH₃.DMS (Hassner & Levy, *J. Amer. Chem. Soc.*, **1965**, 87, 4203-4204) or Staudinger conditions using P(OEt)₃ (Koziara & Zwierzak, *Synth.*, **1992**, 1063-1065) were ineffective.


However, treatment of azide **19** with propane-1,3-dithiol and Et₃N in refluxing methanol (Bayley et al., *Tet. Lett.*, **1978**, 39, 3633-3634) provided the intermediate amine which was protected without isolation with di-*tert*-butyldicarbonate to give carbamate **20** in 93% yield for the two steps. Oxidation of **20** with MCPBA gave 1,4-dioxide **21** in 40% yield as well as recovered starting material (50%). Deprotection of **21** with trifluoroacetic acid gave amine **22** in 91% yield. Coupling of **22** with 4-(1H-imidazol-1-ylcarbonyl)acridine gave compound **23** in 97% yield.

Scheme 7

Reagents: (yield %)

- 5 a) Et_3N , DCM, 63%;
- b) MsCl , Et_3N , DCM;
- c) NaN_3 , DMF, 89% from **24**;
- d) propane-1,3-dithiol, Et_3N , MeOH;
- e) BOC_2O , THF, 93% from **25**;
- 10 f) MCPBA, NaHCO_3 , DCM, 40% + 50% SM;
- g) $\text{CF}_3\text{CO}_2\text{H}$, DCM, 91%;
- h) acridine 4-carboxylic acid, CDI, DMF; **28**, THF, 97%.


Similarly, reaction of **22** with the imidazolides of 8-quinolinecarboxylic acid, 2-phenyl-1H-benzimidazole-4-carboxylic acid (Denny et al., *J. Med. Chem.* **1990**, 33, 814-819) and 2-(4-pyridinyl)-8-quinolinecarboxylic acid (Atwell et al., *J. Med. Chem.* **1989**, 32, 396-401) gave compounds of Formula I: **24**, **25**, and **26** respectively (Scheme 8).

Scheme 8**Reagents:**

- a) quinoline 8-carboxylic acid, CDI, DMF; **22**, DCM, 84%;
- 5 b) 2-phenylbenzimidazole 4-carboxylic acid, CDI, DMF; **22**, DCM, 86%;
- c) 2-pyridylquinoline 8-carboxylic acid, CDI, DMF; **22**, DCM, 70%.

Reaction of chloride **3** with *tert*-butyl bis(3-aminopropyl)carbamate and protection of the intermediate primary amine with trifluoroacetic anhydride gave the trifluoroacetamide **27** in 39% for the two steps (Scheme 9). Oxidation of **27** with MCPBA gave the 1,4-dioxide **28** (8% with 65% recovered starting material).

Deprotection of **28** gave amine **29** in good yield which was coupled to 4-(1*H*-imidazol-1-ylcarbonyl)acridine to give compound **30**, a compound of Formula I.


Scheme 9

Reagents: (yield %)

- 5 a) Et₃N, DCM;
- b) (CF₃CO)₂O, DCM, 22% from **3**;
- c) MCPBA, NaHCO₃, DCM, 8% + 65% SM;
- d) K₂CO₃, MeOH, H₂O, 74%;
- e) acridine 4-carboxylic acid, CDI, DMF; **30**, DCM, 67%; HCl, MeOH, 90%.


10

Similarly, reaction of amine **29** with imidazolides of phenazine and 9-methylphenazine followed by deprotection under acidic conditions gave compounds **31** and **32**, respectively (Scheme 10).

Scheme 10

Reagents: (yield %)

- 5 a) phenazine 1-carboxylic acid, CDI, DMF; **29**, DCM, 40%.
 b) HCl, MeOH, 85%.
 c) 9-methylphenazine 1-carboxylic acid, CDI, DMF; **29**, DCM, 40%.
 d) HCl, MeOH, 86%.
- 10 Reaction of chloride **3** with amine **33**, prepared from *N*¹-(2-aminoethyl)-1,2-ethanediamine gave the 1-oxide **34** (Scheme 11). Compound **34** was protected as carbamate **35** and oxidized with MCPBA to give dioxide **36**. Deprotection and coupling of the intermediate amine with the imidazolide of acridine 4-carboxylic acid gave compound **37**, a compound of Formula I.

Scheme 11

Reagents: (yield %)

a) $\text{CF}_3\text{CO}_2\text{Et}$, ether, 61%;

5 b) $(\text{BOC})_2\text{O}$, quant;

c) aq. NH_3 , MeOH, quant;

d) **3**, Et_3N , DME, 72%;

e) $(\text{BOC})_2\text{O}$, DCM, 52%;


f) MCPBA, DCM, 39% + 62% SM;

10 g) HCl , MeOH, 76%;

h) acridine-4-carboxylic acid, CDI, DMF, 99%.

Reaction of chloride **3** with N^1 -(3-aminopropyl)- N^1 -methyl-1,3-propanediamine and protection of the intermediate amine gave acetamide **38** in 43% yield (Scheme 12).

15 Oxidation of **38** with trifluoroperacetic acid under acidic conditions resulted in selective aromatic N-oxidation to give 1,4-dioxide **39** (27%) and recovered starting material **38** (24%). Deprotection of **39** gave amine **40** which was coupled with 4-(1*H*-imidazol-1-ylcarbonyl)acridine to give compound **41**, a compound of Formula I, in 66% yield.


Scheme 12

Reagents: (yield %)

- 5 a) Et₃N, DCM;
 b) (CF₃CO)₂O, DCM, 43% from **3**;
 c) MCPBA, NaHCO₃, DCM, 27% + 24% SM;
 d) NH₄OH, MeOH, quant.;
 e) acridine 4-carboxylic acid, CDI, DMF; **40**, DCM, 66%.

10

Similarly, reaction of **40** with the imidazolides of 8-quinolinecarboxylic acid, 2-(4-pyridinyl)-8-quinolinecarboxylic acid (Atwell et al., *J. Med. Chem.* **1989**, 32, 396-401), 5-methyl-4-acridine carboxylic acid, and 9-methyl-4-phenazinecarboxylic acid gave compounds **42**, **43**, **44**, and **45** respectively (Scheme 13).


Scheme 13

Reagents: (yield %)

- 5 a) quinoline 8-carboxylic acid, CDI, DMF; **40**, DCM, 91%;
- b) 2-pyridylquinoline 8-carboxylic acid, CDI, DMF; **40**, DCM, 94%;
- c) 5-methylacridine-4-carboxylic acid, CDI, DMF; **40**, DCM, 88%;
- d) 9-methylphenazine-4-carboxylic acid, CDI, DMF; **40**, DCM, 90%.

- 10 Reaction of 4-amino-3-nitrophenol with cyanamide under acidic conditions followed by condensation under basic conditions gave the phenol **46** (Friebe et. al. US Patent 5,856,325, Jan 5, 1999), which was alkylated under basic conditions to give ether **47** (Scheme 14). Diazotization of **47** gave **48**, which was chlorinated with POCl₃ to give chloride **49**. Coupling of chloride **49** with amine **50** gave the 1-oxide **51**. Protection of
- 15 **51** as the trifluoroacetamide **52** and oxidation with trifluoroperacetic acid gave the dioxide **53**. Deprotection of **53** gave intermediate amine **54**, which was coupled with the imidazolidine of acridine-4-carboxylic acid to give compound **55**, a compound of Formula I.

Scheme 14




Reagents: (yield %)

- a) NH_2CN , HCl ; NaOH , 97%;
- 5 b) $\text{MeOCH}_2\text{CH}_2\text{Br}$, K_2CO_3 , DMF, 77%;
- c) NaNO_2 , HCl , 68%;
- d) POCl_3 , 83%;
- e) Et_3N , DME, 98%;
- f) $\text{CF}_3\text{CO}_2\text{Et}$, H_2O , MeCN , 87%;
- 10 g) $\text{CF}_3\text{CO}_3\text{H}$, DCM, 30%;
- h) aq. NH_3 , MeOH ;
- i) acridine-4-carboxylic acid, CDI, DMF; 54, THF, 79% (two steps).

Reaction of chloride **3** with allyltributyltin in the presence of tetrakis-
 15 palladiumtriphenylphosphine in DME at reflux temperature gave alkene **56** in high yield
 (Scheme 15). Hydroboration of **56** gave the alcohol **57** which was activated with

methanesulfonyl chloride and reacted with *tert*-butyl 3-aminopropylcarbamate to give the amine **58**. Conversion to the trifluoroacetamide **59** and oxidation with trifluoroperacetic acid gave the 1,4-dioxide **60**, which was deprotected under basic conditions to give amine **61**. Coupling of amine **61** with the imidazolidine of acridine-4-carboxylic acid gave compound **62**, a compound of Formula I. Similarly, coupling of amine **61** with the imidazolidine of phenazine-4-carboxylic acid gave compound **63**, a compound of Formula I.

Scheme 15

Reagents: (yield %)

a) allylSnBu₃, Pd(PPh₃)₄, DME, 93%;

b) 9-BBN, THF; 30% H₂O₂, 3 M NaOH, 87%;

c) MsCl, Et₃N, DCM; *tert*-butyl 3-aminopropylcarbamate, DMF, 48%;

d) HCl, MeOH; CF₃CO₂Et, H₂O, MeCN, 92%;

e) CF₃CO₃H, CF₃CO₂H, CHCl₃, 57%;

f) aq. NH₃, MeOH;


g) acridine-4-carboxylic acid, CDI, DMF; **61**, THF, 40% (two steps);

h) phenazine-4-carboxylic acid, CDI, DMF; **61**, THF, 40% (two steps);

h) phenazine-1-carboxylic acid, CDI, DMF; **61**, THF, 56% (two steps).

Reaction of amine **16** with the imidazolide of *N*-(7-chloro-4-quinoliny)- β -alanine (**64**) (Titus et al, *J. Org. Chem.*, **1948**, *13*, 39-62) gave amide **65**, a compound of Formula I


Scheme 16



Reagents: (yield %)

10 a) *N*-(7-chloro-4-quinoliny)- β -alanine, CDI, DMF; **16**, DMF, 46%.

Scheme 17



15 Reagents: (yield %)

a) NaNO₂, trifluoroacetic acid, 100%;

b) POCl₃, 60%;

c) **68** + **69**, Et₃N, DME, 93%;

d) HCl, MeOH, 100%;


20 e) CF₃CO₂Et, H₂O, MeCN, 92%;

f) CF₃CO₃H, trifluoroacetic acid, DCM, 35% + 40% SM;

- g) aqueous NH_3 , MeOH;
 h) Acridine-4-carboxylic acid, CDI, DMF, 100%.

Similarly, deprotection of **73** and reaction with the imidazolides of 2-(4-pyridinyl)-8-quinolinecarboxylic acid (Atwell et al., *J. Med. Chem.* **1989**, 32, 396-401), phenazine-1-carboxylic acid (Rewcastle et al., *J. Med. Chem.* **1987**, 30, 843-851) and 9-methylphenazine-1-carboxylic acid (Rewcastle et al., *J. Med. Chem.* **1987**, 30, 843-851) gave compounds of Formula I: **75**, **76**, and **77** respectively (Scheme 18).

Scheme 18




Reagents: (yield %)

- a) aqueous NH_3 , MeOH;
 b) 2-(4-pyridyl)quinoline-8-carboxylic acid, CDI, DMF, 89%;
 c) phenazine-1-carboxylic acid, CDI, DMF, 100%;
 d) 9-methylphenazine-1-carboxylic acid, CDI, DMF, 91%.

Diazotization of amine **78** [Hay et al., *J. Med. Chem.* **2003**, 46, 169-182] with sodium nitrite in trifluoroacetic acid gave the alcohol **79** (Scheme 19) which was converted to chloride **80** in POCl_3 . Coupling of chloride **80** with the mono-protected amine **69** gave carbamate **81** which was deprotected under acidic conditions to give amine **82** which was reprotected as the trifluoroacetate **83**. Oxidation of **83** with trifluoroperacetic acid

gave 1,4-dioxide **84** which was deprotected under basic conditions and coupled to the imidazolidine of acridine-4-carboxylic acid (Spicer et al., *Anti-Cancer Drug Des.*, **1999**, 14, 281-289) to give compound **85**.


5 Scheme 19



Reagents: (yield %)

- 10 a) NaNO₂, trifluoroacetic acid, 97%;
- b) POCl₃, 79%;
- c) **69**, Et₃N, DME, 80%;
- d) HCl, MeOH, 99%;
- e) CF₃CO₂Et, H₂O, MeCN, 100%;
- 15 f) CF₃CO₃H, trifluoroacetic acid, DCM, 30% + 49% SM;
- g) aqueous NH₃, MeOH;
- h) acridine-4-carboxylic acid, CDI, DMF, 94%.

- Similarly, deprotection of **84** and reaction with the imidazolidines of 2-(4-pyridinyl)-8-quinolinecarboxylic acid, phenazine-1-carboxylic acid and 9-methylphenazine-1-carboxylic acid gave compounds of Formula I: **86**, **87**, and **88** respectively (Scheme 20).


Scheme 20**5 Reagents: (yield %)**

- a) aqueous NH_3 , MeOH;
- b) 2-(4-pyridyl)quinoline-8-carboxylic acid, CDI, DMF, 100%;
- c) phenazine-1-carboxylic acid, CDI, DMF, 98%;
- d) 9-methylphenazine-1-carboxylic acid, CDI, DMF, 91%.

10


Coupling of chloride **3** with the mono-protected amine **90**, prepared from **89**, gave carbamate **91** which was deprotected under acidic conditions to give amine **92** which was reprotected as the trifluoroacetate **93** (Scheme 21). Oxidation of **93** with trifluoroperacetic acid gave 1,4-dioxide **94** which was deprotected under basic conditions and coupled to the imidazolide of acridine-4-carboxylic acid to give compound **95**.

15

Scheme 21

- 5 Reagents: (yield %)
- a) BOC_2O , THF, 46%;
 - b) **3** + **90**, Et_3N , DME, 52% + 25% SM;
 - c) HCl , MeOH , 100%;
 - d) $\text{CF}_3\text{CO}_2\text{Et}$, H_2O , MeCN , 88%;
 - 10 e) $\text{CF}_3\text{CO}_2\text{H}$, trifluoroacetic acid, DCM, 47% + 6% SM;
 - f) aqueous NH_3 , MeOH ;
 - g) acridine-4-carboxylic acid, CDI, DMF, 94%.


- Similarly, deprotection of **94** and reaction with the imidazolides of 2-(4-pyridinyl)-8-quinolinecarboxylic acid, phenazine-1-carboxylic acid, 9-methylphenazine-1-carboxylic acid and 5-methylacridine-4-carboxylic acid gave compounds **96**, **97**, **98**, and **99** respectively (Scheme 22).
- 15

Scheme 22**5 Reagents: (yield %)**

- a) aqueous NH_3 , MeOH;
- b) 2-(4-pyridyl)quinoline-8-carboxylic acid, CDI, DMF, 97%;
- c) phenazine-1-carboxylic acid, CDI, DMF, 88%;
- d) 9-methylphenazine-1-carboxylic acid, CDI, DMF, 80%;
- e) 5-methylacridine-4-carboxylic acid, CDI, DMF, 100%.

10

Deprotection of trifluoroacetamide **39** under basic conditions and reaction with the imidazolide of phenazine-1-carboxylic acid gave compounds **100** (Scheme 23).

Scheme 23


5 Reagents: (yield %)

a) aqueous NH_3 , MeOH;

b) phenazine-1-carboxylic acid, CDI, DMF, 82%.

Deprotection of **36** and reaction with the imidazolides of 2-(4-pyridinyl)-8-

10 quinolinecarboxylic acid and 5-methylacridine-4-carboxylic acid gave compounds **102** and **103** respectively (Scheme 24).

Scheme 24

15 Reagents: (yield %)

a) HCl/MeOH, 76%;


b) 2-(4-pyridyl)quinoline-8-carboxylic acid, CDI, DMF, 75%;

c) 5-methylacridine-4-carboxylic acid, CDI, DMF, 75%.

20

Coupling of acid **104** (Baird & Dervan, *J. Am. Chem. Soc.* **1996**, 118, 6141–6146) and amine **105** (Baird & Dervan, *J. Am. Chem. Soc.* **1996**, 118, 6141–6146) with EDCI and DMAP gave ester **106** (Scheme 25) which was hydrolysed under basic conditions to give acid **107**. Coupling of acid **107** and 3-dimethylaminopropylamine with EDCI and DMAP gave amide **108**. Reaction of chloride **3** with methyl 4-aminobutanoate gave ester **109** which was oxidised with trifluoroacetic acid to give 1,4-dioxide **110** which was hydrolysed to acid **111**. Deprotection of carbamate **108** followed by coupling to acid **111** with EDCI and DMAP gave compound **112**.

10 Scheme 25




Reagents: (yield %)

- 15 a) **104** + **105**, EDCI, DMAP, DMF, DCM, 64%;
 b) LiOH, THF, MeOH, 94%;
 c) $\text{NH}_2(\text{CH}_2)_3\text{NMe}_2$, EDCI, DMAP, DMF, 76%;
 d) $\text{NH}_2(\text{CH}_2)_3\text{CO}_2\text{Me}$, Et_3N , DME, 81%;
 e) $\text{CF}_3\text{CO}_3\text{H}$, DCM, 33%;
 20 f) NaOH, MeOH, 81%;

- g) HCl/MeOH;
 h) **111**, EDCI, DMAP, DMF, DCM, 9%.

Reaction of the phenol **46** with the protected bromide gave compound **113** (Scheme 26)
 5 which underwent diazotization to the 3-hydroxy intermediate and chlorination with
 POCl₃ to give chloride **114**. Stille coupling with tetraethyltin in the presence of a
 palladium catalyst gave the 3-ethyl compound **115**. Deprotection followed by
 reprotection with dibutyldicarbonate gave compound **116** which was oxidised with
 MCPBA to give dioxide **117**. Deprotection and coupling of the imidazolide of acridine-
 10 4-carboxylic acid is expected to afford compound **118**, a compound of Formula I'.


Scheme 26



- 15 Reagents: (yield %)
 a) CF₃CONHCH₂CH₂Br, K₂CO₃, DMF, 66%;
 b) NaNO₂, CF₃CO₂H, quant.;
 c) POCl₃, quant.;
 d) Et₄Sn, Pd(PPh₃)₄, DME, 79%;
 20 e) aq. NH₃, MeOH, 80%;
 f) BOC₂O, THF, 88%;
 g) MCPBA, DCM, 76%;
 h) aq. HCl, MeOH;
 i) acridine-4-carboxylic acid, CDI, DMF.

Examples of the compounds of the invention

Table 1 gives details on examples of compounds within the scope of the invention, and preparable by the methods of the invention.



I

Table 1

Table 1. Examples of compounds

No.	#	Y ₁	Y ₂	X	A	DNA targeting unit	mp (°C)	Anal
11	3	H	H	NH	(CH ₂) ₆	9-NHAcridine	118-119	C ₃ H ₃ N
12	3	H	H	NH	(CH ₂) ₆	4-NHCOacridine	196-198	C ₃ H ₃ N
13	3	H	H	NH	(CH ₂) ₆	4-NHquinoline	196-198	C ₃ H ₃ N
17	3	H	H	NH	(CH ₂) ₃	4-NHCOacridine	192	C ₃ H ₃ N
23	3	H	H	NH	(CH ₂) ₂ O(CH ₂) ₂	4-NHCOacridine	98-100	C ₃ H ₃ N
24	3	H	H	NH	(CH ₂) ₂ O(CH ₂) ₂	8-NHCOquinoline	168-170	C ₃ H ₃ N
25	3	H	H	NH	(CH ₂) ₂ O(CH ₂) ₂	4-NHCObenz-imidazole-2-phenyl	203-207	C ₃ H ₃ N
26	3	H	H	NH	(CH ₂) ₂ O(CH ₂) ₂	8-NHCOquinoline-2-(4-pyridyl)	128-130	C ₃ H ₃ N
30	3	H	H	NH	(CH ₂) ₃ NH(CH ₂) ₃	4-NHCOacridine	gum	C ₃ H ₃ N
31	3	H	H	NH	(CH ₂) ₃ NH(CH ₂) ₃	1-NHCOphenazine	163-169	C ₃ H ₃ N
32	3	H	H	NH	(CH ₂) ₃ NH(CH ₂) ₃	1-NHCO-9-methyl-phenazine	183-186	HRMS
37	3	H	H	NH	(CH ₂) ₂ NH(CH ₂) ₂	4-NHCOacridine	151-154	C ₃ H ₃ N
41	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	4-NHCOacridine	169-171	HRMS
42	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	8-NHCOquinoline	119-121	C ₃ H ₃ N
43	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	8-NHCO-2-(4-pyridyl)quinoline	179-181	C ₃ H ₃ N
44	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	4-NHCO-5-methylacridine	158-162	C ₃ H ₃ N
45	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCO-9-methylphenazine	138-142	C ₃ H ₃ N
55	3	†	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	4-NHCOacridine	98-103	HRMS
62	3	H	H	CH ₂	(CH ₂) ₂ NMe(CH ₂) ₃	4-NHCOacridine	gum	HRMS
63	3	H	H	CH ₂	(CH ₂) ₂ NMe(CH ₂) ₃	1-NHCOphenazine	173	HRMS

65	3	H	H	NH	(CH ₂) ₃ NHCO(CH ₂) ₂	4-NH-7-Clquinoline	202	HRMS
74	3	7-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	4-NHCOacridine	166-168	C ₃ H ₃ N
75	3	7-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	8-NHCO-2-(4-pyridyl)quinoline	178-180	C ₃ H ₃ N
76	3	7-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCOphenazine	118-122	C ₃ H ₃ N
77	3	7-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCO-9-methyl-phenazine	119-122	C ₃ H ₃ N
85	3	6-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	4-NHCOacridine	158-160	C ₃ H ₃ N
86	3	6-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	8-NHCO-2-(4-pyridyl)quinoline	178-180	C ₃ H ₃ N
87	3	6-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCOphenazine	111-114	C ₃ H ₃ N
88	3	6-Me	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCO-9-methyl-phenazine	80-83	HRMS
95	3	H	H	NH	(CH ₂) ₂ NMe(CH ₂) ₂	4-NHCOacridine	160-162	HRMS
96	3	H	H	NH	(CH ₂) ₂ NMe(CH ₂) ₂	8-NHCO-2-(4-pyridyl)quinoline	130-135	C ₃ H ₃ N
97	3	H	H	NH	(CH ₂) ₂ NMe(CH ₂) ₂	1-NHCOphenazine	163-165	C ₃ H ₃ N
98	3	H	H	NH	(CH ₂) ₂ NMe(CH ₂) ₂	1-NHCO-9-methyl-phenazine	161-163	C ₃ H ₃ N
99	3	H	H	NH	(CH ₂) ₂ NMe(CH ₂) ₂	4-NHCO-5-methylacridine	148-152	C ₃ H ₃ N
100	3	H	H	NH	(CH ₂) ₃ NMe(CH ₂) ₃	1-NHCOphenazine	129-130	C ₃ H ₃ N
102	3	H	H	NH	(CH ₂) ₂ NH(CH ₂) ₂	8-NHCO-2-(4-pyridyl)quinoline	160-165	C ₃ H ₃ N
103	3	H	H	NH	(CH ₂) ₂ NH(CH ₂) ₂	4-NHCO-5-methylacridine	135-140	HRMS
112	3	H	H	NH	(CH ₂) ₃ CONH	3-pyr-5-CONH-3-pyr-5-CONH(CH ₂) ₃ NMe ₂	140-145	C ₃ H ₃ N

Side chain position

[†]7-MeOCH₂CH₂O-

In the following examples representative of the invention and the detailed methods for preparing them:

Elemental analyses were carried out in the Microchemical Laboratory, University of Otago, Dunedin, NZ.

5 Melting points were determined on an Electrothermal 2300 Melting Point Apparatus. IR spectra were recorded on a Midac FT-IR as KBr discs, unless otherwise stated.

NMR spectra were obtained on a Bruker Avance-400 spectrometer at 400 MHz for ^1H and 100 MHz for ^{13}C spectra. Spectra were obtained in CDCl_3 unless otherwise specified, and are referenced to Me_4Si . Chemical shifts and coupling constants were

10 recorded in units of ppm and Hz, respectively. Assignments were determined using COSY, HSQC, and HMBC two-dimensional experiments.

Mass spectra were determined on a VG-70SE mass spectrometer using an ionizing potential of 70 eV at a nominal resolution of 1000. High-resolution spectra were obtained at nominal resolutions of 3000, 5000, or 10000 as appropriate. All spectra were

15 obtained as electron impact (EI) using PFK as the reference unless otherwise stated.

Solutions in organic solvents were dried with anhydrous Na_2SO_4 . Solvents were evaporated under reduced pressure on a rotary evaporator.

Thin-layer chromatography was carried out on aluminium-backed silica gel plates (Merck 60 F₂₅₄) with visualization of components by UV light (254 nm) or exposure to

20 I_2 .

Column chromatography was carried out on silica gel, (Merck 230–400 mesh). All compounds designated for testing were analyzed at >99% purity by reverse phase HPLC using an Agilent 1100 liquid chromatograph, an Alltima C₁₈ (5 μ) stainless steel column (150 mm \times 3.2 mm i.d.) and an Agilent 1100 diode array detector. Chromatograms were

25 run using various gradients of aqueous (0.045 M ammonium formate and formic acid at pH 3.5) and organic (80% MeCN/MilliQ water) phases. DCM refers to

dichloromethane; DME refers to dimethoxyethane, DMF refers to dry dimethyl

formamide; ether refers to diethyl ether; EtOAc refers to ethyl acetate; EtOH refers to

ethanol; MeOH refers to methanol; pet. ether refers to petroleum ether, boiling range 40–

30 60 °C; THF refers to tetrahydrofuran dried over sodium benzophenone ketyl. All solvents were freshly distilled.

Example A.**3-[(6-Aminohexyl)amino]-1,2,4-benzotriazine 1,4-dioxide (7).****3-Chloro-1,2,4-benzotriazine 1-oxide (3).** 2-Nitroaniline (10 g, 72.4 mmol) and

5 cyanamide (14.0 g, 330 mmol) were melted together and cHCl (20 mL) added cautiously. The mixture was heated at 100 °C until the foaming subsided. The mixture was made strongly alkaline with 30% w/v NaOH and heated at 100 °C for 10 min.

The suspension was cooled to 25 °C and the yellow solid filtered, washed with water (20 mL) and dried. A small sample was recrystallized to give 3-amino-1,2,4-

10 benzotriazine 1-oxide (1) mp (MeOH/EtOAc) 267-269 °C; lit. [Arndt, *Ber.* **1913**, 46, 3522-3529] mp (EtOH) 269 °C]. The remainder was dissolved in 2 M HCl (300 mL), cooled to 5 °C, and a solution of NaNO₂ (10 g, 0.145 mol) in water (100 mL) added dropwise. The resulting precipitate was filtered, dissolved in dilute NH₃, filtered, and acidified with cHCl. The precipitate was filtered, washed with water and dried to give
15 3-hydroxy-1,2,4-benzotriazine 1-oxide (2) (5.77 g, 49%) as a yellow powder, mp 209-212 °C; lit. [Arndt, *Ber.* **1913**, 46, 3522-3529] mp (H₂O) 219 °C]; ¹H NMR

[(CD₃)₂SO] δ 8.14 (d, *J* = 8.4 Hz, 1 H, H-8), 7.77-7.81 (m, 1 H, H-6), 7.54 (d, *J* = 8.4 Hz, 1 H, H-5), 7.88-7.92 (m, 3 H, H-7, NH₂); ¹³C NMR [(CD₃)₂SO] δ 160.2, 148.7, 135.6, 129.8, 125.8, 124.6, 119.8. A mixture of the alcohol (2) (5.7 g, 34.9 mmol),

20 *N,N*-dimethylaniline (11 mL, 87.3 mmol), and POCl₃ (23 mL, 244 mmol) was heated at reflux temperature for 1 h then poured on to ice. The resulting solid was filtered and recrystallized to give 3-chloro-1,2,4-benzotriazine 1-oxide (3) (3.77 g, 59%) as a pale yellow powder, mp 119-119.5 °C; lit. [Robbins *et al.*, *J. Chem. Soc.*, **1957**, 3186-3194] (MeOH) 117-118 °C]; ¹H NMR [(CD₃)₂SO] δ 8.38 (dd, *J* = 8.7, 1.0 Hz, 1 H, H-8), 8.16 (ddd, *J* = 8.3, 7.0, 1.3 Hz, 1 H, H-6), 8.06 (dd, *J* = 8.2, 1.0 Hz, 1 H, H-5), 7.90 (ddd, *J* = 8.7, 6.9, 1.3 Hz, 1 H, H-7); ¹³C NMR [(CD₃)₂SO] δ 155.3, 146.9, 137.2, 133.9, 131.5, 128.0, 119.9.

6-*t*-Butyloxycarbamoylhexylamine (4). A solution of di-*t*-butyldicarbonate (18.6 g,

30 85.3 mmol) in dry DCM (100 mL) was added dropwise to a stirred solution of 6-aminohexanol (10.0 g, 85.3 mmol) in dry DCM (100 mL) at 20 °C and stirred for 16 h. The solution was washed with dilute aqueous Na₂CO₃ solution (100 mL), 0.1 M HCl (100 mL), water (100 mL), brine (50 mL), dried and the solvent evaporated. The

residue was dissolved in DCM (250 mL) and Et₃N (15.5 mL, 111 mmol) added. A solution of methanesulfonyl chloride (7.3 mL, 94 mmol) was added dropwise and the mixture stirred at 20 °C for 16 h. The solution was washed with saturated aqueous KHCO₃ (100 mL), water (2 × 100 mL), brine (50 mL), dried, and the solvent
5 evaporated. The residue was dissolved in DMF (100 mL) and NaN₃ (5.55 g, 85.3 mmol) added. The mixture was stirred at 100 °C for 1 h, the solvent evaporated and the residue partitioned between EtOAc (200 mL) and water (200 mL). The organic fraction was washed with water (200 mL), brine (100 mL), dried and the solvent evaporated. The residue was purified by chromatography, eluting with 30%
10 EtOAc/pet. ether, to give the 6-*t*-butyloxycarbamoylhexyl azide (17.5 g, 85%) as a colorless oil, ¹H NMR δ 4.53 (br s, 1 H, OCONH), 3.52 (t, *J* = 6.9 Hz, 2 H, CH₂N), 3.11 (dt, *J* = 6.5, 6.4 Hz, 2 H, CH₂N), 1.57–1.63 (m, 2 H, CH₂), 1.44–1.52 (m, 11 H, CH₂, C(CH₃)₃), 1.30–1.40 (m, 4 H, 2 × CH₂); ¹³C NMR δ 156.0, 79.1, 51.3, 40.4, 29.9, 28.7, 28.4 (3), 26.4, 26.3. A mixture of azide (14.81 g, 61.1 mol) and Pd/C (0.5
15 g) in EtOAc/EtOH (200 mL) was stirred at 20 °C under hydrogen (60 psi) for 1 h. The mixture was filtered through celite, the cake washed with EtOAc (3 × 30 mL) and the solvent evaporated to give **4** (12.82 g, 97%) as a white solid, mp (EtOAc) 89–91 °C; ¹H NMR δ 4.65 (br s, 1 H, OCONH), 3.52 (br s, 2 H, NH₂), 2.69 (t, *J* = 6.9 Hz, 2 H, CH₂N), 1.88 (br s, 2 H, CH₂N), 1.44–1.50 [m, 13 H, 2 × CH₂, C(CH₃)₃], 1.29–1.35
20 (m, 4 H, 2 × CH₂); ¹³C NMR δ 156.0, 78.9, 41.9, 40.4, 33.4, 29.9, 28.3 (3), 26.5, 26.4.

3-[(6-*t*-Butyloxycarbamoylhexyl)amino]-1,2,4-benzotriazine 1-oxide (5). A

solution of amine **4** (12.8 g, 61.1 mmol) in DCM was added to a stirred solution of chloride **3** (3.70 g, 20.4 mmol) and Et₃N (5.7 mL, 40.8 mmol) in DCM (100 mL) and
25 the solution stirred at 20 °C for 96 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (30–100%) of EtOAc/pet. ether, to give 1-oxide **5** (4.77 g, 65%) as a yellow powder, mp (EtOAc/pet. ether) 154–156 °C; ¹H NMR δ 8.26 (d, *J* = 8.6 Hz, 1 H, H-8), 7.70 (dd, *J* = 8.2, 7.2 Hz, 1 H, H-6), 7.59 (d, *J* = 8.5 Hz, 1 H, H-5), 7.27 (dd, *J* = 8.0, 7.5 Hz, 1 H, H-7), 5.34 (br s, 1 H, NH), 4.55
30 (br s, 1 H, OCONH), 3.51 (dd, *J* = 6.8, 6.6 Hz, 2 H, CH₂N), 3.10–3.13 (m, 2 H, CH₂N), 1.64–1.72 (m, 2 H, CH₂), 1.48–1.54 (m, 2 H, CH₂), 1.44 [s, 9 H, C(CH₃)₃], 1.38–1.43 (m, 4 H, 2 × CH₂); ¹³C NMR δ 158.9, 155.5, 148.9, 135.5, 130.8, 126.4,

124.8, 120.4, 79.0, 41.2, 40.3, 30.0, 29.2, 28.4 (3), 26.4, 26.3. Anal. calcd for $C_{18}H_{27}N_5O_3$: C, 59.8; H, 7.5; N, 19.4; found: C, 59.6; H, 7.7; N, 19.2%.

3-[(6-*t*-Butyloxycarbamoyl)hexyl]amino]-1,2,4-benzotriazine 1,4-dioxide (6). A

5 solution of MCPBA (1.48 g, 6.02 mmol) in DCM (20 mL) was added dropwise to a stirred solution of 1-oxide **5** (1.45 g, 4.01 mmol) in DCM (100 mL) at 20 °C and the solution stirred for 4 h. The solution was partitioned between DCM (200 mL) and saturated $KHCO_3$ solution (200 mL). The organic fraction was dried and the solvent evaporated. The residue was purified by chromatography on neutral alumina, eluting
10 with 50% EtOAc/DCM then a gradient (0–10%) MeOH/ $CHCl_3$, to give (i) starting material **5** (0.73 g, 50%), and (ii) 1,4-dioxide **6** (0.55 g, 37%) as a yellow powder, mp (EtOAc/DCM) 132–134 °C; IR (KBr) ν 3367, 3260, 1688, 1622, 1362, 1173 cm^{-1} ; NMR [$(CD_3)_2SO$] δ 8.30 (dd, J = 6.3, 6.1 Hz, 1 H, OCONH), 8.19 (d, J = 8.5 Hz, 1 H, H-8), 8.12 (d, J = 8.5 Hz, 1 H, H-5), 7.91–7.95 (m, 1 H, H-6), 7.53–7.57 (m, 1 H, H-7), 6.76 (br s, 1 H, NH), 3.32–3.39 (m, 2 H, CH_2N), 2.87–2.92 (m, 2 H, CH_2N), 1.56–
15 1.61 (m, 2 H, CH_2), 1.32–1.40 [m, 13 H, 2 \times CH_2 , $C(CH_3)_3$], 1.25–1.31 (m, 2 H, CH_2); ^{13}C NMR [$(CD_3)_2SO$] δ 155.5, 149.7, 138.1, 135.4, 129.8, 126.8, 121.1, 116.8, 77.2, 40.6, 39.8, 29.4, 28.6, 28.2 (3), 25.9, 25.8. Anal. calcd for $C_{18}H_{27}N_5O_4 \cdot \frac{1}{4}H_2O$: C, 56.6; H, 7.3; N, 18.3; found: C, 56.8; H, 7.3; N, 16.8%.

***N*¹-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,6-hexanediamine (7).** HCl gas was

bubbled through a solution of carbamate **6** (204 mg, 0.54 mmol) in MeOH (20 mL) for 2 minutes and the solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue partitioned between $CHCl_3$ (100 mL) and saturated $KHCO_3$ solution
25 (100 mL). The aqueous fraction was further extracted with $CHCl_3$ (3 \times 30 mL), the combined organic extracts dried, and the solvent evaporated to give amine **7** (127 mg, 85%) as a red powder, mp 120–122 °C, IR (KBr) ν 3250, 2926, 1616, 1599, 1410, 1356, 1078 cm^{-1} ; 1H NMR δ 8.34 (d, J = 8.5 Hz, 1 H, H-8'), 8.29 (d, J = 8.6 Hz, 1 H, H-5'), 7.87–7.90 (m, 1 H, H-6'), 7.48–7.52 (m, 1 H, H-7'), 7.13 (s, 1 H, NH), 3.60 (t, J = 7.1 Hz, 2 H, CH_2N), 2.70 (t, J = 6.8 Hz, 2 H, CH_2N), 1.70–1.76 (m, 2 H, CH_2),
30 1.35–1.50 (m, 6 H, 3 \times CH_2); ^{13}C NMR [$(CD_3)_2SO$] δ 149.7, 138.1, 135.4, 129.8, 126.7, 121.0, 116.8, 41.5, 40.6, 33.1, 28.7, 26.1, 26.0. Anal. calcd for $C_{13}H_{19}N_5O_2$: C, 56.3; H, 6.9; N, 25.3; found: C, 56.3; H, 6.8; N, 22.2%. The compound was dissolved

in MeOH, treated with HCl gas, and the solvent evaporated. The residue was crystallized to give the dihydrochloride of **7** as a red powder, mp (MeOH/EtOAc) 150 °C (dec.).

5 ***N*¹-(1-Oxido-1,2,4-benzotriazin-3-yl)-1,6-hexanediamine (8)**. HCl gas was bubbled through a solution of carbamate **5** (1.0 g, 2.77 mmol) in MeOH (80 mL) for 2 minutes and the solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue partitioned between CHCl₃ (100 mL) and Na₂CO₃ solution (100 mL). The aqueous fraction was further extracted with CHCl₃ (3 × 30 mL), the combined organic extracts
0 dried, and the solvent evaporated to give amine **8** (0.63 g, 87%) as a red powder, mp 132–134 °C, ¹H NMR δ 8.25 (dd, *J* = 8.6, 1.0 Hz, 1 H, H-8'), 7.66–7.71 (m, 1 H, H-6'), 7.59 (d, *J* = 8.4 Hz, 1 H, H-5'), 7.26–7.30 (m, 1 H, H-7'), 5.48 (br s, 1 H, NH), 3.52 (dd, *J* = 6.9, 6.3 Hz, 2 H, H-1), 2.69 (dd, *J* = 6.8, 6.6 Hz, 2 H, H-6), 1.64–1.71 (m, 2 H, H-2), 1.35–1.48 (m, 8 H, H-3, H-4, H-5, NH₂); ¹³C NMR δ 159.0, 148.9,
5 135.5, 130.8, 126.4, 124.7, 120.4, 42.0, 41.3, 33.6, 29.3, 26.6, 26.5. Anal. calcd for C₁₃H₁₉N₅O: C, 59.7; H, 7.3; N, 26.8; found: C, 59.5; H, 7.5; N, 26.5%.

Oxidation of *N*¹-(1-oxido-1,2,4-benzotriazin-3-yl)-1,6-hexanediamine (8).

Trifluoroacetic anhydride (11.9 mL) was added to a stirred solution of amine **8** (1.1 g, 4.2 mmol) in DCM (100 mL) and the solution stirred at 20 °C for 30 min. The solution
10 was cooled to 5 °C and 35% H₂O₂ (11.9 mL, ca 105 mmol) added dropwise and the mixture stirred vigorously for 16 h. The mixture was concentrated to 30 mL (CAUTION) and partitioned between DCM (100 mL) and sat. aqueous KHCO₃ solution (50 mL). The aqueous fraction was extracted with DCM (3 × 50 mL), the
15 combined organic fraction dried and the solvent evaporated (CAUTION). The residue was purified by chromatography, eluting with a gradient (0–10%) MeOH/(40–0%) EtOAc/DCM, to give (i) 2,2,2-trifluoro-*N*-{6-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]hexyl}acetamide (**9**) (0.77 g, 51%) as a yellow solid, mp (EtOAc/DCM) 188–189 °C; IR (KBr) ν 3306, 1699, 1588, 1570, 1176 cm⁻¹; ¹H NMR δ 8.27 (dd, *J* =
30 8.7, 1.3 Hz, 1 H, H-8'), 7.70 (ddd, *J* = 8.5, 6.9, 1.3 Hz, 1 H, H-6'), 7.59 (d, *J* = 8.5 Hz, 1 H, H-5'), 7.29 (ddd, *J* = 8.7, 6.9, 1.3 Hz, 1 H, H-7'), 6.33 (br s, 1 H, NH), 5.22 (s, 1 H, CONH), 3.51 (q, *J* = 6.9 Hz, 2 H, H-1), 3.38 (q, *J* = 6.8 Hz, 2 H, H-6), 1.66–1.73 (m, 2 H, H-5), 1.59–1.65 (m, 2 H, H-2), 1.40–1.47 (m, 4 H, H-3, H-4); ¹³C NMR δ

158.7, 156.8 (q, $J = 37$ Hz), 148.6, 135.0, 130.2, 126.0, 124.1, 119.8, 115.7 (q, $J = 288$ Hz), 40.6, 39.2, 28.6, 28.2, 25.9, 25.8. Anal. calcd for $C_{15}H_{18}F_3N_5O_2$: C, 50.4; H, 5.1; N, 19.6; found: C, 50.7; H, 4.9; N, 19.6%, and:

(ii) *N*-{6-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]hexyl}-2,2,2-trifluoroacetamide

- 5 (10) (346 mg, 22%) as a red solid, mp (MeOH/DCM) 163–165 °C; IR (KBr) ν 3437, 3266, 1699, 1634, 1178 cm^{-1} ; 1H NMR $[(CD_3)_2SO]$ δ 9.42 (s, 1 H, NH), 8.31 (t, $J = 6.0$ Hz, 1 H, CONH), 8.20 (d, $J = 8.7$ Hz, 1 H, H-8'), 8.12 (d, $J = 8.6$ Hz, 1 H, H-5'), 7.91–7.95 (m, 1 H, H-6'), 7.53–7.57 (m, 1 H, H-7'), 3.40 (q, $J = 6.7$ Hz, 2 H, H-1), 3.18 (q, $J = 6.6$ Hz, 2 H, H-6), 1.58–1.64 (m, 2 H, H-2), 1.46–1.53 (m, 2 H, H-5),
 10 1.28–1.38 (m, 4 H, H-3, H-4); ^{13}C NMR $[(CD_3)_2SO]$ δ 156.0 (q, $J = 36$ Hz), 149.7, 138.1, 135.4, 129.8, 126.7, 121.1, 116.8, 115.9 (q, $J = 288$ Hz), 40.5, 39.0, 28.5, 28.1, 25.8, 25.7. Anal. calcd for $C_{15}H_{18}F_3N_5O_3$: C, 48.3; H, 4.9; N, 18.8; found: C, 48.5; H, 4.7; N, 18.0%.

- 15 **Oxidation of 2,2,2-trifluoro-*N*-{6-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]hexyl}acetamide (9).** Trifluoroacetic anhydride (4.0 mL, 28.6 mmol) was added dropwise to a stirred suspension of 35% H_2O_2 (2.2 mL, ca. 23 mmol) in DCM (20 mL) at 5 °C and the mixture was stirred for 15 min. The mixture was dried and added to a stirred solution of 1-oxide **9** (409 mg, 1.14 mmol) in DCM (50 mL) and the
 20 solution stirred at 20 °C for 48 h. The solution was partitioned between sat. aqueous $KHCO_3$ (50 mL) and $CHCl_3$ (50 mL). The aqueous fraction was extracted with $CHCl_3$ (3 \times 40 mL), the combined organic fraction dried, and the solvent evaporated (CAUTION). The residue was purified by chromatography, eluting with a gradient (0–10%) MeOH/(40–0%) EtOAc/DCM, to give (i) starting material **9** (250 mg, 61%);
 25 and (ii) 1,4-dioxide **10** (124 mg, 29%), spectroscopically identical to a sample obtained above.

- N*-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)-1,6-hexanediamine (7).** 1 M NaOH solution (2.8 mL, 2.8 mmol) was added to a stirred solution of trifluoroacetamide **10**
 30 (209 mg, 0.56 mmol) in MeOH (20 mL) and the mixture stirred at 20 °C for 16 h. The solvent was evaporated and the residue partitioned between sat. aqueous $KHCO_3$ (70 mL) and $CHCl_3$ (70 mL). The aqueous fraction was extracted with $CHCl_3$ (3 \times 30

mL), the combined organic fraction dried, and the solvent evaporated to give amine 7 (129 mg, 83%), spectroscopically identical with the sample obtained above.

Example B.

5 *N*¹-(9-AcridinyI)-*N*⁶-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,6-hexanediamine (11).

A solution of amine 7 (64 mg, 0.23 mmol) and 9-methoxyacridine (53 mg, 0.25 mmol) in MeOH (10 mL) was stirred at reflux temperature for 10 h. The solvent was evaporated and the residue purified by chromatography on neutral alumina, eluting with a gradient (0–5%) of MeOH/CHCl₃, to give compound 11 (63 mg, 60%) as a red solid, IR (KBr) ν 3293, 1593, 1414, 1362 cm⁻¹; ¹H NMR δ 8.30 (d, *J* = 8.5 Hz, 1 H, H-8''), 8.29 (d, *J* = 8.5 Hz, 1 H, H-5''), 8.11 (d, *J* = 8.6 Hz, 2 H, H-1', H-8'), 8.04 (d, *J* = 8.6 Hz, 2 H, H-4', H-5'), 7.84 (ddd, *J* = 8.5, 7.2, 1.2 Hz, 1 H, H-6''), 7.59–7.64 (m, 2 H, H-3', H-6'), 7.57 (ddd, *J* = 8.5, 7.2, 1.2 Hz, 1 H, H-7''), 7.31–7.35 (m, 2 H, H-2', H-7'), 7.15 (br s, 1 H, NH), 3.84 (dd, *J* = 7.2, 7.1 Hz, 2 H, CH₂N), 3.57 (dt, *J* = 6.7, 6.5 Hz, 2 H, CH₂N), 1.78–1.85 (m, 2 H, CH₂), 1.67–1.74 (m, 2 H, CH₂), 1.43–1.53 (m, 4 H, 2 \times CH₂), NH not observed; ¹³C NMR δ 151.9 (2), 149.8, 148.0, 138.1, 135.8, 130.3, 130.2 (2), 128.1 (2), 127.1, 123.0 (2), 122.9 (2), 121.6, 117.2, 116.1 (2), 50.4, 41.2, 31.4, 29.2, 26.4, 26.3; MS (FAB⁺) *m/z* 455 (MH⁺, 20%), 439 (10); HRMS (FAB⁺) calcd for C₂₆H₂₇N₆O₂ (MH⁺) *m/z* 455.2196, found 455.2182. The compound was dissolved in MeOH and treated with HCl gas and the solvent evaporated. The residue was crystallized from MeOH/EtOAc to give the hydrochloride of 11, mp (MeOH/EtOAc) 118–119 °C. Anal. calcd for C₂₆H₂₆N₆O₂·2HCl·½H₂O: C, 58.2; H, 5.5; N, 15.7; found: C, 57.8; H, 5.5; N, 15.3%.

25 Example C.

N-{6-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]hexyl}-4-acridinecarboxamide (12). A solution of the amine 7 (447 mg, 1.6 mmol) in THF (20 mL) and DMF (10 mL) was added dropwise to a stirred solution of acridine-4-carboxylic acid imidazolide (440 mg, 1.61 mmol) in THF (20 mL) at 5 °C and the solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give compound 12 (708 mg, 91%) as a red solid, mp (EtOAc) 196–198 °C; ¹H NMR δ 11.88 (s, 1 H, NH), 8.99 (dd, *J* = 7.1, 1.5 Hz, 1 H, H-3), 8.90 (s, 1 H, H-9), 8.30 (d, *J* = 8.4 Hz, 1 H,

H-8'), 8.28 (d, $J = 8.7$ Hz, 1 H, H-5'), 8.17 (d, $J = 9.1$ Hz, 1 H, H-5), 8.14 (dd, $J = 8.4$, 1.5 Hz, 1 H, H-1), 8.04 (d, $J = 8.1$ Hz, 1 H, H-8), 7.84–7.91 (m, 2 H, H-6, H-6'), 7.66 (dd, $J = 8.3$, 7.2 Hz, 1 H, H-2), 7.59 (ddd, $J = 7.9$, 7.0, 0.9 Hz, 1 H, H-7), 7.59 (ddd, $J = 8.4$, 7.2, 1.2 Hz, 1 H, H-7'), 7.11 (br dd, $J = 5.7$, 5.5 Hz, 1 H, CONH), 3.71 (dd, $J = 6.9$, 5.6 Hz, 2 H, CH₂N), 3.63 (dd, $J = 6.9$, 6.7 Hz, 2 H, CH₂N), 1.83–1.89 (m, 2 H, CH₂), 1.74–1.81 (m, 2 H, CH₂), 1.55–1.68 (m, 4 H, 2 × CH₂); ¹³C NMR δ 164.6, 149.7, 147.0, 145.5, 145.0, 138.7, 138.1, 135.4, 134.5, 132.8, 132.0, 129.8, 128.5, 128.3, 126.8, 126.5, 126.4, 125.6, 125.3, 121.1, 116.8, 40.6, 39.0, 29.0, 29.6, 26.5, 26.0. Anal. calcd for C₂₇H₂₆N₆O₀·H₂O: C, 64.8; H, 5.6; N, 16.8; found: C, 65.0; H, 5.5; N, 17.1%.

Example D.

***N*-{6-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]hexyl}-4-quinolinecarboxamide (13).** A solution of 4-quinolinecarboxylic acid (308 mg, 1.78 mmol) and CDI (346 mg, 2.13 mmol) in DMF (20 mL) were stirred at 50 °C for 1 h. The solvent was evaporated and the residue recrystallised from DCM/pet. ether to give 4-(1*H*-imidazol-1-ylcarbonyl)quinoline which was used directly without characterisation. A solution of the amine 7 (494 mg, 1.78 mmol) in DMF (10 mL) was added dropwise to a stirred solution of imidazolide in THF (20 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give compound **13** (619 mg, 80%) as a red powder, mp (MeOH/DCM) 196–198 °C; ¹H NMR [(CD₃)₂SO] δ 8.96 (d, $J = 4.3$ Hz, 1 H, H-2), 8.75 (t, $J = 5.5$ Hz, 1 H, CONH), 8.32 (t, $J = 6.1$ Hz, 1 H, NH), 8.20 (d, $J = 8.6$ Hz, 1 H, H-8''), 8.12 (d, $J = 8.6$ Hz, 1 H, H-5''), 8.11 (d, $J = 8.7$ Hz, 1 H, H-5), 8.06 (d, $J = 8.4$ Hz, 1 H, H-8), 7.92 (ddd, $J = 8.4$, 7.1, 1.3 Hz, 1 H, H-6''), 7.80 (ddd, $J = 8.4$, 7.1, 1.0 Hz, 1 H, H-7), 7.66 (ddd, $J = 8.5$, 7.0, 1.0 Hz, 1 H, H-6), 7.55 (ddd, $J = 8.5$, 7.1, 1.3 Hz, 1 H, H-7''), 7.52 (d, $J = 4.4$ Hz, 1 H, H-3), 3.39–3.43 (m, 2 H, H-1'), 3.3–3.35 (m, 2 H, H-6'), 1.65–1.70 (m, 2 H, H-2'), 1.56–1.73 (m, 2 H, H-5'), 1.38–1.45 (m, 4 H, H-3', H-4'); ¹³C NMR [(CD₃)₂SO] δ 166.4, 150.2, 149.7, 147.8, 142.4, 138.1, 135.4, 129.8, 129.6, 129.2, 127.1, 126.7, 125.3, 124.1, 121.0, 118.8, 116.7, 40.6, 38.9, 28.8, 28.6, 26.1, 25.9. Anal. calcd for C₂₃H₂₄N₆O₃: C, 63.9; H, 5.6; N, 19.4; found: C, 63.9; H, 5.4; N, 19.5%.

Example E.

***N*-{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}-4-acridinecarboxamide (17).**

***tert*-Butyl 3-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]propylcarbamate (14).** A

5 solution of chloride **3** (4.0 g, 22.0 mmol), *tert*-butyl 3-aminopropylcarbamate (5.76 g, 33.0 mmol) and Et₃N (4.6 mL, 33.0 mmol) in DCM (150 mL) was stirred at 20 °C for 5 d. The solvent was evaporated, and the residue purified by chromatography, eluting with 20% EtOAc/DCM, to give 1-oxide **14** (5.21 g, 74%) as a yellow powder, mp (EtOAc/DCM) 145–147 °C; ¹H NMR [(CD₃)₂SO] δ 8.13 (dd, *J* = 8.6, 1.1 Hz, 1 H, H-8'), 7.84 (s, 1 H, NH), 7.78 (ddd, *J* = 8.4, 7.1, 1.1 Hz, 1 H, H-6'), 7.56 (d, *J* = 8.4 Hz, 1 H, H-5'), 7.32 (ddd, *J* = 8.6, 7.1, 1.1 Hz, 1 H, H-7'), 6.83 (t, *J* = 5.3 Hz, 1 H, NHCO₂), 3.32–3.36 (m, 2 H, H-1), 2.99–3.04 (m, 2 H, H-3), 1.66–1.73 (m, 2 H, H-2), 1.37 [s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 158.9, 155.6, 148.2, 135.7, 130.0, 125.9, 124.4, 119.9, 77.4, 38.2, 37.5, 28.9, 28.2 (3). Anal. calcd for C₁₅H₂₁N₅O₃: C, 56.4; H, 6.6; N, 21.9; found: C, 56.4; H, 6.6; N, 22.1%.

***tert*-Butyl 3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propylcarbamate (15).** A

10 solution of MCPBA (6.74g, 27.3 mmol) in DCM (80 mL) was added dropwise to a stirred solution of 1-oxide **14** (5.82 g, 18.2 mmol) in DCM (300 mL) and NaHCO₃ (3.1 g, 36.5 mmol). The mixture was stirred at 20 °C for 1 h, partitioned between DCM (400 mL) and sat. aqueous KHCO₃ solution (100 mL). The organic fraction was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–10%) of MeOH/40%EtOAc/DCM, to give (i) starting material **14** (2.63 g, 45%) and (ii) 1,4-dioxide **15** (1.47 g, 24%) as a red solid, mp (EtOAc/MeOH) 134–136 °C; ¹H NMR [(CD₃)₂SO] δ 8.30 (t, *J* = 6.2 Hz, 1 H, NH), 8.20 (d, *J* = 8.5 Hz, 1 H, H-8'), 8.13 (d, *J* = 8.5 Hz, 1 H, H-5'), 7.93 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-6'), 7.57 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-7'), 6.86 (t, *J* = 5.6 Hz, 1 H, NHCO₂), 3.38–3.42 (m, 2 H, H-1), 2.98–3.02 (m, 2 H, H-3), 1.68–1.74 (m, 2 H, H-2), 1.37 [s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 155.6, 149.7, 138.1, 135.4, 129.8, 126.8, 121.0, 116.8, 77.4, 38.2, 37.1, 28.9, 28.1 (3). Anal. calcd for C₁₅H₂₁N₅O₄•¼EtOAc: C, 53.8; H, 6.5; N, 19.6; found: C, 53.5; H, 6.5; N, 19.5%.

***N*¹-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)-1,3-propanediamine (16).** HCl saturated MeOH (20 mL) was added to a solution of carbamate **15** (1.47 mg, 4.38 mmol) in MeOH (30 mL) and the solution stirred at 20 °C for 16 h. The solution was evaporated and the residue dissolved in water (20 mL) the solution neutralized with KHCO₃ and extracted with CHCl₃ (5 × 50 mL). The combined organic fraction was dried and the solvent evaporated to give compound **16** (0.82 g, 80%) as a red solid, mp (MeOH) 121–123 °C; ¹H NMR [(CD₃)₂SO] δ 8.24 (d, *J* = 8.4 Hz, 1 H, H-8'), 8.13 (d, *J* = 8.6 Hz, 1 H, H-5'), 7.99 (ddd, *J* = 8.6, 7.1, 1.0 Hz, 1 H, H-6'), 7.61 (ddd, *J* = 8.4, 7.1, 1.0 Hz, 1 H, H-7'), 4.01 (br s, 3 H, NH, NH₂), 3.48 (t, *J* = 6.7 Hz, 2 H, H-1), 2.66 (t, *J* = 7.0 Hz, 2 H, H-3), 1.73–1.77 (m, 2 H, H-2); MS (FAB⁺) *m/z* 236 (MH⁺, 6%), 220 (10), 204 (5); HRMS calcd for C₁₀H₁₄N₅O₂ (MH⁺) *m/z* 236.1148, found 236.1139.

***N*-{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}-4-acridinecarboxamide**

(17). A solution of amine **16** (128 mg, 0.54 mmol) in DCM (5 mL) was added dropwise to a stirred solution of 4-(1*H*-imidazol-1-ylcarbonyl)acridine (156 mg, 0.57 mmol) in DCM (10 mL) at 5 °C and the solution was stirred at 20 °C for 6 d. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give compound **17** (102 mg, 80%) as a red gum, ¹H NMR [(CD₃)₂SO] δ 11.39 (t, *J* = 5.5 Hz, 1 H, CONH), 9.31 (s, 1 H, H-9), 8.71 (dd, *J* = 7.1, 1.4 Hz, 1 H, H-3), 8.48 (br s, 1 H, NH), 8.38 (dd, *J* = 8.4, 1.4 Hz, 1 H, H-1), 8.33 (d, *J* = 9.2 Hz, 1 H, H-5), 8.22 (d, *J* = 8.5 Hz, 1 H, H-8), 8.13 (d, *J* = 8.4 Hz, 1 H, H-8'), 8.06 (d, *J* = 8.7 Hz, 1 H, H-5'), 7.87–7.95 (m, 2 H, H-6, H-6'), 7.76 (dd, *J* = 8.4, 7.1 Hz, 1 H, H-2), 7.68 (dd, *J* = 8.5, 7.2 Hz, 1 H, H-7), 7.54 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-7'), 3.64–3.70 (m, 4 H, 2 CH₂N), 2.05–2.10 (m, 2 H, CH₂); ¹³C NMR (CD₃OD) δ 168.9, 152.2, 151.5 (2), 141.0, 140.5, 140.1 (2), 138.8, 137.9, 135.7, 133.7, 131.5, 130.1, 128.9, 128.5, 128.1, 127.0, 123.0, 121.9, 121.6, 40.6, 38.1, 29.6. An analytical sample was recrystallized as the dihydrochloride salt, mp (MeOH/EtOAc) 192 °C. Anal. calcd for C₂₄H₂₀N₆O₃·2HCl·½H₂O: C, 55.2; H, 4.4; N, 16.1; found: C, 55.3; H, 4.5; N, 16.1%.

Example F.

***N*-(2-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethoxy}ethyl)-4-acridinecarboxamide (23).**

3-{[2-(2-Hydroxyethoxy)ethyl]amino}-1,2,4-benzotriazine 1-oxide (18). A solution
5 of chloride **3** (3.0 g, 16.52 mmol) in DCM (50 mL) was added to a stirred solution of
2-(aminoethoxy)ethanol (2.49 mL, 24.8 mmol) and Et₃N (3.45 mL, 24.8 mmol) in
DCM (80 mL) and the solution stirred at 20 °C for 16 h. The solvent was evaporated
and the residue purified by chromatography, eluting with 40% EtOAc/DCM, to give
1-oxide **18** (2.62 g, 63%) as a yellow powder, mp (DCM/EtOAc) 131–131.5 °C; ¹H
10 NMR δ 8.25 (dd, *J* = 8.7, 1.2 Hz, 1 H, H-8), 7.68 (ddd, *J* = 8.4, 7.2, 1.5 Hz, 1 H, H-6),
7.57 (d, *J* = 8.4 Hz, 1 H, H-5), 7.28 (ddd, *J* = 8.7, 7.2, 1.3 Hz, 1 H, H-7), 6.02 (br s, 1
H, NH), 3.74–3.80 (m, 6 H, 3 × CH₂O), 3.64–3.67 (m, 2 H, CH₂N), 2.71 (t, *J* = 5.9
Hz, 1 H, OH); ¹³C NMR δ 158.9, 149.7, 135.5, 130.9, 126.4, 124.9, 120.4, 72.4, 69.5,
61.7, 41.9. Anal. calcd for C₁₁H₁₄N₄O₃: C, 52.8; H, 5.6; N, 22.4; found: C, 52.9; H,
15 5.7; N, 22.6%.

3-{[2-(2-Azidoethoxy)ethyl]amino}-1,2,4-benzotriazine 1-oxide (19).

Methanesulfonyl chloride (0.82 mL, 10.6 mmol) was added dropwise to a stirred
solution of alcohol **18** (2.41 g, 9.63 mmol) and Et₃N (1.74 mL, 12.5 mmol) in DCM
20 (100 mL) at 5 °C and the solution stirred at 20 °C for 1 h. The solution was diluted
with DCM (100 mL) and washed with water (3 × 50 mL), brine (50 mL), dried and the
solvent evaporated. The residue was dissolved in DMF (50 mL) and NaN₃ (0.69 g,
10.6 mmol) added. The mixture was heated at 100 °C for 2 h, cooled to 30 °C and the
solvent evaporated. The residue was partitioned between EtOAc (100 mL) and water
25 (100 mL). The organic fraction was washed with brine (50 mL), dried, and the solvent
evaporated. The residue was purified by chromatography, eluting with 50%
EtOAc/pet. ether, to give azide **19** (2.35 g, 89%) as yellow crystals, mp (EtOAc/pet.
ether) 102–104 °C; ¹H NMR δ 8.27 (dd, *J* = 8.7, 1.4 Hz, 1 H, H-8), 7.70 (ddd, *J* = 8.6,
7.1, 1.5 Hz, 1 H, H-6), 7.59 (d, *J* = 8.6 Hz, 1 H, H-5), 7.29 (ddd, *J* = 8.6, 7.1, 1.4 Hz, 1
30 H, H-7), 5.70 (br s, 1 H, NH), 3.71–3.78 (m, 4 H, 2 × CH₂O), 3.69 (dd, *J* = 5.3, 4.8
Hz, 2 H, CH₂N₃), 3.41 (dd, *J* = 5.1, 4.9 Hz, 2 H, CH₂N); ¹³C NMR δ 158.9, 148.7,
135.5, 131.1, 126.5, 125.0, 120.4, 70.0, 69.6, 50.7, 41.1. Anal. calcd for C₁₁H₁₃N₇O₂:
C, 48.0; H, 4.8; N, 35.6; found: C, 48.3; H, 4.6; N, 35.7%.

3-[[2-(2-*tert*-Butyloxycarbamoylethoxy)ethyl]amino}-1,2,4-benzotriazine 1-oxide

(**20**). Propane-1,3-dithiol (5.7 mL, 57.0 mmol) was added dropwise to a stirred solution of azide **19** (1.57 g, 5.70 mmol) and Et₃N (7.95 mL, 57 mmol) in MeOH (100 mL) under N₂ and the solution heated at reflux temperature for 8 h. The solution was cooled to 30 °C and partitioned between 1 M HCl (100 mL) and Et₂O (100 mL). The aqueous fraction was adjusted to pH 12 with 7 M NaOH solution and extracted with DCM (3 × 50 mL). The organic fraction was dried and the solvent evaporated. The residue was dissolved in THF (100 mL) and a solution of di-*tert*-butyldicarbonate (1.87 g, 8.55 mmol) in THF (50 mL) added dropwise. The solution was stirred at 20 °C for 16 h, the solvent evaporated and the residue purified by chromatography, eluting with 40% EtOAc/pet. ether, to give carbamate **20** (1.85 g, 93%) as a yellow solid, mp (EtOAc/pet. ether) 134–137 °C; ¹H NMR δ 8.26 (dd, *J* = 8.4, 0.9 Hz, 1 H, H-8), 7.71 (ddd, *J* = 8.3, 7.1, 1.4 Hz, 1 H, H-6), 7.59 (d, *J* = 8.3 Hz, 1 H, H-5), 7.29 (ddd, *J* = 8.4, 7.1, 1.3 Hz, 1 H, H-7), 5.74 (br s, 1 H, NH), 4.93 (br s, 1 H, NH), 3.67–3.73 (m, 4 H, 2 × CH₂O), 3.56 (t, *J* = 5.2 Hz, 2 H, CH₂N), 3.29–3.36 (m, 2 H, CH₂N), 1.45 [s, 9 H, C(CH₃)₃]; ¹³C NMR δ 159.9, 155.9, 148.7, 135.5, 131.0, 126.5, 125.0, 120.4, 79.4, 70.2, 69.2, 41.1, 40.4, 28.4 (3). Anal. calcd for C₁₆H₂₃N₅O₄: C, 55.0; H, 6.6; N, 20.1; found: C, 55.3; H, 6.8; N, 20.1%.

3-[[2-(2-*tert*-Butyloxycarbamoylethoxy)ethyl]amino}-1,2,4-benzotriazine 1,4-

dioxide (21). A solution of MCPBA (1.57 g, 6.35 mmol) in DCM (50 mL) was added dropwise to a stirred solution of carbamate **20** (1.85 g, 5.29 mmol) in DCM (100 mL) and NaHCO₃ (0.89 g, 10.6 mmol) and the mixture was stirred at 20 °C for 6 h. The suspension was filtered through celite, the solvent evaporated and the residue purified by chromatography, eluting with a gradient of (0–5%) MeOH/(40–0%) EtOAc/DCM, to give (i) starting material **20** (926 mg, 50%), spectroscopically identical with an authentic sample, and (ii) 1,4-dioxide **21** (702 mg, 40%) as a red solid, mp (EtOAc) 139–140 °C; ¹H NMR δ 8.33 (d, *J* = 8.7 Hz, 1 H, H-8), 8.30 (d, *J* = 8.7 Hz, 1 H, H-5), 7.88 (ddd, *J* = 8.7, 7.2, 1.2 Hz, 1 H, H-6), 7.43–7.50 (m, 2 H, H-7, NH), 5.06 (br s, 1 H, NH), 3.78–3.83 (m, 2 H, CH₂O), 3.69 (dd, *J* = 5.1, 5.0 Hz, 2 H, CH₂O), 3.56 (dd, *J* = 5.1, 5.0 Hz, 2 H, CH₂N), 3.29–3.36 (m, 2 H, CH₂N), 1.43 [s, 9 H, C(CH₃)₃]; ¹³C NMR δ 156.0, 149.8, 138.5, 135.9, 130.6, 129.5, 121.6, 117.4, 79.4, 70.3, 68.9, 41.3,

40.3, 28.3 (3); MS (FAB⁺) m/z 366 (MH⁺, 40%), 350 (5) 310 (20); HRMS (FAB⁺) calcd for C₁₆H₂₄N₅O₅ (MH⁺) m/z 366.1777, found 366.1767. Anal. calcd for C₁₆H₂₃N₅O₅·½H₂O: C, 51.3; H, 6.5; N, 18.7; found: C, 51.3; H, 6.2; N, 16.9%.

5 **3-{[2-(2-Aminoethoxy)ethyl]amino}-1,2,4-benzotriazine 1,4-dioxide (22).**

Trifluoroacetic acid (1.66 mL, 34.6 mmol) was added dropwise to a stirred solution of 1,4-dioxide **21** (632 mg, 1.73 mmol) in DCM (50 mL) and the solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue partitioned between sat. aqueous KHCO₃ solution (100 mL) and CHCl₃ (100 mL). The aqueous phase was extracted
10 with CHCl₃ (8 × 50 mL), the combined organic fractions dried, and the solvent evaporated. The residue was crystallized from CHCl₃ to give the amine **22** (406 mg, 91%) as a red solid, mp (CHCl₃) 124 °C (dec.); ¹H NMR δ 8.26 (d, J = 8.9 Hz, 1 H, H-8), 8.23 (d, J = 8.9 Hz, 1 H, H-5), 7.79 (dd, J = 8.8, 7.8 Hz, 1 H, H-6), 7.45 (dd, J = 8.9, 7.7 Hz, 1 H, H-7), 3.75 (dd, J = 5.0, 4.8 Hz, 2 H, CH₂O), 3.66 (dd, J = 5.0, 4.9
15 Hz, 2 H, CH₂O), 3.47 (dd, J = 5.1, 5.0 Hz, 2 H, CH₂N), 2.82 (dd, J = 5.1, 5.0 Hz, 2 H, CH₂N), NH and NH₂ not observed; ¹³C NMR δ 149.8, 138.3, 135.8, 130.5, 127.2, 121.6, 117.4, 73.0, 68.9, 41.7, 41.3; MS (FAB⁺) m/z 266 (MH⁺, 20%), 250 (5); HRMS (FAB⁺) calcd for C₁₁H₁₆N₅O₃ (MH⁺) m/z 266.1253, found 266.1230. Anal. calcd for C₁₁H₁₅N₅O₃·¼H₂O: C, 49.0; H, 5.8; N, 26.0; found: C, 49.0; H, 5.7; N, 24.7%.

20

***N*-(2-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethoxy}ethyl)-4-**

acridinecarboxamide (23). A solution of the amine **22** (54 mg, 0.20 mmol) in THF (2 mL) was added dropwise to a stirred solution of 4-(1*H*-imidazol-1-ylcarbonyl)acridine (58 mg, 0.21 mmol) in THF (5 mL) at 5 °C and the solution stirred at 20 °C for 16 h.

25 The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0-5%) of MeOH/DCM, to give compound **23** (93 mg, 97%) as a red solid, mp (EtOAc) 98–100 °C; ¹H NMR δ 12.14 (s, 1 H, CONH), 8.96 (dd, J = 7.1, 1.5 Hz, 1 H, H-3'), 8.82 (s, 1 H, H-9), 8.25 (d, J = 8.4 Hz, 1 H, H-8'), 8.16 (d, J = 8.4 Hz, 1 H, H-5'), 8.11–8.13 (m, 2 H, H-1, H-5), 7.94 (d, J = 8.2 Hz, 1 H, H-8), 7.76–7.84 (m, 2
30 H, H-6, H-6'), 7.66 (dd, J = 8.4, 7.1 Hz, 1 H, H-2), 7.44–7.52 (m, 2 H, H-7, H-7'), 7.36 (br s, 1 H, NH), 3.85–3.95 (m, 8 H, 2 × CH₂O, 2 × CH₂N); ¹³C NMR δ 166.1, 149.8, 147.2, 146.3, 138.1, 137.6, 135.5, 135.3, 132.4, 131.3, 130.4, 128.8, 128.3, 128.0, 127.1, 126.8, 126.2, 125.8, 125.4, 121.5, 117.3, 70.2, 68.9, 41.1, 39.5; MS

(FAB⁺) *m/z* 471 (MH⁺, 5%), 455 (4); HRMS (FAB⁺) calcd for C₂₅H₂₃N₆O₄ (MH⁺) *m/z* 471.1781, found 471.1790. Anal. calcd for C₂₅H₂₂N₆O₄·½H₂O: C, 62.6; H, 4.8; N, 17.5; found: C, 63.0; H, 4.7; N, 17.5%.

5 Example G.

N-(2-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethoxy}ethyl)-8-

quinolinecarboxamide (24). A solution of 8-quinolinecarboxylic acid (308 mg, 1.78 mmol) and CDI (346 mg, 2.13 mmol) in DMF (20 mL) were stirred at 50 °C for 1 h. The solvent was evaporated and the residue recrystallised from DCM/pet. ether to give
 10 4-(1*H*-imidazol-1-ylcarbonyl)quinoline (50 mg, 0.21 mmol) which was used directly without characterisation. A solution of the amine **22** (57 mg, 0.21 mmol) in DCM (10 mL) was added dropwise to a stirred solution of imidazolidine (50 mg, 0.21 mmol) in DCM (5 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–
 5 5%) of MeOH/DCM, to give compound **24** (74 mg, 84%) as a red powder, mp (MeOH/DCM) 168–170 °C; ¹H NMR δ 11.51 (br s, 1 H, NH), 9.01 (dd, *J* = 4.2, 1.9 Hz, 1 H, H-2), 8.85 (dd, *J* = 7.3, 1.6 Hz, 1 H, H-4), 8.30 (d, *J* = 8.3 Hz, 1 H, H-8”), 8.23–8.26 (m, 2 H, H-7, H-5”), 7.93 (dd, *J* = 8.1, 1.5 Hz, 1 H, H-5), 7.86 (ddd, *J* = 8.4, 7.0, 1.8 Hz, 1 H, H-6”), 7.67 (dd, *J* = 7.9, 7.5 Hz, 1 H, H-6), 7.46–7.51 (m, 2 H, H-3, H-7”), 7.46 (br s, 1 H, NH), 3.78–3.85 (m, 8 H, 2 × CH₂O, 2 × CH₂N); ¹³C NMR δ
 10 166.0, 149.8, 149.6, 145.6, 138.3, 137.6, 135.7, 133.8, 131.9, 130.5, 128.7, 128.4, 127.2, 126.4, 121.6, 120.9, 117.4, 70.3, 68.9, 41.4, 39.6; MS (FAB⁺) *m/z* 421 (MH⁺, 8%), 405 (5), 389 (1); HRMS (FAB⁺) calcd for C₂₁H₂₁N₆O₄ (MH⁺) *m/z* 421.1624, found 421.1615. Anal. calcd for C₂₁H₂₀N₆O₄·½MeOH: C, 59.2; H, 5.1; N, 19.3;
 15 found: C, 59.2; H, 4.8; N, 19.2%.

Example H.

N-(2-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethoxy}ethyl)-2-phenyl-1*H*-

benzimidazole-4-carboxamide (25). A solution of 2-phenyl-1*H*-benzimidazole-4-
 10 carboxylic acid (396 mg, 1.67 mmol) and CDI (270 mg, 1.67 mmol) in DMF (10 mL) was stirred at 50 °C for 1 h. The solvent was evaporated and the residue recrystallised from DCM/pet. ether to give 4-(1*H*-imidazol-1-ylcarbonyl)-2-phenyl-1*H*-benzimidazole (309 mg, 0.21 mmol) which was used directly without characterisation. A solution of

the amine **22** (56 mg, 0.21 mmol) in DCM (5 mL) was added dropwise to a stirred solution of imidazolide (61 mg, 0.21 mmol) in DCM (5 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give compound **25** (89 mg, 86%) as a red powder, mp (DCM) 203–207 °C; ¹H NMR [(CD₃)₂SO] δ 13.30 (br s, 1 H, NH), 10.24 (s, 1 H, NH), 8.18–8.24 (m, 3 H, H-2', H-6', NH), 8.13 (d, *J* = 8.7 Hz, 1 H, H-8''), 8.04 (d, *J* = 8.5 Hz, 1 H, H-5''), 7.90–7.94 (m, 1 H, H-6''), 7.87 (d, *J* = 7.9 Hz, 1 H, H-5), 7.72 (d, *J* = 7.9 Hz, 1 H, H-7), 7.52–7.58 (m, 3 H, H-3', H-5', H-7''), 7.46–7.48 (m, 1 H, H-4'), 7.34 (t, *J* = 7.9 Hz, 1 H, H-6), 3.78–3.82 (m, 2 H, CH₂O), 3.74–3.77 (m, 2 H, CH₂O), 3.63–3.78 (m, 4 H, 2 × CH₂N); ¹³C NMR [(CD₃)₂SO] δ 164.5, 151.7, 149.7, 141.0, 138.0, 135.4, 135.1, 130.4, 129.9, 128.9 (2), 128.8, 126.9, 126.6 (2), 122.5, 122.3, 122.0, 121.0, 116.7, 114.8, 69.1, 68.2, 40.3, 38.8; MS (FAB⁺) *m/z* 486 (MH⁺, 4%), 470 (2); HRMS (FAB⁺) calcd for C₂₅H₂₄N₇O₄ (MH⁺) *m/z* 486.1890, found 486.1903. Anal. calcd for C₂₅H₂₃N₇O₄: C, 61.8; H, 4.8; N, 20.2; found: C, 61.6; H, 4.7; N, 20.0%.

Example I.

***N*-(2-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethoxy}ethyl)-2-(4-pyridinyl)-8-quinolinecarboxamide (26).** A solution of 2-(4-pyridinyl)-8-

quinolinecarboxylic acid (268 mg, 1.07 mmol) and CDI (173 mg, 1.07 mmol) in DMF (10 mL) were stirred at 50 °C for 1 h. The solvent was evaporated and the residue recrystallized from DCM/pet. ether to give 8-(1*H*-imidazol-1-ylcarbonyl)-2-(4-pyridinyl)quinoline (238 mg, 0.86 mmol) which was used directly without characterization. A solution of the amine **22** (39 mg, 0.15 mmol) in DCM (5 mL) was added dropwise to a stirred solution of imidazolide (41 mg, 0.15 mmol) in DCM (5 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give compound **26** (51 mg, 70%) as a red powder, mp (DCM) 128–130 °C; ¹H NMR [(CD₃)₂SO] δ 12.01 (br s, 1 H, NH), 10.83 (t, *J* = 5.3 Hz, 1 H, NH), 8.70 (dd, *J* = 4.5, 1.5 Hz, 2 H, H-3', H-5'), 8.63 (d, *J* = 8.6 Hz, 1 H, H-7), 8.58 (dd, *J* = 7.3, 1.5 Hz, 1 H, H-4), 8.23 (d, *J* = 8.7 Hz, 1 H, H-5), 8.19 (dd, *J* = 8.7, 1.5 Hz, 1 H, H-8''), 8.09 (dd, *J* = 4.5, 1.5 Hz, 2 H, H-2', H-6'), 7.95 (d, *J* = 8.5 Hz, 1 H, H-5''), 7.82–7.90 (m, 2 H, H-3, H-6''), 7.76 (t, *J* = 8.7 Hz, 1 H, H-6), 7.47 (ddd, *J* = 8.7, 7.0,

1.6 Hz, 1 H, H-7''), 3.70-3.78 (m, 6 H, 2 × CH₂O, CH₂N), 3.49-3.54 (m, 2 H, CH₂N); MS (FAB⁺) *m/z* 498 (MH⁺, 10%), 482 (5); HRMS (FAB⁺) calcd for C₂₆H₂₄N₇O₄ (MH⁺) *m/z* 498.1890, found 498.1898. Anal. calcd for C₂₆H₂₃N₇O₄·H₂O: C, 60.7; H, 4.9; N, 22.0; found: C, 60.6; H, 4.9; N, 19.0%.

5

Example J.

N-[3-({3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-4-acridinecarboxamide (30).

tert-Butyl 3-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]propyl{3-

- 10 [(trifluoroacetyl)amino]propyl}carbamate (27). A solution of chloride 3 (1.34 g, 7.41 mmol) in DCM (50 mL) was added dropwise to a stirred solution of *tert*-butyl bis(3-aminopropyl)carbamate (2.57 g, 11.1 mmol) and Et₃N (1.55 mL, 11.1 mmol) in DCM (200 mL) at 20 °C. The solution was stirred at 20 °C for 3 d. The solvent was evaporated and the residue purified by chromatography, eluting with 50%
15 EtOAc/acetone, to give a crude oil (2.31 g). Trifluoroacetic anhydride (3.5 mL, 24.3 mmol) was added dropwise to a stirred solution of crude amine in pyridine (50 mL) at 5 °C. The solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (30-50%) of EtOAc/pet. ether, to give trifluoroacetamide 27 (0.51 g, 22%) as a yellow solid, mp (EtOAc/pet.
20 ether) 89-90 °C; ¹H NMR δ 8.22-8.26 (m, 2 H, H-8, NH), 7.71 (br dd, *J* = 8.4, 7.0 Hz, 1 H, H-6), 7.59 (d, *J* = 8.4 Hz, 1 H, H-5), 7.29 (br dd, *J* = 8.5, 7.0 Hz, 1 H, H-7), 5.45 (br s, 1 H, NH), 4.12 (br dd, *J* = 6.6, 6.5 Hz, 2 H, CH₂N), 3.26-3.37 (m, 6 H, 3 × CH₂N), 1.84-1.95 (m, 2 H, CH₂), 1.71-1.77 (m, 2 H, CH₂), 1.48 [s, 9 H, C(CH₃)₃]; ¹³C NMR δ 158.9, 157.3 (q, *J* = 37 Hz), 156.8, 148.0, 135.6, 130.9, 126.5, 125.1,
25 120.4, 116 (q, *J* = 288 Hz), 80.8, 44.5, 43.0, 38.8, 35.8, 29.7, 28.3 (3), 27.1; MS (FAB⁺) *m/z* 473 (MH⁺, 60%), 457 (10), 373 (100); HRMS (FAB⁺) calcd for C₂₀H₂₈F₃N₆O₄ (MH⁺) *m/z* 473.2124, found 473.2136. Anal. calcd for C₂₀H₂₇F₃N₆O₄: C, 50.8; H, 5.8; N, 17.8; found: C, 50.5; H, 5.7; N, 17.8%.

- 30 *tert*-Butyl 3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl{3-[(trifluoroacetyl)amino]propyl}carbamate (28). A solution of MCPBA (2.12 g, 8.6 mmol) in DCM (50 mL) was added dropwise to a stirred solution of 1-oxide 27 (3.13 g, 6.6 mmol) in DCM (250 mL) and NaHCO₃ (1.1 g, 13.2 mmol). The mixture was

stirred at 20 °C for 16 h, partitioned between DCM (200 mL) and sat. aqueous KHCO₃ solution (100 mL). The organic fraction was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–4%) of MeOH/40%EtOAc/DCM, to give (i) starting material **27** (2.04 g, 65%) and (ii) 1,4-dioxide **28** (252 mg, 8 %) as a red solid, ¹H NMR δ 8.34 (d, *J* = 8.7 Hz, 1 H, H-8), 8.30 (d, *J* = 8.4 Hz, 1 H, H-5), 8.25 (br s, 1 H, NH), 7.88 (br dd, *J* = 8.4, 7.0 Hz, 1 H, H-6), 7.52 (br dd, *J* = 8.7, 7.0 Hz, 1 H, H-7), 7.20 (br s, 1 H, NH), 3.62 (dt, *J* = 6.8, 6.7 Hz, 2 H, CH₂N), 3.26–3.38 (m, 6 H, 3 × CH₂N), 1.92–1.98 (m, 2 H, CH₂), 1.73–1.79 (m, 2 H, CH₂), 1.49 [s, 9 H, C(CH₃)₃]; ¹³C NMR δ 157.3 (q, *J* = 37 Hz), 156.8, 149.8, 138.2, 135.9, 130.5, 127.4, 121.7, 117.4, 116.1 (q, *J* = 288 Hz), 80.9, 44.4, 43.2, 38.9, 31.9, 29.7, 28.4 (3), 22.7; MS (FAB⁺) *m/z* 489 (MH⁺, 10%), 473 (12), 373 (15); HRMS (FAB⁺) calcd for C₂₀H₂₈F₃N₆O₅ (MH⁺) *m/z* 489.2073, found 489.2086.

***tert*-Butyl 3-aminopropyl{3-[(1,4-dioxido-1,2,4-benzotriazin-3-**

yl)amino]propyl}carbamate (29**).** A mixture of trifluoroacetamide **28** (541 mg, 1.11 mmol) and K₂CO₃ (0.77 g, 5.54 mmol) in MeOH (20 mL) and water (5 mL) was heated at reflux temperature for 1 h. The mixture was partitioned between CHCl₃ (50 mL) and water (30 mL). The aqueous fraction was extracted with CHCl₃ (3 × 30 mL), the combined organic fraction dried, and the solvent evaporated to give amine **29** (322 mg, 74%) as a red oil, ¹H NMR [(CD₃)₂SO] δ 10.50 (br s, 1 H, NH), 8.21 (d, *J* = 8.7 Hz, 1 H, H-8), 8.13 (d, *J* = 8.6 Hz, 1 H, H-5), 7.94 (br dd, *J* = 8.6, 7.5 Hz, 1 H, H-6), 7.56 (br dd, *J* = 8.6, 7.5 Hz, 1 H, H-7), 7.20 (br s, 2 H, NH₂), 3.39 (t, *J* = 6.9 Hz, 2 H, CH₂N), 3.11–3.21 (m, 6 H, 3 × CH₂N), 1.78–1.86 (m, 2 H, CH₂), 1.49–1.58 (m, 2 H, CH₂), 1.39 [s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 154.7, 149.7, 138.1, 135.4, 129.8, 127.9, 121.0, 116.7, 78.3, 44.3, 43.9, 38.8, 38.4, 32.2, 31.6, 27.9 (3); MS (FAB⁺) *m/z* 393 (MH⁺, 15%), 377 (9), 338 (3); HRMS (FAB⁺) calcd for C₁₈H₂₉N₆O₄ (MH⁺) *m/z* 393.2250, found 393.2249.

***N*-[3-({3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-4-**

acridinecarboxamide (30**).** A solution of 4-acridinecarboxylic acid (846 mg, 4.35 mmol) and CDI (846 mg, 5.21 mmol) in DMF (20 mL) were stirred at 50 °C for 1 h. The solvent was evaporated and the residue recrystallized from DCM/pet. ether to give 4-(1*H*-imidazol-1-ylcarbonyl)acridine (746 mg, 63%) which was used directly

without characterization. A solution of the amine **29** (320 mg, 0.82 mmol) in DCM (10 mL) was added dropwise to a stirred solution of imidazolide (234 mg, 0.86 mmol) in THF (25 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give *tert*-butyl 3-[(4-acridinylcarbonyl)amino]propyl{3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}carbamate (330 mg, 67%) as a red gum, ¹H NMR δ 11.92 (br s, 1 H, CONH), 8.98 (dd, *J* = 7.2, 1.5 Hz, 1 H, H-3), 8.89 (s, 1 H, H-9), 8.26–8.32 (m, 3 H, H-5, H-5', H-8'), 8.16 (d, *J* = 8.3 Hz, 1 H, H-1), 8.07 (d, *J* = 8.8 Hz, 1 H, H-8), 7.82–7.89 (m, 3 H, H-3, H-6, H-6'), 7.65–7.69 (m, 1 H, H-7'), 7.58–7.62 (m, 1 H, H-7), 7.48 (br s, 1 H, NH), 3.72 (dt, *J* = 6.6, 6.0 Hz, 2 H, CH₂N), 3.61 (dt, *J* = 6.6, 6.4 Hz, 2 H, CH₂N), 3.38–3.50 (m, 4 H, 2 × CH₂N), 2.04–2.08 (m, 2 H, CH₂), 1.88–1.94 (m, 2 H, CH₂), 1.40 [s, 9 H, C(CH₃)₃]; MS (FAB⁺) *m/z* 598 (MH⁺, 8%), 582 (6); HRMS (FAB⁺) calcd for C₃₂H₃₆N₇O₅ (MH⁺) *m/z* 598.2778, found 598.2772.

HCl saturated MeOH (30 mL) was added to a solution of carbamate (328 mg, 0.55 mmol) in MeOH (30 mL) and the solution stirred at 20 °C for 16 h. The solution was evaporated and the residue dissolved in water (20 mL) the solution neutralized with KHCO₃ and extracted with CHCl₃ (5 × 50 mL). The combined organic fraction was dried and the solvent evaporated to give compound **30** (247 mg, 90%) as a red solid, ¹H NMR [(CD₃)₂SO] δ 11.38 (t, *J* = 5.5 Hz, 1 H, CONH), 10.50 (br s, 1 H, NH), 9.28 (s, 1 H, H-9), 8.71 (dd, *J* = 7.1, 1.5 Hz, 1 H, H-3), 8.35 (dd, *J* = 8.4, 1.5 Hz, 1 H, H-1), 8.24 (d, *J* = 8.7 Hz, 1 H, H-5), 8.19 (d, *J* = 8.3 Hz, 1 H, H-8), 8.14 (d, *J* = 8.5 Hz, 1 H, H-8'), 8.03 (d, *J* = 8.5 Hz, 1 H, H-5'), 7.92–7.96 (m, 1 H, H-6), 7.83–7.88 (m, 1 H, H-6'), 7.75 (dd, *J* = 8.3, 7.1 Hz, 1 H, H-2), 7.65–7.68 (m, 1 H, H-7), 7.48–7.54 (m, 1 H, H-7'), 7.38 (s, 1 H, NH), 3.64 (dt, *J* = 6.9, 5.9 Hz, 2 H, CH₂N), 3.46 (t, *J* = 6.7 Hz, 2 H, CH₂N), 2.79 (t, *J* = 6.9 Hz, 2 H, CH₂N), 2.70 (t, *J* = 6.5 Hz, 2 H, CH₂N), 1.88–1.94 (m, 2 H, CH₂), 1.76–1.82 (m, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 164.7, 149.6, 147.0, 145.4, 138.5, 138.0, 135.3, 134.4, 132.6, 131.8, 129.7, 128.5, 128.4, 128.3, 126.7, 126.4, 126.3, 125.5, 125.2, 121.0, 116.7, 47.1, 46.9, 39.6, 37.2, 29.3, 28.2; MS (FAB⁺) *m/z* 498 (MH⁺, 15%), 482 (5); HRMS (FAB⁺) calcd for C₂₇H₂₈N₇O₃ (MH⁺) *m/z* 498.2254, found 498.2258. Anal. calcd for C₂₇H₂₇N₇O₃·2H₂O: C, 60.8; H, 5.9; N, 18.4; found: C, 60.7; H, 5.6; N, 17.1%.

Example K.

N-[3-({3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-1-

phenazinecarboxamide hydrochloride (31). A solution of the amine **29** (223 mg,

0.57 mmol) in THF (10 mL) was added dropwise to a stirred solution of 1-(1*H*-

imidazol-1-ylcarbonyl)phenazine (171 mg, 0.63 mmol) in THF (25 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue

purified by chromatography, eluting with a gradient (0–3%) of MeOH/DCM, to give

tert-butyl 3-[(1-phenazinecarbonyl)amino]propyl{3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}carbamate (137 mg, 40%) as a red gum, ¹H NMR δ 11.22 and

11.06 (2 × s, 1 H, CONH), 9.03 (dd, *J* = 7.1, 1.4 Hz, 1 H, H-2), 8.64 and 8.27 (2 × s, 1 H, NH), 8.42 (d, *J* = 8.2 Hz, 1 H, H-9), 8.29–8.37 (m, 3 H, H-4, H-6, H-8"), 7.86–

8.03 (m, 5 H, H-3, H-7, H-8, H-5", H-6"), 7.49–7.56 (m, 1 H, H-7"), 3.69–3.77 (m, 2 H, CH₂N), 3.63–3.68 (m, 2 H, CH₂N), 1.93–2.10 (m, 4 H, 2 × CH₂N), 1.67 and 1.63

[2 × s, 9 H, C(CH₃)₃], 1.45–1.53 (m, 4 H, 2 × CH₂); MS (FAB⁺) *m/z* 599 (MH⁺, 12%),

583 (3); HRMS (FAB⁺) calcd for C₃₁H₃₅N₈O₅ (MH⁺) *m/z* 599.2730, found 599.2733.

HCl saturated MeOH (5 mL) was added to a solution of carbamate (135 mg, 0.23

mmol) in MeOH (20 mL) and the solution stirred at 20 °C for 16 h. The solution was

evaporated and the residue dissolved in water (20 mL) the solution neutralized with

dil. aqueous NH₃ and extracted with CHCl₃ (5 × 50 mL). The combined organic

fraction was dried and the solvent evaporated to give compound **31** (97 mg, 85%) as a

red solid, which was converted to the HCl salt and recrystallized, mp (MeOH/EtOAc)

163–169 °C; ¹H NMR [(CD₃)₂SO] δ 10.29 (t, *J* = 5.8 Hz, 1 H, CONH), 9.26 (br s, 2

H, NH₂⁺Cl⁻), 8.97 (t, *J* = 6.1 Hz, 1 H, NH), 8.59 (dd, *J* = 9.0, 2.0 Hz, 1 H, H-2), 8.55

(dd, *J* = 9.0, 2.0 Hz, 1 H, H-9), 8.41 (dd, *J* = 8.7, 1.3 Hz, 1 H, H-4), 8.28 (dd, *J* = 7.9,

2.0 Hz, 1 H, H-6), 8.19 (d, *J* = 8.2 Hz, 1 H, H-8"), 7.98–8.08 (m, 5 H, H-3, H-7, H-8,

H-5", H-6"), 7.60 (ddd, *J* = 8.7, 7.1, 1.4 Hz, 1 H, H-7"), 3.65–3.69 (m, 2 H, H'),

3.55–3.59 (m, 2 H, H-3"), 3.04–3.13 (m, 4 H, H-3', H-1"), 2.03–2.15 (m, 4 H, H-2', H-

2"); ¹³C NMR [(CD₃)₂SO] δ 164.8, 149.8, 142.6, 142.5, 141.3, 139.9, 137.5, 136.5,

133.4, 132.6, 131.8, 131.6, 131.0, 130.5, 130.2, 129.5, 129.0, 127.5, 121.8, 116.2,

44.6, 44.2, 38.1, 36.4, 25.9, 25.1. Anal. calcd for C₂₆H₂₇ClN₈O₃·MeOH: C, 57.2; H,

5.5; N, 19.8; found: C, 57.3; H, 5.8; N, 20.0%.

Example L

N-[3-({3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-9-methyl-1-phenazinecarboxamide hydrochloride (32). A solution of the amine 29 (265 mg, 0.68 mmol) in THF (10 mL) was added dropwise to a stirred solution of 1-(1*H*-imidazol-1-ylcarbonyl)-9-methylphenazine (214 mg, 0.74 mmol) in THF (25 mL) at 5 °C and the solution was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (5–10%) of MeOH/DCM, to give *tert*-butyl 3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl(3-[(9-methyl-1-phenazinyl)carbonyl]amino)propyl)carbamate (168 mg, 40%) as a red gum, MS (FAB⁺) *m/z* 613 (MH⁺, 20%), 597 (5), 513 (15), 497 (5); HRMS (FAB⁺) calcd for C₃₂H₃₇N₈O₅ (MH⁺) *m/z* 613.2887, found 613.2881.

HCl saturated MeOH (5 mL) was added to a solution of carbamate (168 mg, 0.27 mmol) in MeOH (20 mL) and the solution stirred at 20 °C for 16 h. The solution was evaporated and the residue dissolved in water (20 mL) the solution neutralized with dil. aqueous NH₃ and extracted with CHCl₃ (5 × 50 mL). The combined organic fraction was dried and the solvent evaporated to give compound 32 (121 mg, 86%) as a red solid, which was converted to the HCl salt and recrystallized, mp (MeOH/EtOAc) 183–186 °C; ¹H NMR [(CD₃)₂SO] δ 10.45 (t, *J* = 5.8 Hz, 1 H, CONH), 9.18 (br s, 2 H, NH₂⁺Cl⁻), 8.73 (t, *J* = 6.2 Hz, 1 H, NH), 8.63 (dd, *J* = 7.0, 1.4 Hz, 1 H, H-2), 8.39 (dd, *J* = 8.7, 1.4 Hz, 1 H, H-4), 8.18 (d, *J* = 8.7 Hz, 1 H, H-8"), 8.03–8.11 (m, 3 H, H-3, H-5", H-7), 7.92 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-6"), 7.87–7.93 (m, 2 H, H-6, H-8), 7.57 (ddd, *J* = 8.7, 7.1, 1.3 Hz, 1 H, H-7"), 3.61–3.66 (m, 2 H, H'), 3.51–3.57 (m, 2 H, H-3"), 2.98–3.10 (m, 4 H, H-3', H-1"), 2.86 (s, 3 H, CH₃), 2.08–2.15 (m, 2 H, H-2'), 1.99–2.05 (m, 2 H, H-2"); ¹³C NMR [(CD₃)₂SO] δ 164.5, 149.7, 142.6, 142.3, 140.4, 138.7, 137.7, 136.6, 136.0, 133.7, 132.7, 131.5, 131.2, 130.2, 130.1, 130.0, 127.1, 127.0, 121.0, 116.4, 44.8, 44.7, 37.9, 36.6, 26.2, 25.1, 17.5; MS (FAB⁺) *m/z* 513 (MH⁺, 20%), 497 (5); HRMS (FAB⁺) calcd for C₂₇H₂₉N₈O₃ (MH⁺) *m/z* 513.2363, found 513.2352.

Example M

N-[2-({2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}amino)ethyl]-4-acridinecarboxamide (37).

tert-Butyl bis-(2-aminoethyl)carbamate (33). Diethylenetriamine (9.9 mL, 96 mmol) was added to a solution of CF₃CO₂Et (22.8 mL, 192 mmol) in dry ether (80

mL) at 5 °C and the reaction mixture was stirred at 20 °C for 20 h. The resulting white precipitate was filtered and washed with cold ether (100 mL), dried under vacuum to give 2,2,2-trifluoro-*N*-[2-({2-[(trifluoroacetyl)amino]ethyl}amino)ethyl]acetamide (17.26 g, 61%), ¹H NMR [(CD₃)₂SO] δ 7.26 (br, 2 H, 2 × CONH), 3.43 (br s, 4 H, 2 × CH₂), 2.86 (t, *J* = 5.8 Hz, 4 H, 2 × CH₂); ¹³C NMR [(CD₃)₂SO] δ 157.7 (q, *J* = 37 Hz), 115.8 (q, *J* = 288 Hz), 47.3 (2), 39.3 (2).

Di-*tert* butyldicarbonate (8.26 g, 37.8 mmol) was added to a solution of acetamide (10.15 g, 34.4 mmol) in THF (100 mL) at 0 °C and the mixture was stirred at 20 °C for 20 h. Saturated aqueous NH₄Cl (80 mL) added and the mixture stirred at 20 °C for 5 h. The mixture was extracted with DCM (3 × 50 mL), dried, and the solvent evaporated to give *tert*-butyl bis{2-[(trifluoroacetyl)amino]ethyl}carbamate (13.5 g, 100 %), ¹H NMR [(CD₃)₂SO] δ 9.47 (br, 1 H, CONH), 9.40 (br, 1 H, CONH), 3.30 (m, 8 H, 4 × CH₂), 1.38 [s, 9H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 156.4 (q, *J* = 36 Hz), 154.7, 115.8 (q, *J* = 288 Hz), 78.9, 45.4, 45.0, 37.7, 37.4, 27.7 (3).

Conc. ammonia (50 mL) was added to a solution of carbamate (14.0 g, 35.5 mmol) in MeOH (100 mL) and heated at reflux temperature for 20 hr. The solvent was evaporated to give diamine **33** as a yellow foam, ¹H NMR [(CD₃)₂SO] δ 3.39 (t, *J* = 6.4 Hz, 4 H, 2 × CH₂), 2.94 (t, *J* = 6.4 Hz, 4 H, 2 × CH₂), 1.42 [s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 154.9, 79.9, 45.1 (2), 37.4 (2), 27.9 (3).

Di-*tert*-butyl 2-aminoethyl{2-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]ethyl}dicarbamate (35).

A solution of chloride **3** (1.0 g, 5.5 mmol), diamine **33** (4.47 g, 22.0 mmol), and Et₃N (2.24 g, 22 mmol) in DME (20 mL) was heated at 90 °C for 3 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–4%) of MeOH/DCM to give (i) *tert*-butyl bis{2-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]ethyl}carbamate (0.34 g, 25%), ¹H NMR δ 8.17 (br d, *J* = 8.5 Hz, 2 H, H-8), 7.71–7.62 (m, 2 H, H-6), 7.52 (br d, *J* = 8.3 Hz, 2 H, H-8), 7.26–7.22 (m, 2 H, H-7), 6.15 (br s, 1 H, NH), 5.95 (br s, 1 H, NH), 3.71 (br q, *J* = 5.8 Hz, 4 H, 2 × CH₂), 3.37 (br s, 4 H, 2 × CH₂), 1.50 [s, 9 H, C(CH₃)₃]; ¹³C NMR δ 158.9, 156.3 (2), 148.6 (2), 135.5 (2), 130.9 (2), 126.4 (2), 124.9 (2), 120.3 (2), 80.7, 47.7 (2), 40.9 (2), 28.4 (3); and (ii) crude *tert*-butyl 2-aminoethyl{2-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]ethyl}carbamate **34** (1.38 g, 72 %) as a yellow foam.

Di-*tert*-butyldicarbonate (2.7 g, 12.4 mmol) was added to a solution of carbamate **34** (1.38 g, 4.0 mmol) in THF (50 mL) and the solution stirred at 20 °C for 36 h. Water (100 mL) was added and the mixture stirred at 20 °C for 1 h. The mixture was extracted with DCM (3 × 50 mL), the organic fraction dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–7%) MeOH/DCM, to give carbamate **35** (0.94 g, 52 %) as a yellow powder, mp (DCM/hexane) 160–163 °C; ¹H NMR [(CD₃)₂SO] δ 8.14 (dd, *J* = 8.3, 0.7 Hz, 1 H, H-8), 8.00 and 7.92 (2 × br s, 1 H, CONH), 7.79 (dd, *J* = 7.5, 1.2 Hz, 1 H, H-6), 7.58 (br d, *J* = 7.5 Hz, 1 H, H-5), 7.34 (dd, *J* = 7.7, 1.2 Hz, 1 H, H-7), 6.81 (br s, 1 H, NH), 3.43–3.47 (m, 2 H, CH₂), 3.37 (m, 2 H, CH₂), 3.22 (t, *J* = 6.2 Hz, 2 H, CH₂), 3.03–3.07 (m, 2 H, CH₂), 1.34 [s, 9 H, C(CH₃)₃], 1.34 and 1.27 [2 × s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 158.9, 155.5, 154.7, 148.2, 135.7, 130.1, 126.0, 124.6, 119.8, 78.4, 77.5, 47.0, 46.2, 38.5, 38.1, 28.1(3), 27.8 (3). Anal. calcd for C₂₁H₃₂N₆O₅ C, 56.2; H, 7.2; N, 18.7; found C, 56.5; H, 7.5; N, 18.8%.

Di-*tert*-butyl 2-aminoethyl{2-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}dicarbamate (36).

MCPBA (247 mg, 1.0 mmol) was added to a solution of 1-oxide **35** (300 mg, 0.67 mmol) in DCM (10 mL) and the mixture was stirred at 20 °C for 16 h. The mixture was partitioned between dil. aqueous NH₃ (50 mL) and DCM (50 mL) and the aqueous fraction extracted with DCM (3 × 30 mL). The combined organic fraction was dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–2%) of MeOH/DCM, to give (i) starting material (188 mg, 62%) and (ii) 1,4-dioxide **36** (122 mg 39%), mp (DCM/hexane) 128–134 °C; ¹H NMR [(CD₃)₂SO] δ 8.39 and 8.32 (2 × br s, 1 H, CONH), 8.21 (dd, *J* = 8.7, 0.7 Hz, 1 H, H-8), 8.14 (t, *J* = 8.0 Hz, 1 H, H-5), 7.94 (t, *J* = 7.6 Hz, 1 H, H-6), 7.57 (t, *J* = 7.9 Hz, 1 H, H-7), 6.77 (br s, 1 H, NH), 3.50–3.54 (m, 2 H, CH₂), 3.41–3.44 (m, 2 H, CH₂), 3.20 (t, *J* = 6.5 Hz, 2 H, CH₂), 3.03 (br q, *J* = 5.6 Hz, 2 H, CH₂), 1.33 [s, 9 H, C(CH₃)₃], 1.33 and 1.26 [2 × s, 9 H, C(CH₃)₃]; ¹³C NMR [(CD₃)₂SO] δ 155.5, 154.6, 149.8, 138.1, 135.5, 129.9, 127.0, 121.0, 116.81, 78.6, 77.4, 46.8, 46.1, 38.5, 38.0, 28.8, 27.7; HRMS calcd for C₂₁H₃₃H₆O₆ (M⁺) *m/z* 465.2462, found 465.2456.

N*-[2-({2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}amino)ethyl]-4-*acridinecarboxamide (37).** A solution of carbamate **36** (252 mg, 0.54 mmol) in HClsaturated MeOH (10 mL) was stirred at 20 °C for 24 h. The solvent was evaporated and the residue partitioned between aqueous NH₃ (20 mL) and DCM (50 mL). The

5 aqueous fraction was extracted with DCM (5 × 20 mL) and the combined organic extracts dried. The solvent was evaporated to give *N*¹-(2-aminoethyl)-*N*²-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,2-ethanediamine (109 mg, 76%), ¹H NMR δ 8.33 (d, *J* = 8.7 Hz, 1 H, H-8), 8.30 (d, *J* = 8.8 Hz, 1 H, H-5), 7.87 (ddd, *J* = 8.5, 7.1, 1.0 Hz, 1 H, H-6), 7.50 (ddd, *J* = 8.4, 7.1, 1.2 Hz, 1 H, H-7), 3.70 (t, *J* = 5.9 Hz, 2 H, CH₂), 2.98 (t, *J* = 5.9 Hz, 2 H, CH₂), 2.84 (t, *J* = 5.6 Hz, 2 H, CH₂), 2.74 (t, *J* = 5.6 Hz, 2 H, CH₂), 2 × NH and NH₂ not observed.

A solution of *N*¹-(2-aminoethyl)-*N*²-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,2-ethanediamine (96 mg, 0.36 mmol) and 4-(1*H*-imidazol-1-ylcarbonyl)acridine (119 mg, 0.43 mmol) in DMF (5 mL) was stirred at 20 °C for 5 h. The solvent was

15 evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–7%) MeOH/DCM, to give compound **37** (168 mg, 99%) as a red solid, mp (DCM/hexane) 151–154 °C; ¹H NMR δ 11.98 (t, *J* = 5.3 Hz, 1 H, CONH), 8.82 (dd, *J* = 8.2, 1.4 Hz, 1 H, ArH), 8.81 (s, 1 H, ArH), 8.08–8.17 (m, 4 H, 4 × ArH), 7.97 (d, *J* = 8.3 Hz, 1 H, ArH), 7.75–7.82 (m, 2 H, 2 × ArH), 7.60 (dd, *J* = 8.2, 7.2 Hz, 1 H, ArH), 7.54 (ddd, *J* = 8.3, 7.3, 0.9 Hz, 1 H, ArH), 7.40 (ddd, *J* = 8.6, 7.2, 1.2 Hz, 1 H, ArH), 3.86 (br q, *J* = 5.8 Hz, 2 H, CH₂), 3.70 (t, *J* = 5.8 Hz, 2 H, CH₂), 3.17 (br q, *J* = 6.3 Hz, 4 H, 2 × CH₂), 2 × NH not observed; ¹³C NMR δ 166.5, 149.7, 147.4, 146.2, 138.0, 137.7, 135.6, 135.3, 132.5, 131.4, 130.2, 128.9, 128.0, 126.9 (2), 126.7, 126.3, 125.9, 125.4, 121.5, 117.2, 48.6, 47.8, 40.6, 39.3; HRMS (FAB⁺) calcd for C₂₅H₂₄N₇O₃ (MH⁺) *m/z* 470.1941 found 470.1934. Anal. calcd for C₂₅H₂₃N₇O₃·1½H₂O: C, 60.5; H, 5.3; N, 19.8; found C, 60.5; H, 5.0; N, 20.0%.

Example N***N*-{3-[{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-****yl)amino]propyl}(methyl)amino]propyl}-4-acridinecarboxamide (41).****2,2,2-Trifluoro-*N*-[3-(methyl{3-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]acetamide (38).** A solution of chloride **3** (2.07 g,11.4 mmol), *N*¹-(3-aminopropyl)-*N*¹-methyl-1,3-propanediamine (3.31 g, 22.8 mmol)

and Et₃N (3.2 mL, 22.8 mmol) in DCM (200 mL) was stirred at 20 °C for 2 d. The solvent was evaporated and the residue dissolved in MeCN (150 mL). Ethyl trifluoroacetate (5.4 mL, 45.6 mmol) and water (0.8 mL, 45.6 mmol) added and the solution heated at reflux temperature for 16 h. The solvent was evaporated, and the residue purified by chromatography, eluting with a gradient (0–1%) of Et₃N/(0–10%) MeOH/DCM, followed by further chromatography, eluting with 10% MeOH/DCM, to give 1-oxide **38** (1.89 g, 43%) as a yellow solid, mp (DCM) 111–115 °C; ¹H NMR δ 9.04 (br s, 1 H, NH), 8.25 (dd, *J* = 8.7, 1.4 Hz, 1 H, H-8'), 7.70 (ddd, *J* = 8.4, 7.1, 1.4 Hz, 1 H, H-6'), 7.57 (d, *J* = 8.4 Hz, 1 H, H-5'), 7.29 (ddd, *J* = 8.7, 7.1, 1.1 Hz, 1 H, H-7'), 6.17 (br s, 1 H, NH), 3.58 (dd, *J* = 6.6, 5.8 Hz, 2 H, CH₂N), 3.49 (br t, *J* = 6.0 Hz, 2 H, CH₂N), 2.52–2.58 (m, 4 H, 2 × CH₂N), 2.27 (s, 3 H, NCH₃), 1.84–1.90 (m, 2 H, CH₂), 1.75–1.82 (m, 2 H, CH₂); ¹³C NMR δ 158.9, 157.3 (q, *J* = 36 Hz), 148.8, 135.6, 130.8, 126.4, 124.9, 120.4, 116.1 (q, *J* = 288 Hz), 57.1, 56.4, 41.3, 40.3 (2), 26.3, 24.4; MS (FAB⁺) *m/z* 387 (MH⁺, 100%), 371 (8), 338 (30); HRMS (FAB⁺) calcd for C₁₆H₂₂F₃N₆O₂ (MH⁺) *m/z* 387.1756, found 387.1765. Anal. calcd for C₁₆H₂₁F₃N₆O₂·½MeOH: C, 49.2; H, 5.8; N, 20.9; found: C, 49.1; H, 5.5; N, 20.7%.

***N*-{3-[(3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl)(methylamino)propyl]-2,2,2-trifluoroacetamide (39).**

Trifluoroacetic anhydride (4.13 mL, 29.2 mmol) was added to a stirred solution of 1-oxide **38** (1.13 g, 2.92 mmol) in CHCl₃ (50 mL) and the solution stirred at 20 °C for 30 min. The solution was cooled to –10 °C and 70% H₂O₂ (2 mL) (CAUTION) added dropwise. The solution was stirred at 20 °C for 30 d, partitioned between CHCl₃ (50 mL) and sat. aqueous KHCO₃ (50 mL). The aqueous fraction was extracted with CHCl₃ (3 × 30 mL), the combined organic fraction dried and the solvent evaporated (CAUTION: safety shield). The residue was purified by chromatography, eluting with 10% MeOH/DCM, to give (i) starting material **38** (275 mg, 24%) and (ii) 1,4-dioxide **39** (319 mg, 27%) as a red gum, ¹H NMR [(CD₃)₂SO] δ 9.44 (br s, 1 H, NH), 8.45 (t, *J* = 5.9 Hz, 1 H, NH), 8.20 (d, *J* = 8.8 Hz, 1 H, H-8'), 8.12 (d, *J* = 8.6 Hz, 1 H, H-5'), 7.93 (ddd, *J* = 8.6, 7.1, 1.2 Hz, 1 H, H-6'), 7.57 (ddd, *J* = 8.8, 7.1, 1.3 Hz, 1 H, H-7'), 3.42–3.47 (m, 2 H, CH₂N), 3.21–3.25 (m, 2 H, CH₂N), 2.39 (t, *J* = 6.7 Hz, 2 H, CH₂N), 2.32 (t, *J* = 6.9 Hz, 2 H, CH₂N), 2.16 (s, 3 H, NCH₃), 1.72–1.80 (m, 2 H, CH₂), 1.61–1.68 (m, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 155.9 (q, *J* = 36 Hz), 149.7,

138.1, 135.4, 129.8, 126.7, 121.0, 116.7, 115.9 (q, $J = 288$ Hz), 54.9, 54.6, 41.4, 39.5, 37.6, 25.9, 25.8; MS (FAB⁺) m/z 403 (MH⁺, 25%), 387 (5); HRMS (FAB⁺) calcd for C₁₆H₂₂F₃N₆O₃ (MH⁺) m/z 403.1706, found 403.1695.

- 5 ***N*-{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}(methylamino)propyl}-4-acridinecarboxamide (41).** A solution of trifluoroacetamide **39** (175 mg, 0.44 mmol) and NH₄OH (5 mL) in MeOH (20 mL) was stirred at 30 °C for 4 h. The solvent was evaporated and the residue dried to give *N*¹-(3-aminopropyl)-*N*³-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-*N*¹-methyl-1,3-propanediamine (**40**) as a red gum, ¹H NMR [(CD₃)₂SO] δ 8.43 (br s, 1 H, NH), 8.21 (d, $J = 8.5$ Hz, 1 H, H-8'), 8.13 (d, $J = 8.4$ Hz, 1 H, H-5'), 7.94 (ddd, $J = 8.4, 7.1, 1.2$ Hz, 1 H, H-6'), 7.75 (br s, 2 H, NH₂), 7.57 (ddd, $J = 8.7, 7.2, 1.3$ Hz, 1 H, H-7'), 3.45 (t, $J = 6.8$ Hz, 2 H, CH₂N), 3.20–3.25 (m, 2 H, CH₂N), 2.88 (dd, $J = 7.4, 7.2$ Hz, 2 H, CH₂N), 2.40–2.46 (m, 2 H, CH₂N), 2.20 (s, 3 H, NCH₃), 1.77–1.83 (m, 2 H, CH₂), 1.68–1.75 (m, 2 H, CH₂); MS (FAB⁺) m/z 307 (MH⁺, 2%), 291 (5); HRMS (FAB⁺) calcd for C₁₄H₂₃N₆O₃ (MH⁺) m/z 307.1883, found 307.1883. The amine **40** was dissolved in DCM (5 mL) and added to a stirred solution of 4-(1*H*-imidazol-1-ylcarbonyl)acridine (125 mg, 0.46 mmol) in THF (20 mL) and the solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by
- 20 chromatography, eluting with a gradient (0–1%) of Et₃N/(0–15%) MeOH/DCM, to give compound **41** (146 mg, 66%) as a red solid, mp (EtOAc/DCM) 169–171 °C; ¹H NMR [(CD₃)₂SO] δ 11.41 (t, $J = 5.3$ Hz, 1 H, CONH), 9.31 (s, 1 H, H-9), 8.69 (dd, $J = 7.0, 1.4$ Hz, 1 H, H-3), 8.43 (t, $J = 5.6$ Hz, 1 H, NH), 8.38 (d, $J = 7.4$ Hz, 1 H, H-1), 8.32 (d, $J = 8.8$ Hz, 1 H, H-5), 8.21 (d, $J = 8.4$ Hz, 1 H, H-8), 8.16 (d, $J = 8.7$ Hz, 1 H, H-8'), 8.09 (d, $J = 8.7$ Hz, 1 H, H-5'), 7.96 (ddd, $J = 8.7, 7.1, 1.1$ Hz, 1 H, H-6'), 7.91 (dd, $J = 8.8, 7.5$ Hz, 1 H, H-6), 7.74 (dd, $J = 7.4, 7.0$ Hz, 1 H, H-2), 7.69 (br dd, $J = 8.7, 7.1$ Hz, 1 H, H-7'), 7.55 (dd, $J = 8.4, 7.5$ Hz, 1 H, H-7), 3.60–3.65 (m, 2 H, CH₂N), 3.42–3.48 (m, 2 H, CH₂N), 3.39 (s, 3 H, NCH₃), 3.00–3.08 (m, 2 H, CH₂N), 2.60–2.68 (m, 2 H, CH₂N), 2.02–2.08 (m, 2 H, CH₂), 1.92–1.98 (m, 2 H, CH₂); MS (FAB⁺) m/z 512 (MH⁺, 25%), 496 (10); HRMS (FAB⁺) calcd for C₂₈H₃₀N₇O₃ (MH⁺) m/z 512.2410, found 512.2424.
- 30

Example O.

***N*-{3-[[3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl](methylamino)propyl]-8-quinolinecarboxamide (42).**

A solution of 8-quinolinecarboxylic acid (90 mg, 0.5 mmol) and CDI (97 mg, 0.6 mmol) in DMF (5 mL) was stirred at 55 °C for 24 h. The solution was diluted with
 5 dry benzene (10 mL), Sephadex LH-20 (300 mg) was added and the mixture stirred at 20 °C for 1 h. The mixture was filtered and the solvent evaporated. The residue was dissolved in dry THF (5 mL) and a solution of (40) (80 mg, 0.25 mmol) in THF (5 mL) added, and the solution stirred at 20 °C for 70 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous
 10 NH₃/(0–8%) MeOH/DCM, to give compound 42 (110 mg, 91%) as a red powder, mp (DCM/pet. ether) 119–121 °C; ¹H NMR δ 11.39 (br s, 1 H, CONH), 8.96 (dd, *J* = 4.3, 1.8 Hz, 1 H, ArH), 8.74 (dd, *J* = 7.3, 1.5 Hz, 1 H, ArH), 8.34 (dd, *J* = 8.8, 1.8 Hz, 1 H, ArH), 8.25 (d, *J* = 8.3 Hz, 1 H, ArH), 8.17 (d, *J* = 8.6 Hz, 1 H, ArH), 7.98 (br s, 1 H, NH), 7.92 (dd, *J* = 8.1, 1.5 Hz, 1 H, ArH), 7.78 (dd, *J* = 8.1, 1.1 Hz, 1 H, ArH), 7.62
 15 (t, *J* = 7.7 Hz, 1 H, ArH), 7.48 (dd, *J* = 8.3, 1 H, 4.0 Hz), 7.43 (dd, *J* = 7.9, 1.0 Hz, 1 H, ArH), 3.68–3.73 (m, 4 H, 2 × CH₂), 3.05 (br m, 4 H, 2 × CH₂), 2.67 (s, 3 H, CH₃), 2.25–2.17 (m, 4 H, 2 × CH₂); ¹³C NMR δ 166.4, 149.7, 149.6, 145.4, 138.2, 137.7, 135.6, 133.6, 132.0, 130.3, 128.5, 128.4, 127.1, 126.4, 121.5, 121.0, 117.3, 54.7, 54.5, 40.6, 39.4, 37.2, 25.5, 24.5; MS (FAB⁺) *m/z* 462 (MH⁺, 25%), 446 (5); HRMS calcd for C₂₄H₂₈N₇O₃ (MH⁺) *m/z* 462.2254, found 462.2249. Anal. calcd for C₂₄H₂₇N₇O₃: C, 62.5; H, 5.9; N, 21.2; found: C, 62.1; H, 6.0; N, 21.2%.

Example P.

***N*-{3-[[3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl](methylamino)propyl]-2-(4-pyridyl)-8-quinolinecarboxamide (43).** A solution of 2-(4-pyridyl)-
 25 quinoline-8-carboxylic acid (160 mg, 0.62 mmol) and CDI (150 mg, 0.92 mmol) in DMF (10 mL) was stirred at 55 °C for 24 h. The solution was cooled to 20 °C, diluted with dry benzene (15 mL), Sephadex LH-20 (300 mg) was added and the mixture stirred at 20 °C for 1 h. The mixture was filtered and the solvent evaporated. The
 30 residue was dissolved in dry THF (5 mL) and a solution of (39) (90 mg, 0.33 mmol) in THF (5 mL) added, and the solution stirred at 20 °C for 4 days. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–8%) MeOH/DCM, to give compound 43 (160 mg, 94%) as a

red powder, mp (DCM/pet. ether) 179–181 °C; ^1H NMR δ 11.08 (br s, 1 H, CONH), 8.86 (dd, J = 4.5, 1.6 Hz, 2 H, ArH), 8.78 (dd, J = 7.4, 1.5 Hz, 1 H, ArH), 8.37 (d, J = 8.6 Hz, 1 H, ArH), 8.21 (d, J = 8.6 Hz, 1 H, ArH), 8.10 (d, J = 8.6 Hz, 1 H, ArH), 7.95 (dd, J = 8.2, 1.4 Hz, 1 H, ArH), 7.92–7.90 (m, 4 H, NH, 3 \times ArH), 7.98 (ddd, J = 8.6, 7.5, 1.3 Hz, 1 H, ArH), 7.66 (t, J = 7.7 Hz, 1 H, ArH) 7.40 (ddd, J = 8.6, 7.2, 1.2 Hz, 1 H, ArH), 3.74 (br q, J = 6.4 Hz, 2 H, CH₂), 3.61 (br m, 2 H, CH₂), 2.85 (br m, 2 H, CH₂), 2.81 (br m, 2 H, CH₂), 2.45 (s, 3 H, CH₃), 2.17 (br q, J = 7.2 Hz, 2 H, CH₂), 1.98 (br m, 2 H, CH₂); ^{13}C NMR δ 166.1, 154.5, 150.9, 149.7, 146.2, 145.3, 139.0, 138.2, 135.5, 134.4, 131.5, 130.2, 129.4, 127.9, 127.2, 126.9, 121.7, 121.5, 118.7, 117.2, 55.3, 55.2, 41.0, 40.1, 37.7, 26.6, 24.7; HRMS (FAB⁺) calcd for C₂₉H₃₁N₈O₃ (MH⁺) m/z 539.2519, found 539.2527. Anal. calcd for C₂₉H₃₀N₈O₃·½H₂O: C, 64.7; H, 5.6; N, 20.8; found: C, 64.1; H, 5.7; N, 20.6%.

Example Q.

N-{3-[[3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl](methyl)amino]-propyl}-5-methyl-4-acridinecarboxamide (44). A solution of 5-methylacridine-4-carboxylic acid (0.13 g, 0.55 mmol) and CDI (0.21 g, 1.3 mmol) in DMF (5 mL) was stirred at 55 °C for 24 h. The solution was diluted with dry benzene (10 mL), Sephadex LH-20 (300 mg) was added and the mixture stirred at 20 °C for 1 h. The mixture was filtered and the solvent evaporated. The residue was dissolved in dry THF (5 mL) and a solution of **40** (80 mg, 0.27 mmol) in THF (5 mL) added, and the solution stirred at 20 °C for 70 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–8%) MeOH/DCM, to give compound **44** (0.13 g, 88%) as a red powder, mp (DCM/pet. ether) 158–162 °C; ^1H NMR δ 12.08 (br s, 1 H, CONH), 8.83 (d, J = 6.9 Hz, 1 H, ArH), 8.76 (s, 1 H, NH), 8.06 (t, J = 8.9 Hz, 2 H, ArH), 7.97 (br d, J = 8.4 Hz, 2 H, ArH), 7.83 (d, J = 8.4 Hz, 1 H, ArH), 7.66 (d, J = 6.7 Hz, 1 H, ArH), 7.56–7.63 (m, 2 H, ArH), 7.46 (dd, J = 7.6, 6.5 Hz, 1 H, ArH), 7.30 (d, J = 7.9 Hz, 1 H, ArH), 3.77 (br q, J = 6.3 Hz, 2 H, CH₂), 4.83 (br m, 2 H, CH₂), 3.08 (br m, 4 H, 2 \times CH₂), 2.83 (s, 3 H, CH₃), 2.67 (br s, 3 H, CH₃), 2.31 (br m, 2 H, CH₂), 2.15 (br m, 2 H, CH₂); ^{13}C NMR δ 166.5, 149.6, 146.9, 145.1, 137.9, 137.9, 135.8, 135.3, 135.1, 132.4, 131.2, 130.0, 127.9, 126.8, 126.4, 126.3, 126.2, 125.8, 125.2, 121.3, 117.0, 55.1, 54.5, 40.5, 39.2, 37.4, 26.1, 24.5, 19.0; HRMS (FAB⁺) calcd for C₂₉H₃₂N₇O₃ (MH⁺) m/z

526.2593, found 526.2582. Anal. calcd for $C_{29}H_{31}N_7O_3 \cdot 0.5H_2O$: C, 65.2; H, 6.0; N, 18.3; found: C, 65.0; H, 5.8; N, 18.1%.

Example R.

- 5 ***N*-{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}(methylamino)-propyl}-9-methyl-4-phenazinecarboxamide (45).** A solution of 9-methylphenazine-4-carboxylic acid (130 mg, 0.53 mmol) and CDI (100 mg, 0.61 mmol) in DMF (5 mL) was stirred at 55 °C for 6 h. The solution was cooled to 20 °C, diluted with dry benzene (10 mL), Sephadex LH-20 (300 mg) was added and the mixture stirred at 20 °C for 1 h. The mixture was filtered and the solvent evaporated. The residue was dissolved in dry THF (5 mL) and a solution of **40** (80 mg, 0.26 mmol) in THF (5 mL) added, and the solution stirred at 20 °C for 24 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH_3 /(0–8%) MeOH/DCM, to give compound **45** (130 mg, 90%) as a red powder, mp (DCM/pet. ether) 138–142 °C; 1H NMR δ 11.23 (br s, 1 H, CONH), 8.84 (d, J = 6.6 Hz, 1 H, ArH), 8.29 (d, J = 7.6 Hz, 1 H, ArH), 8.07 (d, J = 8.5 Hz, 1 H, ArH), 8.04 (d, J = 8.5 Hz, 1 H, ArH), 7.98 (d, J = 8.6 Hz, 1 H, ArH), 7.85 (t, J = 7.8 Hz, 1 H, ArH), 7.78–7.71 (m, 3 H, ArH, NH), 6.48 (t, J = 7.6 Hz, 1 H, ArH), 7.31 (t, J = 7.7 Hz, 1 H, ArH), 3.78–3.71 (m, 4 H, $2 \times CH_2$), 3.15 (br m, 4 H, $2 \times CH_2$), 2.88 (s, 3 H, CH_3), 2.73 (br s, 3 H, CH_3), 2.32, (br m, 2 H, CH_2), 2.21 (br m, 2 H, CH_2); ^{13}C NMR δ 165.6, 149.6, 143.2, 142.9, 140.7, 139.4, 137.9, 136.4, 135.4, 135.1, 133.7, 131.3, 131.2, 130.1, 129.7, 128.5, 127.7, 127.0, 121.3, 116.9, 54.9, 54.2, 40.2, 38.9, 37.3, 25.7, 24.3, 18.1. Anal. calcd for $C_{28}H_{30}N_8O_3$: C, 63.9; H, 5.9; N, 21.3; HRMS (FAB $^+$) calcd for $(C_{28}H_{31}N_8O_3) (MH^+)$ m/z 527.2519 found 527.2533. Anal. calcd for $C_{28}H_{30}N_8O_3 \cdot 1.75H_2O$: C, 60.3; H, 6.0; N, 20.1; found: C, 60.3; H, 5.6; N, 19.6%.

Example S.

- N*-{3-[(3-[(7-(2-Methoxyethoxy)-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl)(methylamino)propyl]-4-acridinecarboxamide (55).**
- 30 **3-Amino-1,2,4-benzotriazin-7-ol 1-oxide (46).** A mixture of 4-amino-3-nitrophenol (5.0 g, 32.4 mmol) and cyanamide (8.2 g, 194.6 mmol) was heated at 100 °C for 10 min. The resulted solution was cooled to 20 °C and c.HCl (15 mL) was added dropwise, and the mixture was heated at 100 °C for 1.5 h, cooled to 20 °C. A solution

of 30% NaOH (40 mL) was then added and heated at 100 °C for 1 h. The reaction mixture was cooled to 20 °C, diluted with water (20 mL), and the precipitate was filtered, washed with water (100 mL), diethyl ether (100 mL), and dried to give amine **46** (5.45 g, 97%) as a yellow powder, mp > 300 °C [lit. (Friebe et. al. US Patent 5,856,325, Jan 5, 1999) mp (HOAc) >270 °C]; ¹H NMR [(CD₃)₂SO] δ 10.37 (br s, 1 H, OH), 7.48 (dd, *J* = 7.7, 2.6 Hz, 1 H, H-6), 7.40–7.37 (m, 2 H, H-5, H-8), 6.96 (br s, 2 H, NH₂).

7-(2-Methoxyethoxy)-1,2,4-benzotriazin-3-amine 1-oxide (47). A mixture of **46** (1.00 g, 5.8 mmol), dry K₂CO₃ (2.40 g, 17.4 mmol) and 2-bromoethyl methyl ether (2.42 g, 17.4 mmol) in DMF (20 mL) was heated at 80 °C for 2 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–3%) of MeOH/DCM, to give compound **47** (1.06 g, 77 %) as a yellow powder, mp (DCM/pet. ether) 201–203 °C; ¹H NMR [(CD₃)₂SO] δ 8.07(d, *J* = 9.5 Hz, 1 H, H-5), 7.82 (br s, 2 H, NH₂), 7.76 (dd, *J* = 9.5, 2.6 Hz, 1 H, H-6), 7.50 (d, *J* = 2.6 Hz, 1 H, H-8), 4.26, (t, *J* = 4.3 Hz, 2 H, CH₂), 3.72 (t, *J* = 4.3 Hz, 2 H, CH₂), 3.33 (s, 3 H, OCH₃). Anal. calcd for C₁₀H₁₂N₄O₅: C, 50.8; H, 5.1; N, 23.7; found: C, 51.1; H, 5.0; N, 23.7%.

3-Hydroxy-7-(2-methoxyethoxy)-1,2,4-benzotriazine 1-oxide (48). A suspension of **47** (1.00 g, 4.2 mmol) in 2 N HCl (32 mL) was cooled to 5 °C and a solution of NaNO₂ (0.58 g, 8.5 mmol) in water (1.5 mL) was added over 1 h. More NaNO₂ (0.58 g, 8.5 mmol) in water (1.5 mL) was added and the suspension stirred 72 h at 20 °C. The precipitate was filtered and washed with water. The solid was dissolved in 5% aqueous NH₃ and filtered. The filtrate was acidified with conc. HCl to give a precipitate which was filtered dried and purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM to give compound **48** (0.68 g, 68 %) as a yellow solid, mp (DCM/pet.ether) 190–192 °C; ¹H NMR [(CD₃)₂SO] δ 12.52 (br, 1 H, OH), 7.69 (br s, 1 H, H-8), 7.53 (dd, *J* = 8.8, 2.8 Hz, 1 H, H-6), 7.33 (d, *J* = 8.8 Hz, 1 H, H-5), 4.19 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.68 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.33 (s, 3 H, OCH₃); ¹³C NMR [(CD₃)₂SO] δ 154.6, 152.9, 131.8, 129.5, 127.4, 117.8, 101.8, 70.0, 67.9, 58.1. Anal. calcd for C₁₀H₁₁N₃O₄: C, 50.6; H, 4.2; N, 17.7; found: C, 50.5; H, 4.7; N, 17.7.

3-Chloro-7-(2-methoxyethoxy)-1,2,4-benzotriazine 1-oxide (49). A mixture of **48** (1.00 g, 4.3 mmol) in POCl₃ (8 mL) was refluxed for 2 h. Excess reagent was evaporated under vacuum, and ice cold water (50 mL) was added to the residue, then solid Na₂CO₃ (1.0 g) was added portionwise. The resulting precipitate was filtered and purified by chromatography, eluting with a gradient (50–100 %) of DCM/pet. ether, to give compound **49** (0.90 g, 83%) as a pale yellow solid, mp (DCM/pet. ether) 121–125 °C; ¹H NMR [(CD₃)₂SO] δ 8.00 (d, *J* = 9.2 Hz, 1 H, H-5), 7.81 (dd, *J* = 9.2, 2.9 Hz, 1 H, H-6), 7.68 (d, *J* = 2.8 Hz, 1 H, H-8), 4.35 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.74 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.33 (s, 3 H, OCH₃). Anal. calcd for C₁₀H₁₀ClN₃O₃: C, 47.0; H, 3.9; N, 16.4, Cl, 13.9; found: C, 46.9; H, 4.3; N, 16.4; Cl, 13.7.

N¹-(3-Aminopropyl)-N³-[7-(2-methoxyethoxy)-1-oxido-1,2,4-benzotriazin-3-yl]-N¹-methyl-1,3-propanediamine (51). A solution of chloride **49** (0.90 g, 3.5 mmol) *tert*-butyl 3-[(3-aminopropyl)(methyl)amino]propylcarbamate (**50**) (1.60 g, 5.25 mmol) and Et₃N (4 ml) in DME (20 mL) was heated to 90 °C for 4 h. The solvent was evaporated, the residue was dissolved in MeOH (10 mL), and treated with methanolic HCl (100 mL). The reaction mixture was stirred at 20 °C for 20 h, the solvent evaporated and the residue partitioned between DCM and dil. aqueous NH₃. The aqueous layer was extracted with DCM (4 × 25 mL), the combined extracts dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–10%) MeOH/DCM, to give compound **51** (1.25 g, 98%) as a yellow solid, ¹H NMR [(CD₃)₂SO] δ 7.68 (br s, 1 H, NH), 7.55–7.52 (m, 1 H, ArH), 7.50–7.47 (m, 2 H, ArH), 4.20 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.70 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.34 (br m, 2 H, CH₂), 3.32 (s, 3 H, OCH₃), 2.54 (br t, *J* = 6.1 Hz, 2 H, CH₂), 2.35 (t, *J* = 6.9 Hz, 2 H, CH₂), 2.31 (t, *J* = 7.2 Hz, 2 H, CH₂), 2.13 (s, 3 H, NCH₃), 1.70 (br quin, *J* = 6.9 Hz, 2 H, CH₂), 1.47 (br quin, *J* = 7.0 Hz, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 158.3, 155.3, 144.5, 129.8, 138.3, 127.5, 98.9, 70.0, 67.7, 58.1, 55.0, 54.9, 41.8, 39.9, 39.1, 30.7, 26.2; HRMS (FAB⁺) calcd for C₁₇H₂₉N₆O₃ (MH⁺) *m/z* 365.2301, found 365.2311.

2,2,2-Trifluoro-N-{3-[(3-{[7-(2-methoxyethoxy)-1,4-dioxido-1,2,4-benzotriazin-3-yl]amino}propyl)(methyl)amino]propyl}acetamide (52). Ethyl trifluoroacetate (1.2 mL, 9.8 mmol) and H₂O (0.17 mL, 9.8 mmol) were added to a solution of **51** (1.19 g,

3.3 mmol) in CH₃CN and the reaction mixture was heated at reflux for 18 h. The solvent was evaporated and the residue partitioned between aqueous Na₂CO₃ solution and DCM. The aqueous layer was extracted with DCM, the combined organic extracts dried and the solvent evaporated. The residue was purified by chromatography,

5 eluting with a gradient (0–5%) of MeOH/DCM to give compound **52** (1.3 g, 87%) as a yellow solid, mp (DCM/pet. ether) 117–119 °C; ¹H NMR [(CD₃)₂SO] δ 9.43 (br s, 1 H, CONH), 7.66 (br t, *J* = 5.3 Hz, 1 H, NH), 7.54–7.45 (m, 3 H, ArH), 4.20 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.70 (t, *J* = 4.4 Hz, 2 H, CH₂), 3.36–3.27 (m, 2 H, CH₂), 3.30 (s, 3 H, OCH₃), 3.21 (t, *J* = 7.0 Hz, 2 H, CH₂), 2.37 (t, *J* = 6.9 Hz, 2 H, CH₂), 2.31 (t, *J* = 6.9
10 Hz, 2 H, CH₂), 2.14 (s, 3 H, NCH₃), 1.70 (br quin, *J* = 7.0 Hz, 2 H, CH₂), 1.63 (br quin, *J* = 7.0 Hz, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 158.3, 156.5 (q, *J* = 18 Hz), 155.3, 144.5, 129.8, 128.3, 127.5, 115.9 (q, *J* = 288 Hz), 98.9, 70.0, 67.7, 58.1, 54.8, 54.5, 41.5, 39.0, 37.7, 26.2, 25.8; HRMS (EI⁺) calcd for C₁₉H₂₇F₃N₆O₄ (M⁺) *m/z* 460.2046, found 460.2040. Anal. calcd for C₁₉H₂₇F₃N₆O₄ C, 49.6; H, 5.9; N, 18.3; F,
15 12.4; found: C, 49.9; H, 5.9; N, 18.2; F, 12.4%.

2,2,2-Trifluoro-*N*-{3-[(3-{[7-(2-methoxyethoxy)-1,4-dioxido-1,2,4-benzotriazin-3-yl]amino}propyl)(methylamino)propyl]acetamide (53). 70% H₂O₂ (1.05 mL, 21.7 mmol) was added dropwise to a solution of trifluoroacetic anhydride (3.0 mL, 21.7
20 mmol) in DCM (10 mL) at 5 °C. The solution was stirred at 5 °C for 10 min, 20 °C for 10 min, and then cooled to 5 °C. The solution was added dropwise to a solution of 1-oxide **52** (1.0 g, 2.2 mmol) and TFA (0.33 mL, 4.3 mmol) in DCM (50 mL). The reaction mixture was stirred at 20 °C for 18 h. The solution was partitioned between aqueous NaHCO₃ solution and DCM, the aqueous layer extracted further with DCM
25 (5 × 30 mL), the combined extracts dried, and the solvent evaporated. The residue was chromatographed, eluting with a gradient (0–5%) of MeOH/DCM to give compound **53** (0.32 g, 30%) as a red solid, mp (DCM/pet. ether) 91–94 °C; ¹H NMR [(CD₃)₂SO] δ 9.43 (br s, 1 H, CONH), 8.24 (t, *J* = 5.6 Hz, 1 H, NH), 8.05 (d, *J* = 9.5 Hz, 1 H, H-5), 7.60 (dd, *J* = 9.5, 2.7 Hz, 1 H, H-6), 7.50 (d, *J* = 2.6 Hz, 1 H, H-8), 4.26 (t, *J* = 4.3
30 Hz, 2 H, CH₂), 3.72 (t, *J* = 4.3 Hz, 2 H, CH₂), 3.41 (br q, *J* = 6.6 Hz, 2 H, CH₂), 3.33 (s, 3 H, OCH₃), 3.23 (br q, *J* = 6.3 Hz, 2 H, CH₂), 2.38 (t, *J* = 6.7 Hz, 2 H, CH₂), 2.32 (t, *J* = 6.9 Hz, 2 H, CH₂), 2.15 (s, 3 H, NCH₃), 1.75 (br quin, *J* = 6.7 Hz, 2 H, CH₂), 1.65 (br quin, *J* = 6.9 Hz, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 157.0, 156.0 (q, *J* = 6

Hz), 149.0, 134.2, 130.2, 128.2, 120.2, 115.9 (q, $J = 279$ Hz), 99.5, 69.9, 68.0, 58.1, 54.9, 54.6, 41.5, 39.5, 37.7, 25.9, 25.8; HRMS (FAB⁺) calcd for C₁₉H₂₈F₃N₆O₅ (MH⁺) m/z 477.2073, found 477.2074.

- 5 ***N*-{3-[(3-{[7-(2-Methoxyethoxy)-1,4-dioxido-1,2,4-benzotriazin-3-yl]amino}propyl)(methyl)amino]propyl}-4-acridinecarboxamide (55).** A solution of trifluoroacetamide **53** (1.55 g, 0.33 mmol) and aqueous NH₃ (8 mL) in MeOH (10 mL) was stirred at 20 °C for 18 h. The solvent was evaporated and the residue dried to give the intermediate amine **54** as a red solid. The solid was dissolved in dry THF (10 mL) and 4-(1*H*-imidazol-1-ylcarbonyl)acridine (0.18 g, 0.65 mmol) added and solution stirred at 20 °C for 72 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2 %) of aqueous NH₃/(0–5 %MeOH/DCM, to give compound **55** (150 mg, 79%) as a red solid, mp (DCM/pet. ether) 98–103 °C; ¹H NMR [(CD₃)₂SO] δ 11.32 (t, $J = 5.3$ Hz, 1 H, CONH), 9.27 (s, 1 H, NH), 8.68 (dd, $J = 7.0, 1.5$ Hz, 1 H, ArH), 8.33 (dd, $J = 8.4, 1.3$ Hz, 1 H, ArH), 8.27–8.18 (m, 3 H, ArH), 7.96–7.91 (m, 2 H, ArH), 7.72 (dd, $J = 8.2, 7.2$ Hz, 1 H, ArH), 7.66 (t, $J = 7.3$ Hz, 1 H, ArH), 7.50 (dd, $J = 9.5, 2.6$ Hz, 1 H, ArH), 7.40 (d, $J = 2.6$ Hz, 1 H, ArH), 4.23 (t, $J = 4.3$ Hz, 2 H, CH₂), 3.71 (t, $J = 4.4$ Hz, 2 H, CH₂), 3.59 (br q, $J = 6.4$ Hz, 2 H, CH₂), 3.43 (br q, $J = 6.4$ Hz, 2 H, CH₂), 2.56 (t, $J = 7.0$ Hz, 2 H, CH₂), 2.46 (t, $J = 6.7$ Hz, 2 H, CH₂), 2.23 (s, 3 H, NCH₃), 1.91 (br quin, $J = 6.8$ Hz, 2 H, CH₂), 1.78 (br quin, $J = 6.6$ Hz, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 164.6, 156.9, 148.9, 146.9, 145.4, 138.5, 134.3, 134.1, 132.6, 131.8, 130.0, 128.5, 128.3, 128.3, 128.0, 126.4, 126.3, 125.5, 125.1, 118.4, 99.4, 69.9, 68.0, 58.1, 55.2, 55.0, 41.8, 39.7, 37.4, 26.9, 25.8; HRMS (FAB⁺) calcd for C₃₁H₃₆N₇O₅ (MH⁺) m/z 586.2778, found 586.2768. Anal. calcd for C₃₁H₃₅N₇O₅: C, 63.6; H, 6.0; N, 16.7; found: C, 62.3; H, 6.1; N, 16.5%.

Example T.

- 30 ***N*-{2-[[3-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)propyl](methyl)amino]propyl}-4-acridinecarboxamide (62).**

3-Allyl-1,2,4-benzotriazine 1-oxide (56). Pd(PPh₃)₄ (640 mg, 0.55 mmol) was added to a stirred solution of chloride **3** (2.0 g, 11.0 mmol) and allyltributyltin (3.8 mL, 12.1 mmol), the solution degassed, and stirred under N₂ at reflux temperature for 6 h. The

solvent was evaporated and the residue purified by chromatography, eluting with 20% EtOAc/pet. ether to give an oil which was purified by chromatography, eluting with 5% EtOAc/DCM, to give alkene **56** (1.92 g, 93%) as a white solid, mp (EtOAc/pet. ether) 57–58 °C, ^1H NMR δ 8.45 (dd, J = 8.6, 1.4 Hz, 1 H, H-8), 8.10 (dd, J = 8.4, 1.4 Hz, 1 H, H-5), 7.94 (ddd, J = 8.4, 7.1, 1.4 Hz, 1 H, H-6), 7.70 (ddd, J = 8.6, 7.1, 1.4 Hz, 1 H, H-7), 6.15–6.24 (m, 1 H, H-2'), 5.31 (dq, J = 17.0, 1.5 Hz, 1 H, H-3'), 5.24 (dq, J = 10.1, 1.5 Hz, 1 H, H-3'), 3.80 (dq, J = 6.8, 1.5 Hz, 2 H, H-1'); ^{13}C NMR δ 165.2, 147.5, 135.6, 133.3, 132.7, 130.1, 128.8, 120.8, 118.5, 41.8. Anal. calcd for $\text{C}_{10}\text{H}_9\text{N}_3\text{O}$: C, 64.2; H, 4.85; N, 22.45; found: C, 63.85; H, 4.9; N, 22.7%.

3-(3-Hydroxypropyl)-1,2,4-benzotriazine 1-oxide (57). A solution of 9-BBN in THF (13.7 mL, 6.8 mmol) was added to a stirred solution of alkene **56** (1.07 g, 5.7 mmol) in THF (50 mL) and the solution stirred at 20 °C for 1 h. A solution of NaOH (3 M; 2.9 mL, 8.5 mmol), followed by 35% H_2O_2 (2.6 mL, 25.6 mmol) were carefully added and the mixture stirred at 20 °C for 1 h. The mixture was diluted with brine (100 mL), extracted with EtOAc (3×100 mL), the combined organic fraction dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (10–50%) of EtOAc/DCM, to give alcohol **57** (1.02 g, 87%) as a white solid, mp (EtOAc/pet. ether) 99–100 °C; ^1H NMR δ 8.46 (dd, J = 8.7, 1.0 Hz, 1 H, H-8), 7.99 (dd, J = 8.5, 1.2 Hz, 1 H, H-5), 7.93 (ddd, J = 8.5, 7.0, 1.0 Hz, 1 H, H-6), 7.70 (ddd, J = 8.7, 7.0, 1.2 Hz, 1 H, H-7), 3.80 (t, J = 6.1 Hz, 2 H, CH_2O), 3.18 (t, J = 7.3 Hz, 2 H, CH_2), 2.15–2.22 (m, 2 H, CH_2), (OH not observed); ^{13}C NMR δ 166.9, 147.3, 135.7, 133.3, 130.1, 128.6, 120.1, 62.1, 34.1, 30.5. Anal. calcd for $\text{C}_{10}\text{H}_{11}\text{N}_3\text{O}_2$: C, 58.5; H, 5.4; N, 20.5; found: C, 58.6; H, 5.5; N, 20.5%.

tert-Butyl 3-{methyl[3-(1-oxido-1,2,4-benzotriazin-3-yl)propyl]amino}propylcarbamate (58). MsCl (0.52 mL, 6.7 mmol) was added dropwise to a stirred solution of alcohol **57** (1.06 g, 5.2 mmol) and Et_3N (1.1 mL, 7.8 mmol) in DCM (50 mL) and the solution stirred at 20 °C for 1 h. The solution was diluted with DCM (50 mL), washed with water (2×30 mL), dried, and the solvent evaporated. The residue was dissolved in dry DMF (20 mL) and *tert*-butyl 3-(methylamino)propylcarbamate (Rennard et al. *Org. Lett.*, **2000**, 2, 2117–2120) (9.7 g, 51.6 mmol) added and the solution stirred at 50 °C for 3 h. The solvent was

evaporated and the residue partitioned between EtOAc (100 mL) and aqueous KHCO_3 solution (100 mL). The organic fraction was washed with water (2×50 mL), dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–10%) of MeOH/DCM, to give compound **58** (0.93 g, 48%) as a pale yellow oil, ^1H NMR δ 8.45 (dd, $J = 8.9, 1.4$ Hz, 1 H, H-8), 8.10 (br d, $J = 8.3$ Hz, 1 H, H-5), 7.93 (ddd, $J = 8.3, 7.0, 1.4$ Hz, 1 H, H-6), 7.70 (ddd, $J = 8.9, 7.0, 1.5$ Hz, 1 H, H-7), 5.38 (br s, 1 H, NH), 3.17–3.22 (m, 2 H, CH_2N), 3.07 (dd, $J = 7.7, 7.4$ Hz, 2 H, CH_2), 2.55–2.60 (m, 2 H, CH_2N), 2.49–2.53 (m, 2 H, CH_2N), 2.28 (s, 3 H, NCH_3), 2.10–2.18 (m, 2 H, CH_2), 1.68–1.73 (m, 2 H, CH_2), 1.42 [s, 9 H, $\text{C}(\text{CH}_3)_3$]; ^{13}C NMR δ 166.7, 156.1, 147.5, 135.6, 133.3, 130.0, 128.7, 120.1, 78.9, 56.7, 55.6, 41.5, 39.3, 34.9, 28.3 (3), 26.6, 25.0; MS (FAB $^+$) m/z 376 (MH^+ , 55%), 360 (5); HRMS (FAB $^+$) calcd for $\text{C}_{19}\text{H}_{30}\text{N}_5\text{O}_3$ (MH^+) m/z 376.2349, found 376.2345.

2,2,2-Trifluoro-*N*-(3-{methyl[3-(1-oxido-1,2,4-benzotriazin-3-

yl)propyl]amino}propyl)acetamide (59**).** A solution of carbamate **58** (0.51 g, 1.35 mmol) in HCl saturated MeOH (30 mL) was stirred at 50 °C for 3 h. The solvent was evaporated and the residue partitioned between dil. aqueous NH_3 (50 mL) and CHCl_3 (50 mL). The aqueous fraction was extracted with CHCl_3 (3×30 mL), the combined organic fraction dried, and the solvent evaporated. The residue was dissolved in MeCN (30 mL) and ethyl trifluoroacetate (0.24 mL, 2.03 mmol) and water (30 μL , 1.5 mmol) added. The solution was stirred at reflux temperature for 16 h, cooled to 20 °C and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–10%) of MeOH/DCM, to give amide **59** (460 mg, 92%) as a pale yellow oil, ^1H NMR [$(\text{CD}_3)_2\text{SO}$] δ 9.41 (br s, 1 H, CONH), 8.37 (d, $J = 8.6$ Hz, 1 H, H-8), 8.07 (ddd, $J = 8.3, 6.9, 1.4$ Hz, 1 H, H-6), 8.02 (dd, $J = 8.3, 1.3$ Hz, 1 H, H-5), 7.83 (ddd, $J = 8.6, 6.9, 1.3$ Hz, 1 H, H-7), 3.17–3.22 (m, 2 H, CH_2N), 2.95 (dd, $J = 7.6, 7.4$ Hz, 2 H, CH_2), 2.42 (br t, $J = 6.8$ Hz, 2 H, CH_2N), 2.33 (br t, $J = 6.7$ Hz, 2 H, CH_2N), 2.16 (s, 3 H, NCH_3), 1.92–2.00 (m, 2 H, CH_2), 1.57–1.64 (m, 2 H, CH_2); ^{13}C NMR [$(\text{CD}_3)_2\text{SO}$] δ 166.2, 156.0 (q, $J = 37$ Hz), 146.9, 136.0, 132.8, 130.4, 128.3, 119.5, 115.9 (q, $J = 288$ Hz), 56.1, 54.4, 41.4, 37.6, 34.2, 25.7, 24.8; MS (EI^+) m/z 371 (M^+ , 7%), 354 (100); HRMS (EI^+) calcd for $\text{C}_{16}\text{H}_{20}\text{F}_3\text{N}_5\text{O}_2$ (M^+) m/z 371.1569, found 371.1560.

N-{3-[[3-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)propyl](methylamino)propyl]-2,2,2-trifluoroacetamide (**60**). H₂O₂ (0.6 mL, 12.2 mmol) was added dropwise to a stirred solution of trifluoroacetic anhydride (1.7 mL, 12.2 mmol) in DCM (10 mL) at 5 °C. The mixture was stirred at 5 °C for 5 min., warmed to 20 °C for 20 min., cooled to 5 °C, and added to a stirred solution of amide **59** (453 mg, 1.2 mmol) and trifluoroacetic acid (0.19 mL, 2.4 mmol) in CHCl₃ (10 mL) at 5 °C. The mixture was stirred at 20 °C for 4 h, diluted with aqueous KHCO₃ (15 mL), and extracted with CHCl₃ (5 × 30 mL). The combined organic fraction was dried, adsorbed on to silica, and the solvent evaporated (CAUTION: use blast shield). The residue was purified by chromatography, eluting with a gradient (0–10%) of MeOH/DCM, to give 1,4-dioxide **60** (268 mg, 57%) as a yellow oil, ¹H NMR [(CD₃)₂SO] δ 9.40 (br s, 1 H, NHCO), 8.34–8.38 (m, 2 H, H-5, H-8), 8.10 (ddd, *J* = 8.7, 7.1, 1.2 Hz, 1 H, H-6), 7.94 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-7), 3.16–3.21 (m, 2 H, CH₂N), 3.04 (dd, *J* = 7.6, 7.4 Hz, 2 H, CH₂), 2.43 (br t, 6.8 Hz, 2 H, CH₂N), 2.32 (br t, *J* = 6.8 Hz, 2 H, CH₂), 2.14 (s, 3 H, NCH₃), 1.87–1.94 (m, 2 H, CH₂), 155–1.62 (m, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 155.9 (q, *J* = 37 Hz), 154.7, 139.3, 135.4, 134.4, 131.7, 120.9, 118.8, 115.8 (q, *J* = 288 Hz), 56.2, 54.3, 41.3, 37.6, 27.6, 25.8, 21.8; MS (FAB⁺) *m/z* 388 (MH⁺, 25%), 372 (5); HRMS (FAB⁺) calcd for C₁₆H₂₁F₃N₅O₃ (MH⁺) *m/z* 388.1597, found 388.1601.

*N*¹-[3-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)propyl]-*N*¹-methyl-1,3-

propanediamine (**61**). Aq. ammonia (5 mL) was added to a stirred solution of amide **60** (169 mg, 0.44 mmol) in MeOH (10 mL) and the solution stirred at 40 °C for 6 h.

The solvent was evaporated to give crude amine **61** as a brown oil, ¹H NMR

[(CD₃)₂SO] δ 8.34–8.39 (m, 2 H, H-5, H-8), 8.14 (ddd, *J* = 8.6, 7.0, 1.1 Hz, 1 H, H-6), 7.96 (ddd, *J* = 8.5, 7.0, 1.2 Hz, 1 H, H-7), 7.61 (br s, 2 H, NH₂), 3.04 (dd, *J* = 7.6, 7.4 Hz, 2 H, CH₂N), 2.85 (br dd, *J* = 7.4, 7.2 Hz, 2 H, CH₂), 2.45 (br t, *J* = 6.9 Hz, 2 H, CH₂N), 2.39 (br t, *J* = 6.7 Hz, 2 H, CH₂N), 2.17 (s, 3 H, NCH₃), 1.88–1.95 (m, 2 H, CH₂), 1.63–1.70 (m, 2 H, CH₂).

N-{3-[[3-(1,4-dioxido-1,2,4-benzotriazin-3-yl)propyl](methylamino)propyl]-4-acridinecarboxamide (**62**). The crude amine **61** was dissolved in dry THF (10 mL) and 4-(1*H*-imidazol-1-ylcarbonyl)acridine (0.18 g, 0.65 mmol) added and solution

stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–10 %) of MeOH/DCM, to give compound **62** (86 mg, 40%) as a yellow gum, which was converted to the hydrochloride salt, a pale green gum, ¹H NMR [(CD₃)₂SO] δ 11.29 (br s, 1 H, CONH), 10.90 (br s, 1 H, NH⁺Cl⁻), 9.38 (s, 1 H, H-9), 8.74 (dd, *J* = 7.0, 1.0 Hz, 1 H, H-3), 8.47 (d, *J* = 8.7 Hz, 1 H, H-1), 8.42 (dd, *J* = 8.4, 1.2 Hz, 1 H, H-5), 8.35 (dd, *J* = 8.6, 0.7 Hz, 1 H, H-8'), 8.31 (d, *J* = 8.7 Hz, 1 H, H-8), 8.21 (d, *J* = 8.4 Hz, 1 H, H-5'), 8.11 (ddd, *J* = 8.7, 7.0, 1.3 Hz, 1 H, H-6), 7.94–8.10 (m, 2 H, H-2, H-6'), 7.78 (dd, *J* = 8.7, 7.0 Hz, 1 H, H-7), 7.67–7.71 (m, 1 H, H-7'), 3.65–3.70 (m, 2 H, CH₂N), 3.30–3.37 (m, 2 H, CH₂N), 3.19–3.28 (m, 2 H, CH₂N), 3.09 (t, *J* = 7.3 Hz, 2 H, CH₂), 2.79 (d, *J* = 4.8 Hz, 3 H, NCH₃), 2.16–2.27 (m, 4 H, 2 × CH₂); ¹³C NMR [(CD₃)₂SO] δ 165.3, 153.0, 146.3, 144.7, 139.6, 139.2, 135.5, 134.7, 134.5, 134.0, 133.0, 132.3, 132.0, 128.4, 128.1, 126.6, 126.3, 125.5, 125.3, 120.9, 118.8, 53.7, 52.7, 39.4, 36.4, 26.8, 23.7, 18.8; MS (FAB⁺) *m/z* 497 (MH⁺, 12%), 481 (3); HRMS (FAB⁺) calcd for C₂₈H₂₉N₆O₃ (MH⁺) *m/z* 497.2301, found 497.2301.

Example U.

N-{3-[[3-(1,4-Dioxido-1,2,4-benzotriazin-3-yl)propyl](methyl)amino]propyl}-1-phenazinecarboxamide (**63**). Aq. ammonia (5 mL) was added to a stirred solution of amide **60** (61 mg, 0.16 mmol) in MeOH (10 mL) and the solution stirred at 40 °C for 6 h. The solvent was evaporated to give crude amine **61** as a brown oil. The crude amine **61** was dissolved in dry THF (10 mL) and 1-(1*H*-imidazol-1-ylcarbonyl)phenazine (100 mg, 0.36 mmol) added and solution stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–10%) of MeOH/DCM, to give compound **63** (44 mg, 56%) as a yellow gum, which was converted to the hydrochloride salt and recrystallised, mp (MeOH/EtOAc) 173 °C (dec.); ¹H NMR [(CD₃)₂SO] δ 10.31 (t, *J* = 5.8 Hz, 1 H, NH), 9.90 (br s, 1 H, NH⁺Cl⁻), 8.61 (dd, *J* = 7.1, 1.4 Hz, 1 H, H-2), 8.48 (dd, *J* = 9.1, 1.4 Hz, 1 H, H-9), 8.42 (dd, *J* = 8.6, 1.4 Hz, 1 H, H-4), 8.34 (d, *J* = 8.4 Hz, 1 H, H-6), 8.30 (dd, *J* = 8.6, 1.1 Hz, 1 H, H-8'), 8.26 (dd, *J* = 8.3, 1.4 Hz, 1 H, H-5'), 7.93–8.13 (m, 5 H, H-3, H-7, H-8, H-6', H-7'), 3.62–3.67 (m, 2 H, CH₂N), 3.30–3.34 (m, 2 H, CH₂N), 3.22–3.38 (m, 2 H, CH₂N), 3.07–3.11 (m, 2 H, CH₂), 2.82 (br s, 3 H, NCH₃), 2.10–2.22 (m, 4 H, 2 × CH₂); ¹³C NMR [(CD₃)₂SO] δ 164.8, 153.1, 142.7, 142.5,

141.2, 140.0, 139.2, 135.6, 134.5, 133.5, 132.7, 132.0, 131.9, 131.6, 130.9, 130.3, 129.4, 129.1, 121.0, 118.8, 54.0, 52.9, 39.4, 36.4, 26.7, 23.8, 19.0; MS (FAB⁺) *m/z* 498 (MH⁺, 20%), 482 (5); HRMS (FAB⁺) calcd for C₂₇H₂₈N₇O₃ (MH⁺) *m/z* 498.2254, found 498.2256.

5

Example V

3-[(7-Chloro-4-quinolinyl)amino]-N-{3-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}propanamide (65).

A solution of *N*-(7-chloro-4-quinolinyl)-β-alanine (64) (Titus et al, *J. Org. Chem.*,
 10 1948, 13, 39-62) (303 mg, 1.2 mmol) and CDI (235 mg, 1.5 mmol) in DMF (5 mL) was stirred at 50 °C for 1 h. The solvent was evaporated and the residue crystallised from DCM/pet. ether to give the imidazolidine (290 mg, 80%), which was used directly. A solution of *N*¹-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,3-propanediamine (16) (92 mg, 390 μmol) and imidazolidine (176 mg, 590 μmol) in DMF (10 mL) was stirred at
 15 20 °C for 3 days, the solvent evaporated and the residue recrystallised from hot MeOH to give compound 65 (84 mg, 46%) as a red powder, mp (MeOH) 202 °C (dec.); ¹H NMR [(CD₃)₂SO] δ 8.40 (d, *J* = 5.4 Hz, 1 H, H-2'), 8.26 (br t, *J* = 6.2 Hz, 1 H, NH), 8.18–8.12 (m, 2 H, H-5, H-8), 8.13 (d, *J* = 8.6 Hz, 1 H, H-5'), 7.99 (br t, *J* = 5.7 Hz, 1 H, NH), 7.93 (ddd, *J* = 8.6, 7.1, 1.2 Hz, 1 H, H-6), 7.75 (d, *J* = 2.2 Hz, 1 H, H-8'), 7.56 (ddd, *J* = 8.6, 7.1, 1.3 Hz, 1 H, H-7), 7.40 (dd, *J* = 8.6, 2.2 Hz, 1 H, H-6'), 7.37 (br t, *J* = 5.4 Hz, 1 H, NH), 6.52 (d, *J* = 5.4 Hz, 1 H, H-3'), 3.49–3.54 (m, 2 H, CH₂N), 3.36–3.41 (m, 2 H, CH₂N), 3.12–3.17 (m, 2 H, CH₂N), 2.47–2.51 (m, 2 H, CH₂), 1.70–1.77 (m, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 170.3, 151.8, 149.7, 149.6, 149.0, 138.1, 135.4, 133.2, 129.8, 127.4, 126.8, 124.0, 123.9, 121.0, 117.4, 116.8,
 20 99.7, 39.0, 38.2, 35.8, 34.3, 28.5; MS (FAB⁺) *m/z* 470 (MH⁺, 5%), 468 (15), 454 (1), 452 (3); HRMS (FAB⁺) calcd for C₂₂H₂₃³⁵ClN₇O₃ (MH⁺) *m/z* 468.1551, found 468.1546; calcd for C₂₂H₂₃³⁷ClN₇O₃ (MH⁺) *m/z* 470.1540, found 470.1535.

Example W.

30 *N*-[3-(Methyl{3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-4-acridinecarboxamide (74).

7-Methyl-1,2,4-benzotriazin-3-ol 1-Oxide (67). NaNO₂ (9.0g, 130.6 mmol) was added in small portions to a stirred solution of 7-methyl-1,2,4-benzotriazin-3-amine 1-

oxide [Hay et al, *J. Med. Chem.* **2003**, 46, 169–182] (**66**) (11.5 g, 65.3 mmol) in TFA (40 mL) at –5 to 0 °C. After the addition was completed stirring was continued for a further 1 h, the mixture was poured into ice (300 g) and stirred 1 h. The resulting pale yellow precipitate was filtered and washed with water. The precipitate was dissolved
5 in 8% aqueous NH₃, filtered and the filtrate was acidified with cHCl. The resulting precipitate was filtered, washed with water and dried to give alcohol **67** (11.5 g, 100%) which was used without further purification.

3-Chloro-7-methyl-1,2,4-benzotriazine 1-Oxide (68). Alcohol **67** (3.15 g, 65.3
10 mmol) was refluxed in POCl₃ (50 mL) for 5 h. The reaction mixture was cooled and carefully poured into ice/water and stirred for 30 min. The resulting precipitate was filtered, air dried and purified by chromatography, eluting with a gradient (50–100%) of DCM/hexane, to give chloride **68** (9.0 g, 66%), mp (DCM/hexane) 174–176 °C; ¹H NMR δ 8.20 (br s, 1 H, H-8), 7.89 (d, *J* = 8.6 Hz, 1 H, H-5), 7.82 (dd, *J* = 8.6, 1.9 Hz,
15 1 H, H-6), 2.61 (s, 3 H, CH₃). Anal. calcd for C₈H₆ClN₃O: C, 49.1; H, 3.1; N, 21.5; Cl, 18.1, found: C, 49.1, H, 3.1; N, 21.5; Cl, 18.5%.

tert-Butyl 3-(Methyl-{3-[(7-methyl-1-oxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propylcarbamate (70). A mixture of chloride **68** (2.18 g,

20 11.1 mmol), *tert*-butyl 3-[(3-aminopropyl)(methylamino)propylcarbamate **69** (4.35 g, 17.8 mmol) and Et₃N (2.3 mL, 16.5 mmol) in DME (25 mL) was heated at 85 °C for 3 h. The solvent was evaporated, the residue was dissolved in DCM (100 mL) and washed with aqueous NH₃. The organic layer was separated and the aqueous layer further extracted with DCM (3 × 30 mL). The combined organic fraction was dried
25 and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM, to give **70** (3.7 g, 93%) as a yellow solid, mp (DCM/hexane) 117–120 °C; ¹H NMR [(CD₃)₂SO] δ 7.94 (s, 1 H, H-8), 7.79 (br s, 1 H, NH), 7.62 (dd, *J* = 8.7, 2.0 Hz, 1 H, H-6), 7.49 (d, *J* = 8.6 Hz, 1 H, H-5), 6.75 (t, *J* = 5.3 Hz, 1 H, NH), 3.34 (br q, *J* = 6.4 Hz, 2 H, CH₂), 2.93 (br q, *J* =
30 6.5 Hz, 2 H, CH₂), 2.41 (s, 3 H, CH₃), 2.35 (t, *J* = 6.9 Hz, 2 H, CH₂), 2.27 (t, *J* = 7.0 Hz, 2 H, CH₂), 2.12 (s, 3 H, CH₃), 1.70 (br quin, *J* = 6.9 Hz, 2 H, CH₂), 1.51 (br quin, *J* = 7.1 Hz, 2 H, CH₂), 1.35 (s, 9 H, 3 × CH₃); ¹³C NMR [(CD₃)₂SO] 158.6, 155.4,

146.8, 137.5, 134.4, 129.5, 125.7, 118.4, 77.2, 54.8, 54.7, 41.6, 39.0, 38.2, 28.1 (3), 27.1, 26.1, 20.6.

***N*¹-(3-Aminopropyl)-*N*¹-methyl-*N*³-(6-methyl-1-oxido-1,2,4-benzotriazin-3-yl)-**

- 5 **1,3-propanediamine (71).** Carbamate **70** (4.1 g, 10.1 mmol) was dissolved in methanolic HCl (50 mL) and stirred for 48 h at 20 °C. Excess reagent and solvent were evaporated and the residue was partitioned between DCM and aqueous NH₃, the organic layer was separated and the aqueous layer was further extracted with DCM (4 × 30 mL). The combined organic fraction was dried and the solvent evaporated.
- 10 The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(3–10%) MeOH/DCM, to give amine **71** (1.14 g, 100%) as a yellow solid, ¹H NMR [(CD₃)₂SO] δ 7.94 (s, 1 H, H-8), 7.81 (br s, 1 H, NH), 7.63 (dd, *J* = 8.7, 1.9 Hz, 1 H, H-6), 7.49 (d, *J* = 8.6 Hz, 1 H, H-5), 3.34 (br q, *J* = 6.3 Hz, 2 H, CH₂), 2.54–2.58 (m, 2 H, CH₂), 2.41 (s, 3 H, CH₃), 2.36 (t, *J* = 6.9 Hz, 2 H, CH₂),
- 15 2.31 (t, *J* = 7.2 Hz, 2 H, CH₂), 2.13 (s, 3 H, CH₃), 1.71 (br quin, *J* = 7.0 Hz, 2 H, CH₂), 1.48 (br quin, *J* = 7.0 Hz, 2 H, CH₂), NH₂ not observed; HRMS (FAB⁺) calcd for C₁₅H₂₅N₆O (MH⁺) *m/z* 305.2090, found 305.2090.

2,2,2-Trifluoro-*N*-[3-(methyl-{3-[(7-methyl-1-oxido-1,2,4-benzotriazin-3-

- 20 **yl)amino]propyl}amino)propyl]acetamide (72).** CF₃CO₂Et (2.43 mL, 20.4 mmol) and H₂O (0.36 mL, 20.4 mmol) were added to a solution of amine **71** (3.1 g, 10.2 mmol) in CH₃CN (50 mL) and the reaction mixture heated at reflux for 20 h. The solvent was evaporated and residue partitioned between DCM and aqueous NaHCO₃. The organic layer was separated and the aqueous layer was further extracted with
- 25 DCM (3 × 50 mL). The combined organic fraction was dried and the solvent evaporated to give acetamide **72** (3.75 g, 92%) as a yellow solid, mp (DCM/hexane) 121–124 °C; ¹H NMR [(CD₃)₂SO] δ 9.44 (br s, 1 H, NH), 7.94 (s, 1 H, H-8), 7.80 (br s, 1 H, NH), 7.62 (dd, *J* = 8.7, 1.9 Hz, 1 H, H-6), 7.48 (d, *J* = 8.6 Hz, 1 H, H-5), 3.22 (br q, *J* = 6.5 Hz, 2 H, CH₂), 2.41 (s, 3 H, CH₃), 2.37 (t, *J* = 7.0 Hz, 2 H, CH₂), 2.32 (t,
- 30 *J* = 6.9 Hz, 2 H, CH₂), 2.15 (s, 3 H, CH₃), 1.71 (br quin, *J* = 7.0 Hz, 2 H, CH₂), 1.63 (br quin, *J* = 6.5 Hz, 2 H, CH₂), CH₂ not observed; ¹³C NMR [(CD₃)₂SO] δ 158.6, 156.3 (q, *J* = 36 Hz), 146.8, 137.6, 134.5, 129.6, 125.7, 118.4, 116.1 (q, *J* = 288 Hz), 54.7, 54.5, 41.5, 39.0, 37.8, 26.1, 25.8, 20.6; HRMS (FAB⁺) calcd for C₁₇H₂₄F₃N₆O₂

(MH⁺) *m/z* 401.1913, found 401.1896. Anal. calcd for C₁₇H₂₃F₃N₆O₂: C, 51.0; H, 5.8; F, 14.2; N, 21.0; found: C, 51.3; H, 5.9; F, 14.0; N, 21.0%.

2,2,2-Trifluoro-*N*-[3-(methyl{3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-

yl)amino]propyl}amino)propyl]acetamide (73). A solution of trifluoroacetic acid [made from trifluoroacetic anhydride (12.4 mL, 89.4 mmol) and 70% H₂O₂ (4.3 mL, 89.4 mmol) in DCM (50 mL)] was added to a solution of acetamide **72** (3.6 g, 8.9 mmol) and trifluoroacetic acid (2.8 mL, 35.8 mol) in DCM (50 mL) at 0 °C and the reaction mixture was stirred at 20 °C for 18 h. The reaction mixture was slowly added to a solution of aqueous NaHCO₃ (100 mL) at 5 °C. The organic layer was separated and the aqueous layer was further extracted with DCM (4 × 30 mL). The combined organic fraction was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–8%) of MeOH/DCM, to give (i) starting material (1.44 g, 40%) and (ii) acetamide **73** (1.30 g, 35%) as a red solid, mp (DCM/hexane) 117–119 °C; ¹H NMR [(CD₃)₂SO] δ 9.44 (br s, 1 H, NH), 8.36 (t, *J* = 5.9 Hz, 1 H, NH), 8.03 (d, *J* = 8.9 Hz, 1 H, H-5), 8.01 (s, 1 H, H-8), 7.78 (dd, *J* = 8.9, 1.6 Hz, 1 H, H-6), 3.42 (br q, *J* = 6.6 Hz, 2 H, CH₂), 3.23 (br q, *J* = 6.5 Hz, 2 H, CH₂), 2.47 (s, 3 H, CH₃), 2.38 (t, *J* = 6.7 Hz, 2 H, CH₂), 2.32 (t, *J* = 6.9 Hz, 2 H, CH₂), 2.15 (s, 3 H, CH₃), 1.75 (br quin, *J* = 6.9 Hz, 2 H, CH₂), 1.65 (br quin, *J* = 7.1 Hz, 2 H, CH₂); ¹³C NMR [(CD₃)₂SO] δ 156.0 (q, *J* = 36 Hz), 149.4, 137.4, 137.2, 136.7, 129.6, 119.4, 116.6, 115.9 (q, *J* = 288 Hz), 54.9, 54.6, 41.5, 39.5, 37.6, 26.0, 25.8, 20.7; HRMS (FAB⁺) calcd for C₁₇H₂₄F₃N₆O₃ (MH⁺) *m/z* 417.1862, found 417.1859. Anal. calcd for C₁₇H₂₃F₃N₆O₃: C, 49.0; H, 5.6; F, 13.7; N, 20.2; found: C, 49.3; H, 5.5; F, 13.6; N, 20.2%.

***N*-[3-(Methyl{3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-**

yl)amino]propyl}amino)propyl]-4-acridinecarboxamide (74). Aqueous NH₃ (5 mL) was added to a solution of acetamide **73** (135 mg, 0.32 mmol) in MeOH (10 mL) and the reaction mixture stirred at 20 °C for 18 h. The solvent was evaporated, the residue was dissolved in DMF (5 mL), 4-(1*H*-imidazol-1-ylcarbonyl)acridine (177 mg, 0.64 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM, to give compound **74** (168 mg, 100%)

as a red solid, mp (DCM/hexane) 166–168 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 11.32 (br s, 1 H, NH), 9.26 (s, 1 H, ArH), 8.68 (d, $J = 6.7$ Hz, 1 H, ArH), 8.37 (t, $J = 5.6$ Hz, 1 H, NH), 8.33 (d, $J = 8.0$ Hz, 1 H, ArH), 8.23 (d, $J = 8.7$ Hz, 1 H, ArH), 8.17 (d, $J = 8.4$ Hz, 1 H, ArH), 7.89–7.96 (m, 3 H, ArH), 7.64–7.74 (m, 3 H, ArH), 3.59 (br q, $J = 6.0$ Hz, 2 H, CH_2), 3.41 (br q, $J = 6.2$ Hz, 2 H, CH_2), 2.56 (t, $J = 7.0$ Hz, 2 H, CH_2), 2.46 (t, $J = 6.9$ Hz, 2 H, CH_2), 2.44 (s, 3 H, CH_3), 2.23 (s, 3 H, CH_3), 1.91 (br quin, $J = 6.7$ Hz, 2 H, CH_2), 1.79 (br quin, $J = 6.6$ Hz, 2 H, CH_2); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 164.7, 149.2, 147.0, 145.4, 138.5, 137.2, 137.1, 136.5, 134.3, 132.6, 131.8, 129.4, 128.5, 128.4, 128.3, 126.4, 126.4, 125.5, 125.2, 119.3, 116.5, 55.3, 55.1, 41.8, 39.5, 37.4, 26.9, 25.9, 20.7; HRMS (FAB $^+$) calcd for $\text{C}_{29}\text{H}_{32}\text{N}_7\text{O}_3$ (MH^+) m/z 526.2567, found 526.2537. Anal. calcd for $\text{C}_{29}\text{H}_{31}\text{N}_7\text{O}_3 \cdot \frac{1}{4}\text{H}_2\text{O}$: C, 65.7; H, 6.0; N, 18.5; found: C, 65.8; H, 5.9; N, 18.7%.

Example X

***N*-[3-(Methyl{3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-2-(4-pyridinyl)-8-quinolinecarboxamide (75).**

Aqueous NH_3 (5 mL) was added to a solution of acetamide **73** (135 mg, 0.32 mmol) in MeOH (5 mL) and the mixture stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL) and 8-(1*H*-imidazol-1-ylcarbonyl)-2-(4-pyridinyl)quinoline (160 mg, 0.64 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated, the residue dissolved in DCM (20 mL) and washed with water (3×15 mL). The organic layer was separated, dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH_3 /(0–3%) MeOH/DCM, to give compound **75** (158 mg, 89%) as a red solid, mp (DCM/hexane) 178–180 °C; ^1H NMR δ 10.89 (t, $J = 4.9$ Hz, 1 H, NH), 8.80–8.84 (m, 3 H, ArH), 8.36 (d, $J = 8.6$ Hz, 1 H, ArH), 8.28 (t, $J = 4.8$ Hz, 1 H, NH), 8.05 (s, 1 H, ArH), 8.02 (d, $J = 8.9$ Hz, 1 H, ArH), 7.90–7.95 (m, 4 H, ArH), 7.67 (t, $J = 7.7$ Hz, 1 H, ArH), 7.56 (dd, $J = 8.9, 1.7$ Hz, 1 H, ArH), 4.73 (br q, $J = 6.5$ Hz, 2 H, CH_2), 3.52 (br q, $J = 6.0$ Hz, 2 H, CH_2), 2.54 (t, $J = 7.3$ Hz, 2 H, CH_2), 2.49 (s, 3 H, CH_3), 2.47 (t, $J = 6.3$ Hz, 2 H, CH_2), 2.24 (s, 3 H, CH_3), 2.01 (br quin, $J = 7.1$ Hz, 2 H, CH_2), 1.74 (br quin, $J = 6.2$ Hz, 2 H, CH_2); ^{13}C NMR δ 165.8, 154.4, 150.8 (2), 149.4, 146.2, 145.4, 138.9, 137.7, 137.6, 136.8, 134.3, 131.2, 130.0, 129.9, 127.9, 127.2, 121.7 (2), 120.1, 118.6, 117.0, 56.6, 55.7, 41.9, 41.3, 38.2, 27.7,

25.6, 21.4; HRMS (FAB⁺) calcd for C₃₀H₃₃N₈O₃ (MH⁺) *m/z* 553.2676, found 553.2669. Anal. calcd for C₃₀H₃₂N₈O₃·½H₂O: C, 64.2; H, 5.9; N, 20.0; found: C, 64.0; H, 5.7; N, 20.0 %.

5 Example Y

N-[3-(Methyl{3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-1-phenazinecarboxamide (**76**). Aqueous NH₃ (6 mL) was added to a solution of acetamide **73** (141 mg, 0.34 mmol) in MeOH (10 mL) and the mixture stirred at 20 °C for 18 h. The solvent was evaporated, the residue was dissolved in DMF (5 mL) and 1-(1*H*-imidazol-1-ylcarbonyl)phenazine (183 mg, 0.68 mmol) was added and mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–4%) MeOH/DCM, to give compound **76** (178 mg, 100%) as a red solid, mp (DCM/hexane) 118–122 °C; ¹H NMR δ 10.86 (br s, 1 H, NH), 8.93 (dd, *J* = 7.0, 1.5, 1 H, ArH), 8.45 (br s, 1 H, NH), 8.34 (dd, *J* = 8.7, 1.5 Hz, 1 H, ArH), 8.21–8.27 (m, 2 H, ArH), 8.30 (s, 1 H, ArH), 7.99 (d, *J* = 8.8 Hz, 1 H, ArH), 7.93 (dd, *J* = 8.7, 7.2 Hz, 1 H, ArH), 7.79–7.83 (m, 2 H, ArH), 7.53 (dd, *J* = 8.9, 1.8 Hz, 1 H, ArH), 3.77 (br q, *J* = 6.4 Hz, 2 H, CH₂), 3.65 (br q, *J* = 5.9 Hz, 2 H, CH₂), 2.67 (t, *J* = 7.4 Hz, 2 H, CH₂), 2.62 (t, *J* = 6.1 Hz, 2 H, CH₂), 2.49 (s, 3 H, CH₃), 2.35 (s, 3 H, CH₃), 2.09 (br quin, *J* = 7.1 Hz, 2 H, CH₂), 1.88 (br quin, *J* = 6.2 Hz, 2 H, CH₂); ¹³C NMR δ 165.1, 149.4, 143.4, 142.9, 141.3, 140.8, 137.8, 137.7, 136.7, 135.1, 135.0, 133.4, 131.6, 131.0, 130.0, 129.8, 129.7, 129.0, 120.1, 116.8, 56.8, 55.8, 42.0, 41.5, 38.2, 27.5, 25.6, 21.4; HRMS (FAB⁺) calcd for C₂₈H₃₁N₈O₃ (MH⁺) *m/z* 527.2519, found 527.2512. Anal. calcd for C₂₈H₃₀N₈O₃·¼H₂O: C, 63.3; H, 5.8; N, 21.1; found: C, 63.2, H, 5.9, N, 21.4%.

Example Z

9-Methyl-*N*-[3-(3-[(7-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-1-phenazinecarboxamide (**77**). Aqueous NH₃ (5 mL) was added to a solution of acetamide **73** (135 mg, 0.32 mmol) in MeOH (5 mL) and the solution stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL) and 1-(1*H*-imidazol-1-ylcarbonyl)-9-methylphenazine (172 mg, 0.6 mmol) was added and stirred at 20 °C for 48 h. The solvent was evaporated

and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH_3 /(0–4%) MeOH/DCM, to give compound **77** (147 mg, 91%) as a red solid, mp (DCM/hexane) 119–122 °C; ^1H NMR δ 11.02 (br s, 1 H, NH), 8.97 (dd, J = 7.2, 1.5 Hz, 1 H, ArH), 8.37 (dd, J = 8.5, 1.4 Hz, 1 H, ArH), 8.22 (br s, 1 H, NH),
 5 8.10 (d, J = 8.7 Hz, 1 H, ArH), 8.05 (s, 1 H, ArH), 8.02 (d, J = 8.9 Hz, 1 H, ArH), 7.96 (dd, J = 8.7, 7.2 Hz, 1 H, ArH), 7.78 (dd, J = 8.6, 6.9 Hz, 1 H, ArH), 7.69–7.73 (m, 1 H, ArH), 7.55 (dd, J = 8.9, 1.6 Hz, 1 H, ArH), 3.77 (br q, J = 6.6 Hz, 2 H, CH_2), 3.65 (br q, J = 6.0 Hz, 2 H, CH_2), 2.93 (s, 3 H, CH_3), 2.61 (t, J = 7.3 Hz, 2 H, CH_2), 2.58 (t, J = 6.1 Hz, 2 H, CH_2), 2.49 (s, 3 H, CH_3), 2.31 (s, 3 H, CH_3), 2.07 (br quin, J = 7.2 Hz, 2 H, CH_2), 1.86 (br quin, J = 6.2 Hz, 2 H, CH_2); ^{13}C NMR δ 165.1, 149.4, 143.2, 143.1, 141.0, 139.7, 137.8, 137.7, 136.8, 136.6, 135.1, 133.3, 131.0, 130.9, 130.0, 129.9, 129.4, 127.7, 120.1, 116.9, 56.4, 55.8, 42.0, 41.2, 38.3, 27.9, 25.7, 21.4, 18.1; HRMS (FAB $^+$) calcd for $\text{C}_{29}\text{H}_{33}\text{N}_8\text{O}_3$ (MH^+) m/z 541.2676, found 541.2669.
 10 Anal. calcd for $\text{C}_{29}\text{H}_{32}\text{N}_8\text{O}_3 \cdot \frac{1}{4}\text{H}_2\text{O}$: C, 63.9; H, 6.0; N, 20.6; found: C, 63.8; H, 5.9; N, 20.8%.

Example AA

***N*-[3-(Methyl{3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-4-acridinecarboxamide (85).**

20 **6-Methyl-1,2,4-benzotriazin-3-ol 1-Oxide (79).** NaNO_2 (2.5 g, 36.3 mmol) was added in small portions to a stirred solution of 6-methyl-1,2,4-benzotriazin-3-amine 1-oxide (**78**) [Hay et. al., *J. Med Chem.* **2003**, 46, 169-182] (3.2 g, 18.2 mmol) in TFA (15 mL) at –5 to 0 °C. After the addition was completed the reaction mixture was stirred for further 1 h and poured into ice (150 g). The resulting pale yellow
 25 precipitate was filtered, washed with water and dried to give compound **79** (3.2 g, 97%), which was used without further purification.

3-Chloro-6-methyl-1,2,4-benzotriazine 1-Oxide (80). Compound **79** (3.2 g, 17.8 mmol) was heated at reflux in POCl_3 (25 mL) for 3 h. Excess reagent was evaporated
 30 and the residue was stirred in ice/water (150 mL) for 20 min. The resulting precipitate was filtered, air dried and purified by chromatography, eluting with a gradient (50–100%) of DCM/pet. ether, to give chloride **80** (2.5 g, 79%) as a white crystalline solid, mp (DCM/hexane) 156–158 °C; ^1H NMR δ 8.30 (d, J = 8.8 Hz, 1 H, H-8), 7.75

(br s, 1 H, H-5), 7.64 (dd, $J = 9.4, 1.6$ Hz, 1 H, H-7), 2.62 (s, 3 H, CH₃). Anal. calcd for C₈H₆ClN₃O: C, 49.1; H, 3.1; N, 21.5; found: C, 49.2; H, 3.1; N, 21.5%.

***tert*-Butyl 3-(Methyl{3-[(6-methyl-1-oxido-1,2,4-benzotriazin-3-**

- 5 **yl)amino]propyl}amino)propylcarbamate (81).** A mixture of chloride **80** (2.23 g, 11.4 mmol), *tert*-butyl-3[(aminopropyl)(methyl)amino]propylcarbamate **69** (Huang et al., *J. Med. Chem.* **1992**, 35, 2414-18) (3.34 g, 14.4 mmol) and triethylamine (2.3 mL, 16.5 mmol) in DME (60 mL) was heated at 85 °C for 3 h. The solvent was evaporated, the residue was dissolved in DCM (100 mL) and washed with aqueous
10 NH₃ (40 mL). The organic layer was separated, the aqueous layer further extracted with DCM (3 × 30 mL), the combined organic fraction dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM to give carbamate **81** (3.7 g, 80%) as a yellow solid, mp (DCM/hexane) 117–120 °C; ¹H NMR [(CD₃)₂SO] δ 8.01 (d, $J = 8.7$
15 Hz, 1 H, H-8), 7.84 (br s, 1 H, NH), 7.37 (br s, 1 H, H-5), 7.15 (dd, $J = 8.8, 1.7$ Hz, 1 H, H-7), 6.75 (t, $J = 5.2$ Hz, 1 H, NH), 3.33–3.37 (m, 2 H, CH₂), 2.94 (br q, $J = 6.5$ Hz, 2 H, CH₂), 2.42 (s, 3 H, CH₃), 2.35 (t, $J = 6.9$ Hz, 2 H, CH₂), 2.28 (t, $J = 7.0$ Hz, 2 H, CH₂), 2.08 (s, 3 H, CH₃), 1.70 (br quin, $J = 6.9$ Hz, 2 H, CH₂), 1.51 (br quin, $J = 6.9$ Hz, 2 H, CH₂), 1.35 (s, 9 H, 3 × CH₃); ¹³C NMR [(CD₃)₂SO] 159.0, 155.5, 148.5,
20 146.6, 128.2, 126.4, 124.8, 119.5, 77.2, 54.8, 54.7, 41.6, 39.0, 38.2, 28.1 (3), 27.1, 26.1, 21.3; HRMS (FAB⁺) calcd for C₂₀H₃₃N₆O₃ (MH⁺) m/z 405.2614, found 405.2616.

***N*¹-(3-Aminopropyl)-*N*¹-methyl-*N*³-(6-methyl-1-oxido-1,2,4-benzotriazin-3-yl)-**

- 25 **1,3-propanediamine (82).** Carbamate **81** (2.1 g, 5.19 mmol) was dissolved in methanolic HCl (50 mL) and stirred 48 h at 20 °C. Excess reagent and solvent were evaporated and the residue partitioned between DCM and aqueous NH₃. The organic layer was separated and the aqueous layer was further extracted with DCM (4 × 30 mL). The combined organic fraction was dried and the solvent evaporated.
30 The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(3–7%) MeOH/DCM, to give amine **82** (1.57 g, 99%) as a yellow solid, mp 118–122 °C (DCM/MeOH); ¹H NMR [(CD₃)₂SO] δ 8.01 (d, $J = 8.8$ Hz, 1 H, H-8), 7.87 (br s, 1 H, NH), 7.37 (br s, 1 H, H-5), 7.16 (dd, $J = 8.8, 1.7$ Hz, 1 H, H-7),

3.14–3.34 (m, 4 H, CH₂, NH₂), 2.54 (t, J = 6.5 Hz, 2 H, CH₂), 2.42 (s, 3 H, CH₃), 2.35 (t, J = 6.9 Hz, 2 H, CH₂), 2.31 (t, J = 7.2 Hz, 2 H, CH₂), 2.13 (s, 3 H, CH₃), 1.71 (br quin, J = 7.0 Hz, 2 H, CH₂), 1.47 (br quin, J = 7.0 Hz, 2 H, CH₂); HRMS (FAB⁺) calcd for C₁₅H₂₅N₆O (MH⁺) m/z 305.2090, found 305.2088.

5

2,2,2-Trifluoro-*N*-[3-(methyl{3-[(6-methyl-1-oxido-1,2,4-benzotriazin-3-

yl)amino]propyl}amino)propyl]acetamide (83). CF₃CO₂Et (0.88 mL, 7.4 mmol)

and H₂O (0.13 mL, 7.4 mmol) were added to a stirred solution of amine **82** (1.5 g, 4.9 mmol) in CH₃CN (50 mL) and the reaction mixture heated at reflux for 20 h. The

10 solvent was evaporated and the residue partitioned between DCM and aqueous NaHCO₃. The organic layer was separated, the aqueous layer further extracted with DCM (3 × 30 mL), the combined organic fraction dried, and the solvent evaporated to give acetamide **83** (1.9 g, 100%) as a yellow solid, mp (DCM/hexane) 127–130 °C; ¹H NMR [(CD₃)₂SO] δ 9.43 (br s, 1 H, NH), 8.01 (d, J = 8.8 Hz, 1 H, H-8), 7.85 (br s, 15 1 H, NH), 7.36 (br s, 1 H, H-5), 7.15 (dd, J = 8.8, 1.5 Hz, 1 H, H-7), 3.23 (br q, J = 6.5 Hz, 2 H, CH₂), 2.43 (s, 3 H, CH₃), 2.38 (t, J = 6.9 Hz, 2 H, CH₂), 2.32, (t, J = 6.9 Hz, 2 H, CH₂), 2.15 (s, 3 H, CH₃), 1.72 (br quin, J = 7.0 Hz, 2 H, CH₂), 1.64 (br quin, J = 7.0 Hz, 2 H, CH₂), CH₂ not observed; ¹³C NMR [(CD₃)₂SO] δ 159.0, 156.0 (q, J = 36 Hz), 148.5, 146.6, 128.2, 126.4, 124.8, 119.5, 115.9 (q, J = 288 Hz), 54.7, 54.5, 20 41.5, 38.9, 37.7, 26.1, 25.8, 21.3; HRMS (FAB⁺) calcd for C₁₇H₂₄F₃N₆O₂ (MH⁺) 401.1913, found 401.1896. Anal. calcd for C₁₇H₂₃F₃N₆O₂: C, 51.0; H, 5.8; F, 14.2; N, 21.0; found: C, 51.1; 6.0; F, 14.2; N, 21.0%.

2,2,2-Trifluoro-*N*-[3-(methyl{3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-

25 **yl)amino]propyl}amino)propyl]acetamide (84).** A solution of trifluoroacetic acid [prepared from trifluoroacetic anhydride (5.7 mL, 6.3 mmol) and 70% H₂O₂ (2.0 mL, 6.3 mmol) in DCM (10 mL)] was added to a suspension of acetamide **83** (1.63 g, 4.1 mmol) and trifluoroacetic acid (0.63 mL, 8.1 mmol) in DCM (20 mL) and the mixture stirred at 20 °C for 18 h. The mixture was slowly added to a cooled solution of 30 aqueous NaHCO₃ (100 mL). The organic layer was separated and the aqueous layer extracted further with DCM (4 × 30 mL). The combined organic fraction was dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–5%) of MeOH/DCM, to give (i) starting material **83** (807 mg, 49%) and

(ii) dioxide **84** (509 mg, 30%) as a red solid, mp (DCM/hexane) 128–131 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 9.43 (br s, 1 H, NH), 8.42 (t, J = 6.0 Hz, 1 H, NH), 8.32 (d, J = 8.9 Hz, 1 H, H-8), 7.93 (s, 1 H, H-5), 7.38 (dd, J = 9.0, 1.7 Hz, 1 H, H-7), 3.43 (br q, J = 6.6 Hz, 2 H, CH_2), 3.22 (br q, J = 6.6 Hz, 2 H, CH_2), 2.53 (s, 3 H, CH_3), 2.38 (t, J = 6.7 Hz, 2 H, CH_2), 2.31 (t, J = 6.9 Hz, 2 H, CH_2), 2.15 (s, 3 H, CH_3), 1.75 (br quin, J = 6.9 Hz, 2 H, CH_2), 1.65 (br quin, J = 7.0 Hz, 2 H, CH_2); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 155.9 (q, J = 36 Hz), 149.8, 146.9, 138.0, 128.8, 128.2, 120.8, 116.1 (q, J = 288 Hz), 115.5, 54.9, 54.6, 41.5, 39.5, 37.7, 26.0, 25.8, 21.6; HRMS (FAB $^+$) calcd for $\text{C}_{17}\text{H}_{24}\text{F}_3\text{N}_6\text{O}_3$ (MH^+) m/z 417.1862, found 417.1868. Anal. calcd for $\text{C}_{17}\text{H}_{23}\text{F}_3\text{N}_6\text{O}_3$: C, 49.0; H, 5.6; F, 13.7; N, 20.2; found: C, 49.3; H, 5.9; F, 14.0; N, 20.4%.

***N*-[3-(Methyl{3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-4-acridinecarboxamide (85).** Aqueous NH_3 (5 mL) was added to a solution of dioxide **84** (125 mg, 0.3 mmol) in MeOH (5 mL) and the reaction mixture was stirred at 20 °C for 18 h. The solvent was evaporated, the residue was dissolved in DMF (5 mL), 4-(1*H*-imidazol-1-ylcarbonyl)acridine (164 mg, 0.6 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH_3 /(0–5%) MeOH/DCM, to give compound **85** (148 mg, 94%) as a red solid, mp (DCM/hexane) 158–160 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 11.30 (t, J = 5.4 Hz, 1 H, NH), 9.23 (s, 1 H, ArH), 8.67 (dd, J = 7.1, 1.4 Hz, 1 H, ArH), 8.48 (t, J = 5.7 Hz, 1 H, NH), 8.32 (dd, J = 8.4, 1.2 Hz, 1 H, ArH), 8.23 (d, J = 8.7 Hz, 1 H, ArH), 8.17 (d, J = 8.4 Hz, 1 H, ArH), 8.00 (d, J = 8.9 Hz, 1 H, ArH), 7.92 (ddd, J = 7.7, 7.3, 1.4 Hz, 1 H, ArH), 7.76 (s, 1 H, ArH), 7.71 (t, J = 7.7 Hz, 1 H, ArH), 7.65 (t, J = 7.4 Hz, 1 H, ArH), 7.31 (dd, J = 9.0, 1.5 Hz, 1 H, ArH), 3.59 (br q, J = 6.3 Hz, 2 H, CH_2), 3.42 (br q, J = 6.4 Hz, 2 H, CH_2), 2.57 (t, J = 7.0 Hz, 2 H, CH_2), 2.47 (t, J = 6.6 Hz, 2 H, CH_2), 2.44 (s, 3 H, CH_3), 2.23 (s, 3 H, CH_3), 1.92 (br quin, J = 6.9 Hz, 2 H, CH_2), 1.79 (br quin, J = 6.6 Hz, 2 H, CH_2); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 164.7, 149.6, 146.9, 146.8, 145.4, 138.4, 137.7, 134.3, 132.6, 131.8, 128.7, 128.6, 128.4, 128.3, 128.0, 126.4, 126.3, 125.5, 125.1, 120.7, 115.2, 55.4, 55.1, 41.7, 39.8, 37.4, 27.0, 25.8, 21.5; HRMS (FAB $^+$) calcd for $\text{C}_{29}\text{H}_{32}\text{N}_7\text{O}_3$ (MH^+) m/z 526.2567, found 526.2535. Anal. calcd for $\text{C}_{29}\text{H}_{31}\text{N}_7\text{O}_3$: C, 66.3; H, 5.9; N, 18.7; found: C, 66.0; H, 6.0; N, 18.8%.

Example AB

***N*-[3-(Methyl{3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-2-(4-pyridinyl)-8-quinolinecarboxamide (86).**

Aqueous NH₃ (5 mL) was added to a solution of dioxide **84** (126 mg, 0.3 mmol) in MeOH (5 mL) and the mixture stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL), 8-(1*H*-imidazol-1-ylcarbonyl)-2-(4-pyridinyl)quinoline (150 mg, 0.6 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated, the residue dissolved in DCM (20 mL) and washed with water (3 × 15 mL). The organic layer was separated, dried, and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–3%) MeOH/DCM, to give compound **86** (165 mg, 100%) as a red solid, mp (DCM/hexane) 178–180 °C; ¹H NMR δ 10.81 (br s, 1 H, NH), 8.77–8.83 (m, 3 H, ArH), 8.40 (br s, 1 H, NH), 8.32 (d, *J* = 8.6 Hz, 1 H, ArH), 8.13 (d, *J* = 8.9 Hz, 1 H, ArH), 7.87–7.93 (m, 5 H, ArH), 7.65 (t, *J* = 7.7 Hz, 1 H, ArH), 7.21 (d, *J* = 8.8 Hz, 1 H, ArH), 3.72 (br q, *J* = 6.3 Hz, 2 H, CH₂), 3.49 (br q, *J* = 5.3 Hz, 2 H, CH₂), 2.54 (t, *J* = 7.2 Hz, 2 H, CH₂), 2.47 (s, 3 H, CH₃), 2.46 (t, *J* = 6.2 Hz, 2 H, CH₂), 2.24 (s, 3 H, CH₃), 2.01 (br quin, *J* = 5.5 Hz, 2 H, CH₂), 1.72 (br quin *J* = 6.1 Hz, 2 H, CH₂); ¹³C NMR δ 165.9, 154.4 (2), 150.8, 149.8, 147.6, 146.2, 145.3, 138.8, 138.1, 134.1, 131.1, 130.3, 128.8, 128.5, 127.8, 127.2, 121.7 (2), 121.3, 118.5, 116.0, 56.7, 55.7, 41.9, 41.3, 38.2, 27.7, 25.5, 22.2; HRMS (FAB⁺) calcd for C₃₀H₃₃N₈O₃ (MH⁺) *m/z* 553.2676, found 553.2673. Anal. calcd for C₃₀H₃₂N₈O₃·½H₂O: C, 64.2; H, 5.9; N, 20.0; found: C, 64.4; H, 6.1; N, 19.5%.

Example AC

***N*-[3-(Methyl{3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-1-phenazinecarboxamide (87).** Aqueous NH₃ (6 mL) was added to a solution of dioxide **84** (145 mg, 0.35 mmol) in MeOH (10 mL) and stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL), 1-(1*H*-imidazol-1-ylcarbonyl)phenazine (183 mg, 0.68 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–4%) MeOH/DCM, to give compound **87** (181 mg, 98%) as a red solid, mp (DCM/hexane) 111–114 °C; ¹H NMR δ 10.82 (br s, 1 H, NH), 8.91 (dd, *J* = 7.2, 1.5

Hz, 1 H, ArH), 8.59 (br s, 1 H, NH), 8.33 (dd, $J = 8.6, 1.5$ Hz, 1 H, ArH), 8.21–8.25 (m, 2 H, ArH), 8.13 (d, $J = 8.9$ Hz, 1 H, ArH), 7.93 (dd, $J = 8.7, 7.1$ Hz, 1 H, ArH), 7.85–7.90 (m, 2 H, ArH), 7.83 (s, 1 H, ArH), 7.22 (dd, $J = 9.0, 1.7$ Hz, 1 H, ArH), 3.78 (br q, $J = 6.4$ Hz, 2 H, CH₂), 3.65 (br q, $J = 5.9$ Hz, 2 H, CH₂), 2.69 (t, $J = 7.3$ Hz, 2 H, CH₂), 2.63 (t, $J = 6.1$ Hz, 2 H, CH₂), 2.45 (s, 3 H, CH₃), 2.36 (s, 3 H, CH₃), 2.10 (br quin, $J = 7.1$ Hz, 2 H, CH₂), 1.89 (br quin, $J = 6.2$ Hz, 2 H, CH₂); ¹³C NMR δ 165.1, 149.8, 147.8, 143.3, 142.9, 141.4, 140.8, 138.0, 135.0, 134.9, 133.3, 131.6, 131.0, 130.0 (2), 129.0, 129.7, 129.0, 121.3, 115.7, 56.9, 55.8, 42.0, 41.6, 38.2, 27.5, 25.5, 22.2; HRMS (FAB⁺) calcd for C₂₈H₃₁N₈O₃ (MH⁺) m/z 527.2519, found 527.2506. Anal. calcd for C₂₈H₃₀N₈O₃·¼H₂O: C, 63.3; H, 5.8; N, 21.1; found: C, 63.3; H, 5.8; N, 21.5%.

Example AD

9-Methyl-N-[3-({3-[(6-methyl-1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}amino)propyl]-1-phenazinecarboxamide (88). Aqueous NH₃ (5 mL) was added to a solution of dioxide **84** (126 mg, 0.3 mmol) in MeOH (5 mL) and the reaction mixture was stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL), 1-(1*H*-imidazol-1-ylcarbonyl)-9-methylphenazine (172 mg, 0.6 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–4%) MeOH/DCM, to give compound **88** (147 mg, 91%) as a red solid, mp (DCM/hexane) 80–83 °C; ¹H NMR δ 10.99 (t, $J = 5.2$ Hz, 1 H, NH), 8.96 (dd, $J = 7.1, 1.5$ Hz, 1 H, ArH), 8.43 (br s, 1 H, NH), 8.36 (dd, $J = 8.5, 1.0$ Hz, 1 H, ArH), 8.14 (d, $J = 8.9$ Hz, 1 H, ArH), 8.10 (dd, $J = 8.8, 1.2$ Hz, 1 H, ArH), 7.96 (dd, $J = 8.8, 7.1$ Hz, 1 H, ArH), 7.96 (dd, $J = 8.7, 6.8$ Hz, 1 H, ArH), 7.77 (dd, $J = 8.7, 6.8$ Hz, 1 H, ArH), 7.69–7.73 (m, 1 H, ArH), 7.22 (dd, $J = 8.9, 1.8$ Hz, 1 H, ArH), 3.77 (br q, $J = 6.6$ Hz, 2 H, CH₂), 3.66 (br q, $J = 6.1$ Hz, 2 H, CH₂), 2.93 (s, 3 H, CH₃), 2.62 (t, $J = 7.2$ Hz, 2 H, CH₂), 2.58 (t, $J = 6.0$ Hz, 2 H, CH₂), 2.48 (s, 3 H, CH₃), 2.32 (s, 3 H, CH₃), 2.08 (br quin, $J = 7.2$ Hz, 2 H, CH₂), 1.86 (br quin, $J = 6.2$ Hz, 2 H, CH₂); ¹³C NMR δ 165.2, 149.8, 147.8, 143.1 (2), 141.0, 139.7, 138.1, 136.6, 135.0, 133.3, 131.0, 130.9, 130.0, 129.5, 129.0, 128.5, 127.7, 121.3, 115.8, 56.4, 55.8, 42.0, 41.3, 38.3, 27.9, 25.7, 22.3, 18.1; HRMS (FAB⁺) calcd for C₂₉H₃₃N₈O₃ (MH⁺) m/z 541.2676, found 541.2668.

Example AE

N-{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}(methylamino)ethyl-4-acridinecarboxamide (**95**).

- 5 *tert*-Butyl 2-[(2-Aminoethyl)(methylamino)ethylcarbamate (**90**). A solution of (BOC)₂O (9.60 g, 44 mmol) in THF (50 mL) was added over a period of 2 h to a solution of bis(diethylamino)methylamine (**89**) (10.32 g, 88 mmol) in THF (50 mL) at 0 °C. The reaction mixture stirred for 30 min then allowed to warm to 20 °C and stirred for 20 h. The reaction mixture was partitioned between DCM and saturated
- 10 aqueous NaCl, the organic layer separated and the aqueous layer further extracted with DCM (3 × 25 mL). The combined organic extract was dried and the solvent evaporated at 30 °C to give carbamate **90** (8.79 g, 46%) as a colourless oil, which was used without further purification.
- 15 *tert*-Butyl 2-(Methyl{2-[(1-oxido-1,2,4-benzotriazin-3-yl)amino]ethyl}amino)ethylcarbamate (**91**). A solution of chloride **3** (2.0 g, 11.0 mmol), carbamate **90** (2.9 g, 13.3 mmol) and triethylamine (3.0 mL, 22.1 mmol) in DME (50 mL) was heated at 85 °C for 3 h. The solvent was evaporated and the residue was partitioned between DCM (100 mL) and aqueous NH₃ (50 mL). The
- 20 DCM layer was separated, the aqueous layer further extracted with DCM (3 × 30 mL), the combined organic fraction dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM, to give (i) starting material **3** (500 mg, 25%) and (ii) carbamate **91** (2.1 g, 52%) as a yellow solid, mp (DCM/hexane) 122–124 °C; ¹H NMR [(CD₃)₂SO]
- 25 δ 8.13 (dd, *J* = 8.6, 1.0 Hz, 1 H, H-8), 7.78 (ddd, *J* = 8.4, 7.0, 1.6 Hz, 1 H, H-6), 7.71 (br s, 1 H, NH), 7.52 (d, *J* = 8.2 Hz, 1 H, H-5), 7.33 (ddd, *J* = 7.8, 7.1, 1.2 Hz, 1 H, H-7), 6.61 (br s, 1 H, NH), 3.43 (br q, *J* = 6.0 Hz, 2 H, CH₂), 3.01 (br q, *J* = 6.2 Hz, 2 H, CH₂), 2.57 (t, *J* = 6.6 Hz, 2 H, CH₂), 2.42 (t, *J* = 6.7 Hz, 2 H, CH₂), 2.23 (s, 3 H, CH₃), 1.35 (s, 9 H, 3 × CH₃); ¹³C NMR [(CD₃)₂SO] δ 158.8, 155.4, 148.2, 135.6,
- 30 129.9, 126.0, 124.4, 119.8, 77.4, 56.4, 55.5, 41.8, 38.5, 37.8, 28.1 (3); HRMS (FAB⁺) calcd for C₁₇H₂₇N₆O₃ (MH⁺) *m/z* 363.2145, found 363.2144. Anal. calcd for C₁₇H₂₆N₆O₃: C, 56.3; H, 7.2; N, 23.2; found: C, 56.5; H, 7.3; N, 23.3%.

***N*¹-(2-Aminoethyl)-*N*¹-methyl-*N*²-(1-oxido-1,2,4-benzotriazin-3-yl)-1,2-**

ethanediamine (92). Carbamate **91** (2.14 g, 5.9 mmol) was dissolved in methanolic HCl (30 mL) and stirred 20 h at 20 °C. Excess reagent and solvent were evaporated and the residue was partitioned between DCM and aqueous NH₃. The organic fraction
 5 was separated and the aqueous fraction was further extracted with DCM (4 × 30 mL). The combined organic fraction was dried and the solvent evaporated to give amine **92** (1.55 g, 100%) as yellow solid which was used without further purification, ¹H NMR [(CD₃)₂SO] δ 8.13 (dd, *J* = 8.6, 1.3 Hz, 1 H, H-8), 7.78 (ddd, *J* = 7.7, 7.0, 1.4 Hz, 2 H, H-6, NH), 7.57 (d, *J* = 8.4 Hz, 1 H, H-5), 7.33 (ddd, *J* = 7.8, 7.1, 1.3 Hz, 1 H, H-7),
 10 3.42–3.68 (m, 2 H, CH₂), 3.20–3.40 (m, 2 H, CH₂), 2.58 (q, *J* = 6.9 Hz, 2 H, CH₂), 2.37 (t, *J* = 6.5 Hz, 2 H, CH₂), 2.22 (s, 3 H, CH₃), 1.42 (br s, 2 H, NH₂).

2,2,2-Trifluoro-*N*-[2-(methyl{2-[(1-oxido-1,2,4-benzotriazin-3-

yl)amino]ethyl}amino)ethyl]acetamide (93). CF₃CO₂Et (2.05 mL, 17.2 mmol) and
 15 H₂O (0.31 mL, 17.2 mmol) was added to a solution of amine **92** (1.5 g, 5.7 mmol) in CH₃CN (50 mL) and the reaction mixture was heated at reflux for 48 h. The reaction mixture was evaporated and residue partitioned between DCM and aqueous NaHCO₃. The DCM layer was separated and the aqueous layer was further extracted with DCM (5 × 30 mL). The combined organic fraction was dried and the solvent evaporated to
 20 give trifluoroacetamide **93** (1.80 g, 88%) as a yellow solid, mp (DCM/hexane) 141–143 °C; ¹H NMR [(CD₃)₂SO] δ 9.29 (br s, 1 H, NH), 8.13 (dd, *J* = 8.6, 1.3 Hz, 1 H, H-8), 7.78 (ddd, *J* = 8.4, 7.0, 1.5 Hz, 1 H, H-6), 7.68 (br s, 1 H, NH), 7.57 (d, *J* = 8.4 Hz, 1 H, H-5), 7.33 (ddd, *J* = 8.5, 7.1, 1.3 Hz, 1 H, H-7), 3.42 (br q, *J* = 6.3 Hz, 2 H, CH₂),
 3.30 (br q, *J* = 6.8 Hz, 2 H, CH₂), 2.61 (t, *J* = 6.7 Hz, 2 H, CH₂), 2.56 (t, *J* = 6.7 Hz, 2
 25 H, CH₂), 2.27 (s, 3 H, CH₃); ¹³C NMR [(CD₃)₂SO] δ 158.8, 156.1 (q, *J* = 36 Hz) 148.3, 135.6, 130.0, 125.9, 124.4, 119.8, 115.8 (q, *J* = 288 Hz), 55.4, 55.1, 41.7, 38.4, 37.2; HRMS (FAB⁺) calcd for C₁₄H₁₈F₃N₆O₂ (MH⁺) *m/z* 359.1443, found 359.1451. Anal. calcd for C₁₄H₁₇F₃N₆O₂: C, 46.9; H, 4.8; N, 23.5; F, 15.9; found: C, 47.2; H, 4.9; N, 23.6; F, 15.8%.

***N*-{2-[2-[(1,4-Dioxido-1,2,4-benzotriazin-3-**

yl)amino]ethyl}(methylamino)ethyl}-2,2,2-trifluoroacetamide (94). A solution of trifluorooperacetic acid [prepared from trifluoroacetic anhydride (6.8 mL, 49 mmol)

and 70% H₂O₂ (2.0 mL, 49 mmol) in DCM (20 mL)] was added to a solution of trifluoroacetamide **93** (1.75 g, 4.9 mmol) and trifluoroacetic acid (0.8 mL, 9.8 mmol) in DCM (20 mL) and the reaction mixture was stirred at 20 °C for 5 h. The reaction mixture was slowly added to a cooled solution of aqueous NaHCO₃ (100 mL). The DCM layer was separated and the aqueous layer was further extracted with DCM (5 × 30 mL). The combined organic fraction was dried and the solvent was evaporated. The residue was purified by chromatography, eluting with a gradient (0–4%) of DCM/MeOH, to give (i) starting material **93** (100 mg, 6%); and (ii) dioxide **94** (859 mg, 47%) as a red solid, mp (DCM/hexane) 141–144 °C; ¹H NMR [(CD₃)₂SO] δ 9.28 (br s, 1 H, NH), 8.20 (d, *J* = 9.1 Hz, 1 H, H-5), 8.12 (d, *J* = 8.6 Hz, 1 H, H-8), 8.03 (t, *J* = 5.8 Hz, 1 H, NH), 7.96 (ddd, *J* = 8.6, 7.2, 1.3 Hz, 1 H, H-6), 7.56 (ddd, *J* = 8.6, 7.1, 1.3 Hz, 1 H, H-7), 3.48 (br q, *J* = 6.3 Hz, 2 H, CH₂), 3.31 (br q, *J* = 6.3 Hz, 2 H, CH₂), 2.63, (t, *J* = 6.6 Hz, 2 H, CH₂), 2.54 (t, *J* = 6.8 Hz, 2 H, CH₂), 2.27 (s, 3 H, CH₃); ¹³C NMR [(CD₃)₂SO] δ 156.1 (q, *J* = 36 Hz), 149.7, 138.0, 135.4, 129.9, 126.9, 121.0, 115.8 (q, *J* = 288 Hz), 116.7, 55.4, 55.0, 41.6, 38.3, 37.1; HRMS (FAB⁺) calcd for C₁₄H₁₈F₃N₆O₃ (MH⁺) *m/z* 375.1393, found 375.1392. Anal. calcd for C₁₄H₁₇F₃N₆O₃: C, 44.9; H, 4.6; N, 22.5; found: C, 44.8; H, 4.6; N, 22.5%.

***N*-{2-[{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-**

yl)amino]ethyl}(methylamino)ethyl}-4-acridinecarboxamide (95**).** Aqueous NH₃ (6 mL) was added to a solution of dioxide **94** (125 mg, 0.33 mmol) in MeOH (6 mL) and the reaction mixture was stirred at 20 °C for 16 h. The solvent was evaporated, the residue dissolved in THF (5 mL), 4-(1*H*-imidazol-1-ylcarbonyl)acridine (180 mg, 0.66 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated, the residue was dissolved in DCM (20 mL) and washed with water (3 × 15 mL). The organic fraction was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–3%) MeOH/DCM, to give compound **95** (132 mg, 94%) as a red solid, mp (DCM/hexane) 160–162 °C; ¹H NMR δ 11.90 (br s, 1 H, NH), 8.95 (dd, *J* = 7.1, 1.5 Hz, 1 H, ArH), 8.72 (s, 1 H, ArH), 8.16 (d, *J* = 8.6 Hz, 1 H, ArH), 8.12 (d, *J* = 8.8 Hz, 1 H, ArH), 8.07 (dd, *J* = 8.4, 1.4 Hz, 1 H, ArH), 7.82–7.86 (m, 2 H, ArH), 7.78 (ddd, *J* = 7.7, 6.7, 1.5 Hz, 1 H, ArH), 7.70 (ddd, *J* = 7.8, 7.1, 1.3 Hz, 1 H, ArH), 7.64 (dd, *J* = 8.2, 7.1 Hz, 1 H, ArH), 7.46 (ddd, *J* = 7.5, 7.2, 0.7 Hz, 1 H, ArH), 7.40 (ddd, *J* =

7.9, 7.1, 1.3 Hz, 1 H, ArH), 7.29 (br s, 1 H, NH), 3.86 (br q, $J = 6.0$ Hz, 2 H, CH₂), 3.70 (br q, $J = 5.6$ Hz, 2 H, CH₂), 2.93 (t, $J = 6.4$ Hz, 2 H, CH₂), 2.89 (t, $J = 6.1$ Hz, 2 H, CH₂), 2.57 (s, 3 H, CH₃); ¹³C NMR δ 166.0, 149.7, 147.4, 146.5, 137.8, 137.4, 135.3, 135.0, 132.1, 131.1, 130.1, 128.8, 128.6, 127.9, 126.8, 126.7, 126.1, 125.8, 125.5, 121.5, 117.2, 56.5, 55.9, 42.5, 39.1, 37.8; HRMS (FAB⁺) calcd for C₂₆H₂₆N₇O₃ (MH⁺) m/z 484.2097, found 484.2102.

Example AF

***N*-{2-[(2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl)(methylamino)ethyl]-2-(4-pyridinyl)-8-quinolinecarboxamide (96)}.** Aqueous NH₃ (6 mL) was added to a solution of dioxide **94** (132 mg, 0.35 mmol) in MeOH (10 mL) and the reaction mixture was stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL), 8-(1*H*-imidazol-1-ylcarbonyl)-2-(4-pyridinyl)quinoline (150 mg, 0.6 mmol) was added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated, the residue was dissolved in DCM (20 mL) and washed with water (3 × 15 mL). The organic layer was separated, dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–3%) MeOH/DCM to give compound **96** (174 mg, 97%) as a red solid, mp (DCM/hexane) 130–135 °C; ¹H NMR δ 11.00 (t, $J = 5.0$ Hz, 1 H, NH), 8.83 (dd, $J = 5.3, 2.2$ Hz, 1 H, ArH), 8.81 (d, $J = 7.7$ Hz, 1 H, ArH), 8.81 (dd, $J = 4.7, 3.0$ Hz, 1 H, ArH), 8.29 (d, $J = 8.6$ Hz, 1 H, ArH), 8.23 (dd, $J = 8.1$ Hz, 1 H, ArH), 7.98 (d, $J = 9.7$ Hz, 1 H, ArH), 7.92–7.93 (m, 2 H, ArH), 7.87–7.90 (m, 2 H, ArH), 7.76 (ddd, $J = 6.1, 5.4, 2.2$ Hz, 1 H, ArH), 7.65 (dd, $J = 7.4, 6.5$ Hz, 1 H, ArH), 7.44 (ddd, $J = 7.9, 7.0, 1.3$ Hz, 1 H, ArH), 7.21 (br, 1 H, NH), 3.79 (br q, $J = 6.1$ Hz, 2 H, CH₂), 3.48 (br q, $J = 5.8$ Hz, 2 H, CH₂), 2.82 (t, $J = 6.3$ Hz, 2 H, CH₂), 2.73 (t, $J = 6.1$ Hz, 2 H, CH₂), 2.43 (s, 3 H, CH₃); ¹³C NMR δ 165.8, 154.6 (2), 150.7, 149.6, 146.4, 145.4, 138.8, 138.0, 135.2, 134.5, 131.3, 130.2, 129.5, 127.9, 127.1, 126.9, 121.8 (2), 121.6, 118.7, 117.3, 56.8, 56.1, 42.3, 38.8, 37.9; HRMS (FAB⁺) calcd for C₂₇H₂₇N₈O₃ (MH⁺) m/z 511.2206, found 511.2208. Anal. calcd for C₂₇H₂₆N₈O₃· $\frac{1}{4}$ H₂O: C, 63.0; H, 5.2; N, 21.8; found: C, 63.0; H, 5.2; N, 21.5%.

Example AG

N*-{2-[2-[(1,4-Dioxido-1,2,4-benzotriazin-3-*yl)amino]ethyl}(methyl)amino]ethyl}-1-phenazinecarboxamide (97).** AqueousNH₃ (6 mL) was added to a solution of dioxide **94** (120 mg, 0.32 mmol) in MeOH (10 mL) and reaction mixture was stirred at 20 °C for 18 h. The solvent was evaporated,the residue dissolved in DMF (5 mL), and 1-(1*H*-imidazol-1-ylcarbonyl)phenazine (172 mg, 0.64 mmol) added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient(0–1%) of aqueous NH₃/(0–3%) MeOH/DCM, to give compound **97** (137 mg, 88%)as a red solid, mp (DCM/hexane) 163–165 °C; ¹H NMR δ 11.07 (br s, 1 H, NH), 8.95(dd, *J* = 7.1, 1.5 Hz, 1 H, ArH), 8.31 (dd, *J* = 8.7, 1.5 Hz, 1 H, ArH), 8.07–8.11 (m, 2H, ArH), 8.03 (dd, *J* = 8.7, 0.7 Hz, 1 H, ArH), 7.93 (dd, *J* = 8.7, 7.1 Hz, 1 H, ArH),7.78 (ddd, *J* = 7.7, 6.8, 1.6, 1 H, ArH), 7.65–7.72 (m, 2 H, ArH), 7.60 (dd, *J* = 8.6, 0.9Hz, 1 H, ArH), 7.38 (ddd, *J* = 7.8, 7.0, 1.5 Hz, 1 H, ArH), 7.19 (br s, 1 H, NH), 3.85(br q, *J* = 5.8 Hz, 2 H, CH₂), 3.68 (br q, *J* = 5.6 Hz, 2 H, CH₂), 2.85–2.91 (m, 4 H, 2 ×CH₂), 2.57 (s, 3 H, CH₃); ¹³C NMR δ 165.0, 149.6, 143.4, 142.7, 141.3, 141.0, 137.6,

135.2 (2), 133.4, 131.2, 130.5, 130.0, 129.9, 129.6, 129.5, 128.8, 126.9, 121.3, 117.0,

56.2, 55.8, 42.3, 38.9, 37.8; HRMS (FAB⁺) calcd for C₂₅H₂₅N₈O₃ (MH⁺) *m/z*485.2050, found 485.2045. Anal calcd for C₂₅H₂₄N₈O₃: C, 62.0; H, 5.0; N, 23.1;

found: C, 61.7; H, 4.7; N 23.1%.

Example AH***N*-{2-[2-[(1,4-Dioxido-1,2,4-benzotriazin-3-****yl)amino]ethyl}(methyl)amino]ethyl}-9-methyl-1-phenazinecarboxamide (98).**Aqueous NH₃ (6 mL) was added to a solution of dioxide **94** (120 mg, 0.32 mmol) in

MeOH (10 mL) and the reaction mixture was stirred at 20 °C for 16 h. The solvent

was evaporated, the residue dissolved in THF (5 mL), and 1-(1*H*-imidazol-1-

ylcarbonyl)-9-methylphenazine (185 mg, 0.6 mmol) was added and the mixture

stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by

chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–3%)MeOH/DCM, to give compound **98** (128 mg, 80%) as a red solid, mp (DCM/hexane)161–163 °C; ¹H NMR δ 11.03 (br s, 1 H, NH), 8.94 (dd, *J* = 7.1, 1.5 Hz, 1 H, ArH),8.29 (dd, *J* = 8.6, 1.5 Hz, 1 H, ArH), 8.12 (dt, *J* = 8.8, 0.7 Hz, 1 H, ArH), 7.94–7.98(m, 1 H, ArH), 7.91 (dd, *J* = 8.6, 7.1 Hz, 1 H, ArH), 7.60–7.70 (m, 4 H, ArH), 7.78

(ddd, $J = 8.3, 7.3, 1.1$ Hz, 1 H, ArH), 7.26 (br s, 1 H, NH), 3.86 (br q, $J = 6.1$ Hz, 2 H, CH₂), 3.60 (br q, $J = 5.5$ Hz, 2 H, CH₂), 2.91 (s, 3 H, CH₃), 2.87 (t, $J = 6.2$ Hz, 2 H, CH₂), 2.80 (t, $J = 5.9$ Hz, 2 H, CH₂), 2.47 (s, 3 H, CH₃); ¹³C NMR δ 165.2, 149.5, 143.1, 141.0, 140.0, 137.6, 136.7, 135.2, 135.1, 133.3, 130.7, 130.6, 129.9 (2), 129.4, 127.6, 126.8, 121.3, 117.0, 56.6, 55.8, 42.3, 38.8, 37.7, 17.9, one resonance not observed; HRMS (FAB⁺) calcd for C₂₆H₂₇N₈O₃ (MH⁺) m/z 499.2206, found 499.2200. Anal. calcd for C₂₆H₂₆N₈O₃: C, 62.6; H, 5.3; N, 22.5; found: C, 62.2; H, 5.3; N, 22.4%.

10 Example AI

N-{2-[{2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}(methyl)amino]ethyl}-5-methyl-4-acridinecarboxamide (**99**).

Aqueous NH₃ (6 mL) was added to a solution of dioxido **94** (125 mg, 0.33 mmol) in MeOH (10 mL) and the reaction mixture was stirred at 20 °C for 24 h. The solvent was evaporated, the residue dissolved in THF (5 mL) and 4-(1*H*-imidazol-1-ylcarbonyl)-5-methylacridine (208 mg, 0.72 mmol) was added and the mixture stirred at 20 °C for 24 h. The solvent was evaporated, the residue dissolved in DCM (20 mL) and washed with water (3 × 15 mL). The organic layer was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–4%) MeOH/DCM, to give compound **99** (180 mg, 100%) as a red solid, mp (DCM/hexane) 148–152 °C; ¹H NMR δ 11.90 (br s, 1 H, NH), 8.94 (dd, $J = 7.1, 1.5$ Hz, 1 H, ArH), 8.70 (s, 1 H, ArH), 8.17 (d, $J = 8.2$ Hz, 1 H, ArH), 8.05 (dd, $J = 8.3, 1.4$ Hz, 1 H, ArH), 7.88 (d, $J = 8.2$ Hz, 1 H, ArH), 7.96 (d, $J = 7.9$, Hz, 1 H, ArH), 7.69–7.73 (m, 1 H, ArH), 7.60–7.65 (m, 2 H, ArH), 7.38–7.43 (m, 2 H, ArH), 7.33 (br s, 1 H, NH), 3.85 (br q, $J = 6.3$ Hz, 2 H, CH₂), 3.61 (br q, $J = 4.2$ Hz, 2 H, CH₂), 2.90 (s, 3 H, CH₃), 2.89 (t, $J = 6.7$ Hz, 2 H, CH₂), 2.56 (t, $J = 6.0$ Hz, 2 H, CH₂), 2.47 (s, 3 H, CH₃); ¹³C NMR δ 166.2, 149.7, 147.0, 145.4, 137.9, 137.8, 135.9, 135.2, 135.1, 132.1, 130.8, 130.1, 128.4, 126.7, 126.6, 126.2, 126.1, 125.8, 125.4, 121.5, 117.2, 56.9, 55.8, 42.3, 39.0, 37.9, 18.8; HRMS (FAB⁺) calcd for C₂₇H₂₈N₇O₃ (MH⁺) m/z 498.2254, found 498.2257. Anal. calcd for C₂₇H₂₇N₇O₃· $\frac{1}{4}$ H₂O: C, 64.6; H, 5.5; N, 19.5; found: C, 64.5; H, 5.5; N, 19.7%.

Example AJ

***N*-{3-[{3-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]propyl}(methyl)amino]propyl}-1-phenazinecarboxamide (100).**

Aqueous NH₃ (6 mL) was added to a solution of trifluoroacetamide **39** (283 mg, 0.70 mmol) in MeOH (10 mL) and the reaction mixture was stirred at 20 °C for 18 h. The solvent was evaporated, the residue dissolved in DMF (5 mL), 1-(1*H*-imidazol-1-ylcarbonyl)phenazine (283 mg, 1.05 mmol) added and the mixture stirred at 20 °C for 48 h. The solvent was evaporated and the residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–3%) MeOH/DCM, to give compound **100** (293 mg, 82%) as a red solid, mp (DCM/hexane) 129–130 °C; ¹H NMR δ 10.85, (br s, 1 H, NH), 8.93 (dd, *J* = 7.1, 1.4 Hz, 1 H, ArH), 8.52 (br, 1 H, NH), 8.33 (dd, *J* = 8.7, 1.4 Hz, 1 H, ArH), 8.21–8.27 (m, 2 H, ArH), 8.11 (d, *J* = 8.7 Hz, 1 H, ArH), 7.93 (dd, *J* = 8.6, 6.5 Hz, 1 H, ArH), 7.86–7.90 (m, 3 H, ArH), 7.70 (t, *J* = 7.8 Hz, 1 H, ArH), 7.42 (t, *J* = 7.8 Hz, 1 H, ArH), 3.77 (br q, *J* = 6.4 Hz, 2 H, CH₂), 3.66 (br q, *J* = 5.7 Hz, 2 H, CH₂), 2.68 (t, *J* = 7.3 Hz, 2 H, CH₂), 2.62 (t, *J* = 6.1 Hz, 2 H, CH₂), 2.36 (s, 3 H, CH₃), 2.10 (br quin, *J* = 7.1 Hz, 2 H, CH₂), 1.89 (br quin, *J* = 6.2 Hz, 2 H, CH₂); ¹³C NMR δ 165.0, 149.8, 143.4, 142.9, 141.4, 140.8, 138.2, 135.4, 135.1, 135.0, 133.4, 131.5, 130.9, 130.1 (2), 129.8, 129.0, 126.7, 121.6, 117.1, 56.7, 55.9, 42.1, 41.5, 38.2, 27.5, 25.6; HRMS (FAB⁺) calcd for C₂₇H₂₉N₈O₃ (M⁺) *m/z* 513.2363, found 513.2365. Anal. calcd for C₂₇H₂₈N₈O₃: C, 54.8; H, 6.0; N, 31.9; found: C, 55.1; H, 5.8; N, 32.3%.

Example AK

***N*¹-(2-aminoethyl)-*N*²-(1,4-dioxido-1,2,4-benzotriazin-3-yl)-1,2-ethanediamine (101).** A solution of carbamate (**36**) (252 mg, 0.54 mmol) in methanolic HCl was stirred at 20 °C for 24 h. Excess reagent and solvent were evaporated and the residue partitioned between aqueous NH₃ and DCM. The organic layer was separated and the aqueous layer was extracted with DCM (15 × 20 mL). The combined organic extract was dried, and the solvent evaporated to give amine **101** (109 mg, 76%) as a gum which was used without further purification, ¹H NMR δ 8.33 (d, *J* = 8.7 Hz, 1 H, ArH), 8.30 (d, *J* = 8.8 Hz, 1 H, ArH), 7.87 (ddd, *J* = 8.5, 7.1, 1.0 Hz, 1 H, ArH), 7.50 (ddd, *J* = 8.4, 7.1, 1.2 Hz, 1 H, ArH), 3.70 (br t, *J* = 5.9 Hz, 2 H, CH₂), 2.98 (br t, *J* = 5.9 Hz, 2 H, CH₂), 2.84 (br t, *J* = 5.6 Hz, 2 H, CH₂), 2.74 (br t, *J* = 5.6 Hz, 2 H, CH₂), 2 × NH, NH₂ not observed.

N-[2-({2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}amino)ethyl]-2-(4-pyridinyl)-8-quinolinecarboxamide (**102**). 8-(1*H*-Imidazol-1-ylcarbonyl)-2-(4-pyridinyl)quinoline (198 mg, 0.78 mmol) was added to a solution of amine (**101**) (105 mg, 0.39 mmol) in DMF (10 mL) and the reaction mixture was stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–4%) MeOH/DCM to give compound **102** (148 mg, 75%) as a red solid, mp (DCM/hexane) 160–165 °C; ¹H NMR [(CD₃)₂SO] δ 10.62 (t, *J* = 5.4 Hz, 1 H, NH), 8.75 (d, *J* = 6.1 Hz, 1 H, ArH), 8.75 (dd, *J* = 4.5, 1.8 Hz, 1 H, ArH), 8.68 (d, *J* = 8.7 Hz, 1 H, ArH), 8.57 (dd, *J* = 7.3, 1.6 Hz, 1 H, ArH), 8.28 (d, *J* = 8.7 Hz, 1 H, ArH), 8.21 (dd, *J* = 8.2, 1.6 Hz, 1 H, ArH), 8.18 (d, *J* = 6.2 Hz, 1 H, ArH), 8.15 (dd, *J* = 3.5, 1.7 Hz, 1 H, ArH), 8.11 (br s, 1 H, NH), 8.07 (dd, *J* = 8.6, 0.9 Hz, 1 H, ArH), 7.98 (dd, *J* = 8.8, 0.7 Hz, 1 H, ArH), 7.88 (ddd, *J* = 7.8, 7.0, 1.4 Hz, 1 H, ArH), 7.77 (dd, *J* = 8.0, 7.4 Hz, 1 H, ArH), 7.51 (ddd, *J* = 7.8, 7.0, 1.5 Hz, 1 H, ArH), 3.61 (br q, *J* = 5.8 Hz, 2 H, CH₂), 3.41–3.45 (m, 2 H, CH₂), 2.90 (t, *J* = 5.9 Hz, 2 H, CH₂), 2.87 (t, *J* = 6.2 Hz, 2 H, CH₂), NH not observed; HRMS (FAB⁺) calcd for C₂₆H₂₅N₇O₃ (MH⁺) *m/z* 497.2050, found 497.2058. Anal. calcd for C₂₆H₂₄N₈O₃: C, 62.9; H, 4.9; N, 22.7; found: C, 62.9; H, 4.9; N, 22.7.

20 Example AL

N-[2-({2-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]ethyl}amino)ethyl]-5-methyl-4-acridinecarboxamide (**103**). 4-(1*H*-Imidazol-1-ylcarbonyl)-5-methylacridine (245 mg, 0.86 mmol) was added to a solution of amine **101** (113 mg, 0.43 mmol) in DMF (5 mL) and the reaction mixture stirred at 20 °C for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of aqueous NH₃/(0–4%) MeOH/DCM, to give compound **103** (156 mg, 75%) as a red solid, mp (DCM/hexane) 135–140 °C; ¹H NMR [(CD₃)₂SO] δ 11.52 (t, *J* = 5.5 Hz, 1 H, CONH), 9.25 (s, 1 H, ArH), 8.74 (dd, *J* = 7.1, 1.6 Hz, 1 H, ArH), 8.36 (dd, *J* = 8.6, 1.5 Hz, 1 H, ArH), 8.18 (br, 1 H, NH), 8.10 (dd, *J* = 8.6, 0.9 Hz, 1 H, ArH), 8.03 (d, *J* = 8.4 Hz, 1 H, ArH), 7.96 (dd, *J* = 8.7, 1.2 Hz, 1 H, ArH), 7.90 (ddd, *J* = 7.8, 5.9, 1.4 Hz, 1 H, ArH), 7.73–7.77 (m, 2 H, ArH), 7.49–7.57 (m, 2 H, ArH), 3.65 (br q, *J* = 6.0 Hz, 2 H, CH₂), 3.47–3.51 (m, 2 H, CH₂), 2.92 (t, *J* = 6.0 Hz, 2 H, CH₂), 2.88 (s, 3 H, CH₃), 2.86 (t, *J* = 6.3 Hz, 2 H, CH₂), NH not observed;

^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 164.8, 149.8, 146.4, 144.6, 138.7, 137.8, 135.4, 135.2, 134.5, 132.6, 131.1, 129.7, 128.2, 126.7, 126.4, 126.2 (2), 125.5, 125.1, 120.9, 116.6, 48.4, 47.8, 40.4, 39.6, 18.2; HRMS (FAB $^+$) calcd for $\text{C}_{26}\text{H}_{26}\text{N}_7\text{O}_3$ (MH^+) m/z 484.2097, found 484.2094.

5

Example AM

***N*-{5-[4-(Dimethylamino)butanoyl]-1-methyl-1*H*-pyrrol-3-yl]-4-({4-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]butanoyl}amino)-1-methyl-1*H*-pyrrole-2-carboxamide (112).**

- 10 **Methyl 4-[(4-[(*tert*-Butoxycarbonyl)amino]-1-methyl-1*H*-pyrrol-2-yl)carbonyl]amino]-1-methyl-1*H*-pyrrole-2-carboxylate (106).** A solution of methyl 4-amino-1-methyl-1*H*-pyrrole-2-carboxylate (**104**) (Baird & Dervan, *J. Am. Chem. Soc.* 1996, 118, 6141–6146) (792 mg, 4.2 mmol) and 4-[(*tert*-butoxycarbonyl)amino]-1-methyl-1*H*-pyrrole-2-carboxylic acid (**105**) (Baird &
- 15 Dervan, *J. Am. Chem. Soc.* 1996, 118, 6141–6146) (1.0 g, 4.2 mmol) in DMF (13 mL) and DCM (3 mL) was treated with EDCI (1.5 g, 6.2 mmol) and DMAP (0.77 g, 5.0 mmol). The reaction mixture was stirred at 20 °C for 18 h, poured into 10% HCl (20 mL) and extracted with EtOAc (3 × 40 mL). The combined organic fraction was washed with saturated aqueous NaHCO_3 , dried and the solvent evaporated. The
- 20 residue was purified by chromatography, eluting with a gradient (0–50%) of EtOAc/pet. ether to give amide **106** (1.0 g, 64%) as a white solid, mp (DCM/pet. ether) 89–90 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 9.82 (s, 1 H, NH), 9.06 (br s, 1 H, NH), 7.44 (d, J = 1.9 Hz, 1 H, ArH), 6.90 (d, J = 2.0 Hz, 1 H, ArH), 6.88 (br s, 1 H, ArH), 6.83 (br s, 1 H, ArH), 3.83 (s, 3 H, CH_3), 3.80 (s, 3 H, CH_3), 3.74 (s, 3 H, CO_2CH_3), 1.46
- 25 (s, 9 H, 3 × CH_3); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 168.7, 158.3, 152.7, 122.9, 122.5, 122.3, 120.6, 118.4, 117.1, 108.3, 103.8, 78.2, 50.8, 36.0, 35.9, 28.1 (3); HRMS (EI^+) calcd for $\text{C}_{18}\text{H}_{24}\text{N}_4\text{O}_5$ (M^+) m/z 376.1747, found 376.1744.

- 30 **4-[(4-Amino-1-methyl-1*H*-pyrrol-2-yl)carbonyl]amino}-1-methyl-1*H*-pyrrole-2-carboxylic Acid (107).** A solution of LiOH (343 mg, 14.3 mmol) in water (7 mL) was added to a solution of amide **106** (1.0 g, 2.7 mmol) in THF/MeOH (3:1, 28 mL) and the mixture heated at 60 °C for 18 h. The mixture was cooled and diluted with EtOAc (150 mL). The aqueous layer was separated, adjusted to the pH 3 with 10% aqueous

HCl and extracted with EtOAc (3 × 40 mL). The combined organic fraction was dried and the solvent evaporated to give acid **107** (900 mg, 94%) as a white solid, mp (DCM/hexane) 138–142 °C; ¹H NMR [(CD₃)₂SO] δ 11.70 (br s, 1 H, CO₂H), 9.78 (s, 1 H, NH), 9.05 (s, 1 H, NH), 7.38 (d, *J* = 1.9 Hz, 1 H, ArH), 6.90 (br s, 1 H, ArH),
 5 6.82 (d, *J* = 2.0 Hz, 2 H, ArH), 3.81 (s, 3 H, CH₃), 3.80 (s, 3 H, CH₃), 1.46 (s, 9 H, 3 × CH₃); ¹³C NMR [(CD₃)₂SO] δ 171.8, 161.8, 158.3, 152.8, 122.6, 122.3, 120.1, 119.4, 117.0, 108.3, 103.7, 78.2, 36.0, 35.9, 28.1 (3); HRMS (FAB⁺) calcd for C₁₇H₂₃N₄O₅ (MH⁺) *m/z* 363.1669, found 363.1653.

10 ***tert*-Butyl 5-({[5-({[3-(Dimethylamino)propyl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-yl]amino}carbonyl)-1-methyl-1*H*-pyrrol-3-ylcarbamate (108).** A solution of acid **107** (900 mg, 2.5 mmol) was treated with EDCI (950 mg, 5.0 mmol), DMAP (758 mg, 6.2 mmol) and 3-dimethylaminopropylamine (507 mg, 5.0 mmol). The reaction mixture was stirred at 20 °C for 18 h, diluted with EtOAc (100 ml) and
 15 washed with 10% aqueous HCl (3 × 20 mL). The combined aqueous fraction was basified with aqueous NH₃, extracted with EtOAc (3 × 40 mL), dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM, to give amide **108** (847 mg, 76%) as a viscous oil, which solidified on standing, mp (DCM/pet.ether) 120–123 °C; ¹H NMR
 20 [(CD₃)₂SO] δ 9.77 (s, 1 H, NH), 9.04 (br s, 1 H, NH), 8.01 (t, *J* = 5.6 Hz, 1 H, NH), 7.15 (d, *J* = 1.8 Hz, 1 H, ArH), 6.87 (br s, 1 H, ArH), 6.81 (d, *J* = 1.8 Hz, 2 H, ArH), 3.80 (s, 3 H, CH₃), 3.79 (s, 3 H, CH₃), 3.18 (br q, *J* = 6.5 Hz, 2 H, CH₂), 2.24 (t, *J* = 7.1 Hz, 2 H, CH₂), 2.13 [s, 6 H, N(CH₃)₂] 1.89 (br quin, *J* = 7.0 Hz, 2 H, CH₂), 1.46 (s, 9 H, 3 × CH₃).

25 **Methyl 4-[(1-Oxido-1,2,4-benzotriazin-3-yl)amino]butanoate (109).** A solution of chloride **3** (1.57 g, 8.7 mmol), methyl 4-aminobutanoate hydrochloride (1.73 g, 11.4 mmol) and Et₃N (3.14 mmol, 22.5 mmol) in DME (50 mL) was heated at 90 °C for 6 h. The solvent was evaporated and the residue was partitioned between DCM (100
 30 mL) and water (50 mL). The organic fraction was separated and the aqueous layer was further extracted with DCM (4 × 30 mL). The combined organic fraction was dried, the solvent evaporated and the residue purified by chromatography, eluting with a gradient (0–2%) of MeOH/DCM, to give ester **109** (1.9 g, 81%) as a yellow solid,

mp (DCM/pet. ether) 122–126 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 8.11 (dd, $J = 8.6, 1.1$ Hz, 1 H, H-8), 7.90 (br s, 1 H, NH), 7.76 (ddd, $J = 7.7, 7.1, 1.5$ Hz, 1 H, H-6), 7.54 (br d, $J = 8.3$ Hz, 1 H, H-5), 7.31 (ddd, $J = 7.9, 7.0, 1.2$ Hz, 1 H, H-7), 3.58 (s, 3 H, OCH_3), 3.34–3.38 (m, 2 H, CH_2), 2.41 (t, $J = 7.4$ Hz, 2 H CH_2), 1.86–1.91 (m, 2 H, CH_2); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 173.2, 159.1, 148.4, 138.2, 135.7, 126.1, 124.6, 120.0, 51.3, 40.0, 30.8, 24.0; HRMS (EI^+) calcd for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_3$ (M^+) m/z 262.1066, found 262.1066. Anal. calcd for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_3$: C, 55.0; H, 5.4; N, 21.4; found: C, 55.1; H, 5.4; N, 21.4%.

Methyl 4-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]butanoate (110). Hydrogen peroxide (70%, 3.1 mL, 65 mmol) was added dropwise to a stirred solution of trifluoroacetic anhydride (9.0 mL, 65 mmol) in DCM (15 mL) at 0 °C and the solution stirred at 0 °C for 10 minutes. The solution was added to a solution of ester **109** (1.7 g, 6.5 mmol) in DCM (30 mL) at 20 °C and stirred for 16 h. The reaction mixture was poured into saturated aqueous NaHCO_3 (100 mL), the organic layer separated and the aqueous layer further extracted with DCM (3×30 mL). The combined organic fraction was dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–2%) of MeOH/DCM, to give (i) starting material **109** (610 mg, 36%); and (ii) 1,4-dioxide **110** (592 mg, 33%) as a red solid, mp (DCM/pet. ether) 169–171 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 8.33 (t, $J = 4.1$ Hz, 1 H, NH), 8.20 (dd, $J = 8.8, 0.7$ Hz, 1 H, H-8), 8.12 (dd, $J = 8.7, 0.7$ Hz, 1 H, H-5), 7.93 (ddd, $J = 7.8, 7.1, 1.4$ Hz, H-6), 7.56 (ddd, $J = 7.8, 7.1, 1.3$ Hz, 1 H, H-7), 3.59 (s, 3 H, OCH_3), 3.42 (br q, $J = 6.6$ Hz, 2 H, CH_2), 2.40 (t, $J = 7.4$ Hz, 2 H, CH_2), 1.88 (br quin, $J = 7.1$ Hz, 2 H, CH_2); ^{13}C NMR $[(\text{CD}_3)_2\text{SO}]$ δ 173.0, 149.8, 138.2, 135.4, 129.9, 126.9, 121.1, 116.8, 51.2, 39.9, 30.5, 23.9; HRMS (EI^+) calcd for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_4$ (M^+) m/z 278.1015, found 278.1014. Anal. calcd for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_4$: C, 51.8; H, 5.1; N, 20.1; found: C, 51.6; H, 4.9; N, 20.1%.

4-[(1,4-Dioxido-1,2,4-benzotriazin-3-yl)amino]butanoic Acid (111). A mixture of 1,4-dioxide **110** (351 mg, 1.26 mmol) and 1 N NaOH (6.3 mL, 6.30 mmol) in MeOH (20 mL) was stirred at 20 °C for 18 h. 10% Aqueous HCl (7 mL) was added and MeOH was evaporated. The resulting red precipitate was filtered, washed with water and dried to give acid **111** (270 mg, 81%) yield, mp (H_2O) 185–188 °C; ^1H NMR $[(\text{CD}_3)_2\text{SO}]$ δ 8.82 (br s, 1 H, NH), 8.19 (dd, $J = 8.8, 0.6$ Hz, 1 H, H-8), 8.11 (dd, $J =$

8.4, 0.9 Hz, 1 H, H-5), 7.92 (ddd, $J = 7.8, 7.1, 1.4$ Hz, 1 H, H-6), 7.54 (ddd, $J = 7.8, 7.2, 1.3$ Hz, 1 H, H-7), 3.38 (t, $J = 6.9$ Hz, 2 H, CH₂), 1.99 (t, $J = 7.0$ Hz, 1 H, CH₂), 1.79 (br quin, $J = 7.0$ Hz, 2 H, CH₂); HRMS (EI) calcd for C₁₁H₁₂N₄O₄ (M⁺) m/z 264.0845, found 264.0850. Anal. calcd for C₁₁H₁₂N₄O₄: C, 50.0, H, 4.6, N, 21.2; found: C, 50.1; H, 4.5, N, 21.2%.

***N*-{5-[4-(Dimethylamino)butanoyl]-1-methyl-1*H*-pyrrol-3-yl}-4-[(1,4-dioxido-1,2,4-benzotriazin-3-yl)amino]butanoyl}amino)-1-methyl-1*H*-pyrrole-2-carboxamide (112).** Carbamate 108 (166 mg, 0.37 mmol) was dissolved in HCl/MeOH (3 mL) and stirred for 16 h. The solvent was evaporated and the residue was dissolved in MeOH (5 mL) and evaporated. This process was repeated two more times. The residue was dissolved in DMF (5 mL) and DCM (2 mL) and acid 111 (264 mg, 0.38 mmol), EDCI (146 mg, 0.76 mmol) and DMAP (93 mg, 0.76 mmol) were added and the mixture stirred for 16 h at 20 °C. The solvent was evaporated and the residue was partitioned between DCM and aqueous NH₃. The resulting precipitate was collected by filtration and purified by chromatography, eluting with a gradient (0–1%) of aqueous NH₃/(0–5%) MeOH/DCM, to give compound 112 (21 mg, 9%) as an orange solid, mp (DCM/pet. ether) 140–145 °C; ¹H NMR [(CD₃)₂SO] δ 9.81 (s, 1 H, ArH), 9.79 (s, 1 H, ArH), 8.35 (t, $J = 6.1$ Hz, 1 H, NH), 8.21 (d, $J = 8.5$ Hz, 1 H, H-8), 8.14 (d, $J = 8.1$ Hz, 1 H, H-5), 8.03 (t, $J = 5.7$ Hz, 1 H, NH), 7.94 (ddd, $J = 7.8, 7.1, 0.8$ Hz, 1 H, H-6), 7.59 (ddd, $J = 7.9, 7.1, 1.3$ Hz, 1 H, H-7), 7.25 (d, $J = 1.8$ Hz, 1 H, ArH), 7.13 (d, $J = 1.8$ Hz, 1 H, ArH), 6.85 (d, $J = 1.8$ Hz, 1 H, ArH), 6.81 (d, $J = 1.8$ Hz, 1 H, ArH), 3.81 (s, 3 H, CH₃), 3.79 (s, 3 H, CH₃), 3.46 (br q, $J = 6.6$ Hz, 2 H, CH₂), 3.19 (br q, $J = 6.5$ Hz, 2 H, CH₂), 2.32 (br q, $J = 6.9$ Hz, 4 H, 2 × CH₂), 2.19 [s, 6 H, N(CH₃)₂], 1.93 (br quin, $J = 7.2$ Hz, 2 H, CH₂), 1.63 (br quin, $J = 7.0$ Hz, 2 H, CH₂), 3-NH not observed; ¹³C NMR [(CD₃)₂SO] δ 169.0, 161.1, 158.3, 149.7, 138.1, 135.3, 129.8, 126.8, 122.9, 122.6, 121.9, 121.8, 121.0, 118.0, 117.6, 116.8, 103.9 (2), 56.8, 44.9 (2), 40.5, 36.9, 35.9, 35.8, 32.9, 26.9, 25.0; HRMS (FAB⁺) calcd for C₂₈H₃₇N₁₀O₅ (MH⁺) m/z 593.2948, found 593.2953. Anal. calcd for C₂₈H₃₆N₁₀O₅·H₂O: C, 55.1; H, 6.3; N, 22.9; found: C, 55.1; H, 6.6, N, 22.2%.

Example AN

***tert*-Butyl 2-[(3-Ethyl-1,4-dioxido-1,2,4-benzotriazin-7-yl)oxy]ethylcarbamate (117).**

***N*-{2-[(3-Amino-1-oxido-1,2,4-benzotriazin-7-yl)oxy]ethyl}-2,2,2-**

trifluoroacetamide (113). A mixture of compound **46** (520 mg, 3.0 mmol), K₂CO₃

5 (833 mg, 6.0 mmol) and *N*-(2-bromoethyl)-2,2,2-trifluoroacetamide (1.25 g, 4.0 mmol) in DMF (20 mL) was stirred at 100 °C for 16 h. The solvent was evaporated and the residue suspended in water. The suspension was extracted with EtOAc (3 × 50 mL), the organic fraction dried and the solvent evaporated. The residue was purified by chromatography, eluting with 5% MeOH/DCM, to give compound **113** (639 mg, 10 66%) as a tan solid, mp (DCM/pet. ether) 234–236 °C. Anal. calcd for C₁₁H₁₀F₃N₅O₃: C, 41.7; H, 3.2; N, 22.1; F, 18.0; found: C, 41.9; H, 3.0; N, 21.9; F, 17.5%.

***N*-{2-[(3-Chloro-1-oxido-1,2,4-benzotriazin-7-yl)oxy]ethyl}-2,2,2-**

trifluoroacetamide (114). A solution of NaNO₂ (652 mg, 9.5 mmol) in water (20 15 mL) was added dropwise to a stirred suspension of amine **113** (1.5 g, 4.7 mmol) in 2 M HCl (75 mL) at 0 °C and the mixture stirred at 20 ° for 16 h. The suspension was filtered, the solid washed with water (2 × 10 mL) and dried to give 2,2,2-trifluoro-*N*-{2-[(3-hydroxy-1-oxido-1,2,4-benzotriazin-7-yl)oxy]ethyl}acetamide (1.44 g, 100%) as a tan solid, mp 202–204 °C. Anal. calcd for C₁₁H₉F₃N₄O₄: C, 41.5; H, 2.9; N, 17.6; 20 F, 17.9; found: C, 41.8; H, 2.9; N, 17.4; F, 17.6%.

A mixture of the 3-hydroxide (1.39 g, 4.1 mmol) and POCl₃ (15 mL) was stirred at 100 °C for 2 h. The solution was cooled and poured into ice/water and stirred for 30 min. The precipitate was filtered, washed with water, and dried. The solid was purified by chromatography, eluting with a gradient (0–10%) of EtOAc/DCM, to give 25 chloride **114** (1.37 g, 100%) as a tan solid, mp 179–181 °C. Anal. calcd for C₁₁H₈ClF₃N₄O₃: C, 39.2; H, 2.4; N, 16.6; F, 16.9; found: C, 39.5; H, 2.5; N, 16.7; F, 16.9%.

***N*-{2-[(3-Ethyl-1-oxido-1,2,4-benzotriazin-7-yl)oxy]ethyl}-2,2,2-**

30 **trifluoroacetamide (115).** Pd(PPh₃)₄ (198 mg, 0.17 mmol) was added to a purged solution of chloride **114** (1.16 g, 3.4 mmol) and Et₄Sn (0.82 mL, 4.1 mmol) in DME (50 mL) and the mixture heated at reflux temperature under N₂ for 16 h. The solvent was evaporated and the residue purified by chromatography, eluting with 1%

MeOH/DCM, to give compound **115** (896 mg, 79%) as a cream solid, mp (DCM) 155–157 °C. Anal. calcd for C₁₃H₁₃F₃N₄O₃: C, 47.3; H, 4.0; N, 17.0; found: C, 47.4; H, 4.2; N, 17.1%.

5 ***tert*-Butyl 2-[(3-Ethyl-1-oxido-1,2,4-benzotriazin-7-yl)oxy]ethylcarbamate (116).**

A solution of compound **115** (370 mg, 1.1 mmol) in 0.5 M K₂CO₃ solution (15 mL) was stirred at 20 °C for 16 h. The solution was extracted with CHCl₃ (3 × 30 mL), the organic fraction dried and the solvent evaporated. The residue was dissolved in THF (50 mL) and di-*tert*-butyl dicarbonate (367 mg, 1.68 mmol) added and the solution
10 stirred at 20 °C for 5 h. The solution was partitioned between EtOAc and water, the organic fraction dried and the solvent evaporated. The residue was purified by chromatography, eluting with 10% EtOAc/DCM, to give carbamate **116** (330 mg, 88%) as a white solid, mp. 101–103 °C.

15 ***tert*-Butyl 2-[(3-Ethyl-1,4-dioxido-1,2,4-benzotriazin-7-yl)oxy]ethylcarbamate (117).**

A mixture of 1-oxide **116** (300 mg, 0.9 mmol) and MCPBA (663 mg, 2.7 mmol) in DCM (20 mL) was stirred at 20 °C for 36 h. The solution was partitioned between dilute aqueous NH₃ and DCM, the organic fraction dried and the solvent evaporated. The residue was purified by chromatography, eluting with a gradient (0–
20 20%) of EtOAc/DCM, to give 1,4-dioxide (239 mg, 76%) as a red powder, mp (DCM/pet. ether) 111–113 °C.

Example AO: Cytotoxicity of Compounds

**Evaluation of the cytotoxicity of compounds by clonogenic assay under aerobic
25 and hypoxic conditions.**

Compounds representative of the invention were evaluated under both aerobic and hypoxic conditions in clonogenic assays, using three cell lines: human colon carcinoma HT-29, murine SCCVII, and human lung adenocarcinoma LXFL.

30 Clonogenic survival was determined using aerobic and hypoxic SCCVII cell suspensions. Drug exposures were performed using continuously stirred and gassed single cell suspensions (10⁶ cells/mL) at 37 °C, equilibrated with 5% CO₂ in air or N₂ for 60 min before drug addition. After a 60 min drug exposure cells were washed by centrifugation and plated to determine colony formation. Cytotoxicity was measured

as the concentration required to reduce plating efficiency to 10% of controls (C_{10}). The hypoxic cytotoxicity ratio (HCR) was determined as the ratio of the C_{10} values under aerobic and hypoxic conditions. The relative hypoxic toxicity (RHT) was determined as the ratio of hypoxic TPZ C_{10} to hypoxic BTO C_{10} . The results of these

5 assays are given in Table 2. Abbreviations used in Table 2 are:

C_{10} = The concentration of drug (in micromolar) to reduce viable cell numbers to 10% of those of control cell cultures grown under the same conditions but not exposed to drug

RHT = Relative hypoxic toxicity is defined as the ratio of concentrations of

10 Tirapazamine/test compound to give equal cell killing under hypoxic conditions.

HCR = Hypoxic cytotoxicity ratio is defined as the ratio of drug concentrations under aerobic and hypoxic condition to produce equal cell survival (10%) determined by clonogenic assay

15 **Table 2. Cytotoxicities of compounds of the invention under hypoxic conditions, hypoxic toxicity relative to Tirapazamine (RHT) and hypoxic selectivity (HCR) in clonogenic assay**

HT 29 cells			
compound	C_{10} hypoxic (μ M)	RHT	HCR
11	0.12	416	83.0
30	0.9	78	33.0
SCVIII cells			
compound	C_{10} hypoxic (μ M)	RHT	HCR
	SN (hypoxic)		
11	0.48	16.7	20.0
17	4.8	1.94	>6.3
30	0.3	20	21.3
41	0.8	12.5	52.5

44	0.16	56.3	>187
43	1.4	5.7	10
45	0.31	29	23.9
55	1.1	10	63.6
95	1.0	11	400
96	2.3	3.9	65
99	0.29	17.2	176
LXFL cells			
compound	C₁₀ hypoxic (μM)	RHT	HCR
11	0.04	450	35.0
30	0.4	50	12.5
41	0.4	37.5	50

The results of Table 2 clearly show that the compounds of the invention show large increases in cytotoxicity compared with Tirapazamine, while retaining selective killing under hypoxic conditions.

Example AP: Cytotoxicity of Compounds

Evaluation of the cytotoxicity of compounds by proliferation assay (IC₅₀) under aerobic and hypoxic conditions.

Compounds representative of the invention were evaluated under both aerobic and hypoxic conditions in a proliferation assay (IC₅₀), using two cell lines: human colon carcinoma HT-29, and human cervical carcinoma SiHa.

Drug exposures were performed in 96-well plates (Nunc) using either a 37 °C humidified incubator (20% O₂, 5% CO₂) or in the incubator compartment (37 °C) of an anaerobic chamber (Shell Lab) where palladium catalyst scrubbed gas (90% N₂, 5% H₂, 5% CO₂) ensures severe anoxia (<0.001% O₂). For each experiment, compounds were simultaneously tested under both oxic and hypoxic conditions against the HT-29 cell line and included TPZ as an independent internal control at the

front and back of the assay ($n = 2$). Final data was pooled from a series of seven independent experiments and is calculated using inter-experimental means. In all cases, 8-methyl-5-nitroquinoline was used as a second internal control to confirm that strict hypoxia was present during the experiment. (Siim et al., *Br. J. Cancer* **1994**, 70, 596–603). Cell cultures were grown in α MEM (Gibco) containing 5% heat inactivated FCS and maintained in exponential growth phase. For each individual experiment an appropriate number of cells were seeded (HT-29 = 1000) into wells in α MEM + 10% FCS + 10 mM added glucose + 100 μ M 2'-deoxycytidine (2'dCyd), and allowed to attach for 3 h. High glucose (final concentration 17 mM) and the presence of 2'-dCyd minimize hypoxia-induced cell cycle arrest. Replicates were then exposed to BTOs, using 2-fold serial dilutions in triplicate, for a further 4 h. Subsequently cells were washed free of compound using complete media (without glucose/2'-dCyd) and allowed to grow for 5 (oxic) or 6 (anoxic) days. Plates were stained as described previously (Wilson et al., *J. Med. Chem.* **1989**, 32, 31–38) and IC_{50} values determined.

IC_{50} = The concentration of drug (in micromolar) to reduce viable cell numbers to 50% of those of control cell cultures grown under the same conditions but not exposed to drug

RHT = Relative hypoxic toxicity is defined as the ratio of concentrations of Tirapazamine/test compound to give equal cell killing under hypoxic conditions.

HCR = Hypoxic cytotoxicity ratio is defined as the ratio of drug concentrations under aerobic and hypoxic condition to produce equal cell survival (50%) determined by proliferation assay

Table 3. Cytotoxicities of compounds of the invention under hypoxic conditions, hypoxic toxicity relative to Tirapazamine (RHT) and hypoxic selectivity (HCR) in proliferation assay

HT-29 IC₅₀			
Compound	IC₅₀ hypoxic (μM)	RHT	HCR
11	0.016	370	38
30	0.065	90	167
31	0.356	163	5.3
37	0.079	74	160
41	0.043	134	154
42	0.517	11.2	86.2
43	0.113	51.4	119
44	0.226	25.7	72.7
45	0.018	321	31
55	0.124	47	97
62	0.021	274	157
63	0.034	167	129
74	0.130	44.7	95
75	0.200	29	92
85	0.222	26	134
86	0.225	66	168
95	0.18	71	74
96	0.19	31	77
98	0.135	114	45
99	0.41	14	25
102	0.49	12	54
103	0.035	357	83
SiHa IC₅₀			
Compound	IC₅₀ hypoxic (μM)	RHT	HCR
30	0.031	121	41

55	0.05	72	124
75	0.07	53	110
86	0.105	35	136
95	0.076	48	89
96	0.10	37	78
98	0.075	63	30
102	0.16	23	53
103	0.01	309	118

The results of Table 3 clearly show that the compounds of the invention show large increases in cytotoxicity compared with Tirapazamine, while retaining selective killing under hypoxic conditions.

5


Wherein the foregoing description reference has been made to reagents, or integers having known equivalents thereof, then those equivalents are herein incorporated as if individually set forth.

10

While this invention has been described with reference to certain embodiments and examples, it is to be appreciated that further modifications and variations can be made to embodiments and examples without departing from the scope of the invention.

What we claim is:

1. A compound of Formula I,



wherein

Y_1 and Y_2 at one or more of the available carbons 5-8 on the benzo ring: are each independently selected from the following groups: halo, H, R, OH, OR, NO_2 , NH_2 , NHR , NR_2 , SH, SR, SO_2R , CF_3 , CN, CO_2H , CO_2R , CHO, COR, CONH_2 , CONHR or CONRR , cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

wherein each R is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and wherein the said optional substituents are each independently selected from; halo, OH, OR^1 , NO_2 , NH_2 , NHR^1 , NR^1R^1 , SH, SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN, CO_2H , CO_2R^1 , CHO, COR^1 , CONH_2 , CONHR^1 , CONR^1R^1 ;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR^1 , NH_2 , NHR^1 , NR^1R^1 , SH, SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN, CO_2H , CO_2R^1 , CHO, COR^1 , CONH_2 , CONHR^1 , CONR^1R^1 , and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;


wherein each R^1 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH_2 , NHR^2 , NR^2_2 or N(OH)R^2 wherein each R^2 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH, and

wherein X is selected from NH, NMe, CH₂, SO, SO₂, or O;

A is an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³

- 5 wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted or extended by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted
10 C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the DNA-targeting unit is any moiety of a molecular weight below 700 Daltons that
15 has an association constant (K) for binding to double-stranded random-sequence DNA of >10³ M⁻¹ at an ionic strength of 0.01 M at 20 °C, or a pharmacologically acceptable salt thereof.

2. The compound of Formula I as claimed in claim 1 wherein the DNA-targeting
20 unit is selected from one of formulae II- XVI,



wherein in structures **XI-XVI** R^6 is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and
 5 wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NO_2 , NH_2 , NHR^7 , NR^7R^7 , SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$;
 R^6 can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents
 10 are each independently selected from; halo, OH, OR^7 , NH_2 , NHR^7 , NR^7R^7 , SH, SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O,

N or S;

wherein each R^7 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^8 , NH_2 , NHR^8 , NR^8_2 or $N(OH)R^8$ wherein each R^8 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH;

wherein D represents up to four of the following groups as substituents at any available ring carbon position; H, R^9 , hydroxy, alkoxy, halogen, NO_2 , NH_2 , NHR^9 , NR^9_2 , SH, SR^9 , SO_2R^9 , CF_3 , CN, CO_2H , CO_2R^9 , CHO, COR^9 , $CONH_2$, $CONHR^9$ or $CONR^9R^9$, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino, wherein each R^9 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^{10} , NH_2 , NHR^{10} , NR^{10}_2 or $N(OH)R^{10}$ wherein each R^{10} is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH; and wherein any available ring carbon position of formulae **II** - **XVI** is optionally replaced by -N- when the valency and configuration of the formula allows, the point of attachment of formulae **II**- **XVI** to the A group defined above is represented by \blacklozenge ; and

wherein in formulae **XI**, **XII**, , m is selected from 2, 3 or 4, and

wherein in formulae **XI**, **XII**, **XV** and **XVI**, J is selected from CH or N; and wherein in formulae **XIII** and **XIV** n is selected from 0, 1 or 2; and wherein in formulae **XV** and **XVI** o is selected from 1 and 2.

3. The compound of Formula I as claimed in claim 2 wherein the DNA targeting unit is selected from one of formulae IV, V, VI, VII, VIII, or IX.

4. The compound of Formula I as claimed in claim 2 or claim 3 wherein D of the DNA targeting unit of Formulae **II** - **X** is H or Me.

5. The compound of Formula I as claimed in any one of claims 1 to 4 wherein X is NH or CH_2 .

6. The compound of Formula I as claimed in any one of claims 1 to 5 wherein Y_1

and Y₂ each represent H.

7. The compound of Formula I as claimed in any one of claims 1 to 5 wherein Y₁ represents OMe.
8. The compound of Formula I as claimed in any one of claims 1 to 7 wherein A is selected from -(CH₂)₆NH-, -(CH₂)₃NH(CH₂)₃NHCO-, -(CH₂)₃NMe(CH₂)₃NHCO-, -(CH₂)₃NH-, -(CH₂)₂NH(CH₂)₂NHCO- or -(CH₂)₂NMe(CH₂)₂NHCO-.
9. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₆NH-, the DNA targeting unit represents formula VII and D is H.
10. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₃NH(CH₂)₃NHCO-, the DNA targeting unit represents formula VIII and D is H.
11. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NH(CH₂)₂NHCO-, the DNA targeting unit represents formula VIII and D is H.
12. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₃NMe(CH₂)₃NHCO-, the DNA targeting unit represents formula VIII and D is H.
13. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₃NMe(CH₂)₃NHCO-, the DNA targeting unit represents formula IV and D is H.
14. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₃NMe(CH₂)₃NHCO-, the DNA targeting unit represents formula VI and D is H.

15. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is Me.
- 5 16. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula IX and D is Me.
- 10 17. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is 7-MeOCH₂CH₂O-, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H.
- 15 18. The compound of Formula I as claimed in claim 2 wherein X is CH₂-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H.
- 20 19. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is $-(CH_2)_2NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula XI and D is H.
- 20 20. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is 7-Me, Y₂ is H, A is $-(CH_2)_3NMeH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H.
- 25 21. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is 7-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VI and D is H.
- 30 22. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is 6-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H.
23. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is 6-Me, Y₂ is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents

formula VI and D is H.

24. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NMe(CH₂)₂NHCO-, the DNA targeting unit represents
5 formula VIII and D is H.

25. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NMe(CH₂)₂NHCO-, the DNA targeting unit represents
10 formula VI and D is H.


26. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NMe(CH₂)₂NHCO-, the DNA targeting unit represents
formula XI and D is Me.

27. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NMe(CH₂)₂NHCO-, the DNA targeting unit represents
15 formula VIII and D is Me.

28. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NH(CH₂)₂NHCO-, the DNA targeting unit represents formula
20 VI and D is H.

29. The compound of Formula I as claimed in claim 2 wherein X is NH-, Y₁ is H, Y₂ is H, A is -(CH₂)₂NH(CH₂)₂NHCO-, the DNA targeting unit represents formula
25 VIII and D is Me.

30. A compound of Formula I',



wherein

Y₁ represents at one or more of the available carbons 5-8 on the benzo ring the following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

Y₃ is selected from the following groups halo, H, R, OR, NH₂, NHR, NR₂, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

wherein each R of groups Y₁ and Y₃ is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from: halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from: halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;


wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

wherein A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂ or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₂₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, wherein each

R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and

wherein the DNA-targeting unit is any moiety of a molecular weight below 700 Daltons that has an association constant (K) for binding to double-stranded random-sequence DNA of $>10^3 \text{ M}^{-1}$ at an ionic strength of 0.01 M at 20 °C, or a pharmacologically acceptable salt thereof.

31. The compound of Formula I' as claimed in claim 30 wherein the DNA-targeting unit is selected from one of formulae **II- XVI**,



wherein in structures **XI** - **XVI** R^6 is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and
 5 wherein the optional substituents are each independently selected from; halo, OH, OR^7 , NO_2 , NH_2 , NHR^7 , NR^7R^7 , SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$;

R^6 can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents
 10 are each independently selected from; halo, OH, OR^7 , NH_2 , NHR^7 , NR^7R^7 , SH, SR^7 , imidazolyl, R^7 -piperazinyl, morpholino, SO_2R^7 , CF_3 , CN, CO_2H , CO_2R^7 , CHO, COR^7 , $CONH_2$, $CONHR^7$, $CONR^7R^7$, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N

or S;

wherein each R^7 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^8 , NH_2 , NHR^8 , NR^8_2 or $N(OH)R^8$

5 wherein each R^8 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH;

D represents up to four of the following groups as substituents at any available ring carbon position; H, R^9 , hydroxy, alkoxy, halogen, NO_2 , NH_2 , NHR^9 , NR^9_2 , SH, SR^9 , SO_2R^9 , CF_3 , CN, CO_2H , CO_2R^9 , CHO, COR^9 , $CONH_2$, $CONHR^9$ or $CONR^9R^9$,

10 cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino, wherein each R^9 independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR^{10} , NH_2 , NHR^{10} , NR^{10}_2 or $N(OH)R^{10}$ wherein each R^{10} is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 ,
15 CN, CO_2H or SH; and wherein any available ring carbon position of formulae II- XVI can also be optionally replaced by -N- when the valency and configuration of the formula allows, the point of attachment of formulae II- XVI to the A group defined above is represented by \blacklozenge ; and

wherein in formulae XI and XII, m is selected from 2, 3 or 4, and

20 wherein in formulae XI, XII, XV or XVI J is selected from CH or N; and

wherein in formulae XIII and XIV n is selected from 0, 1 or 2, and

wherein in formulae XV and XVI o is selected from 1 or 2.

25 32. The compound of Formula I' as claimed in claim 31 wherein the DNA targeting unit is selected from one of formulae III - IX.

33. The compound of Formula I' as claimed in claim 31 or claim 32 wherein D of the DNA targeting unit of Formulae II - X is H or Me.

30 34. The compound of Formula I' as claimed in any one of claims 30 to 33 wherein X is O, NH or CH_2 .

35. The compound of Formula I' as claimed in any one of claims 30 to 34 wherein

Y_1 represents H.

36. The compound of Formula I' as claimed in any one of claims 30 to 35 wherein A is selected from $-(CH_2)_6NH-$, $-(CH_2)_3NH(CH_2)_3NHCO-$, $-(CH_2)_3NMe(CH_2)_3NHCO-$, $-(CH_2)_3NH-$, $-(CH_2)_2NH(CH_2)_2NHCO-$ or $-(CH_2)_2NMe(CH_2)_2NHCO-$.

37. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VI and D is H.

38. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VI and D is H;

39. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H;

40. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula VI and D is H;

41. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_3NH(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

42. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_3NMe(CH_2)_3NHCO-$, the DNA targeting unit represents formula VIII and D is H;

43. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y_1 is H, A is $-(CH_2)_2NH(CH_2)_2NHCO-$, the DNA targeting unit represents formula VIII

and D is H;

44. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_2\text{NMe}(\text{CH}_2)_2\text{NHCO}-$, the DNA targeting unit represents formula
VIII and D is H;

45. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_3\text{NH}(\text{CH}_2)_3\text{NHCO}-$, the DNA targeting unit represents formula
VIII and D is Me;

46. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_3\text{NMe}(\text{CH}_2)_3\text{NHCO}-$, the DNA targeting unit represents formula
VIII and D is Me;

47. The compound of Formula I' as claimed in claim 31 X is O-, Y₁ is H, A is
 $-(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{NHCO}-$, the DNA targeting unit represents formula VIII
and D is Me;

48. The compound of Formula I' as claimed in claim 31 X is O-, Y₁ is H, A is
 $-(\text{CH}_2)_2\text{NMe}(\text{CH}_2)_2\text{NHCO}-$, the DNA targeting unit represents formula VIII
and D is Me;

49. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_3\text{NH}(\text{CH}_2)_3\text{NHCO}-$, the DNA targeting unit represents formula IX
and D is Me.

50. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_3\text{NMe}(\text{CH}_2)_3\text{NHCO}-$, the DNA targeting unit represents formula
IX and D is Me;

51. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H,
A is $-(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{NHCO}-$, the DNA targeting unit represents formula IX
and D is Me;

52. The compound of Formula I' as claimed in claim 31 wherein X is O-, Y₁ is H, A is $-(CH_2)_2NMe(CH_2)_2NHCO-$, the DNA targeting unit represents formula XI and D is Me

5

53. The compounds of Formula I' as claimed in any one of claims 30 to 52, wherein Y₃ represents CH₃, -CH₂CH₃ or NHCH₂CH₂N(CH₃)₂.

10

54. A method of therapy for treating cancers including the step of administering a compound of Formula I as defined in any one of claims 1 to 29 or a compound of Formula I' as defined in any one of claims 30 to 53 or a mixture thereof in a therapeutically effective amount to tumour cells in a subject.

15

55. The method of therapy according to claim 54 wherein the tumour cells are in a hypoxic environment.

20

56. The method of therapy according to claim 54 or claim 55 further including the step of administering radiotherapy to the tumor cells before, during or after the administration of the compound of Formula I as defined in any one of claims 1 to 29 or a compound of Formula I' as claimed in any one of claims 30 to 53 or a mixture thereof to the tumour cells.

25

57. The method of therapy according to any one of claims 54 to 56 further including the step of administering one or more chemotherapeutic agents to the tumor cells before, during or after the administration of the compound of Formula I as defined in any one of claims 1 to 29 or a compound of Formula I' as defined in any one of claims 30 to 53 or a mixture thereof to the tumour cells.

30


58. The method according to any one of claims 54 to 57 wherein the therapy can be administered alone or in combination with other chemotherapeutic agents or treatments, either simultaneously or sequentially dependent upon the condition to be treated.

59. The method according to claim 58 wherein the chemotherapeutic treatment is radiation therapy.

60. The method according to claim 59 wherein the chemotherapeutic agents are selected from one or more of :Cisplatin or other platinum-based derivatives, Temozolomide or other DNA methylating agents, Cyclophosphamide or other DNA alkylating agents, Doxorubicin, mitoxantrone, camptothecin or other topoisomerase inhibitors, Methotrexate, gemcitabine or other antimetabolites.

61. A pharmaceutical composition including a therapeutically effective amount of a compound of formula I as claimed in any one of claims 1 to 29 or a compound of formula I' as claimed in any one of claims 30 to 53 or a mixture thereof, a pharmaceutically acceptable excipient, adjuvant, carrier, buffer or stabiliser.

62. A method of making a compound of formula XVII



wherein

Y₁ and Y₂ at one or more of the available carbons 5-8 on the benzo ring: are each independently selected from the following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;


wherein each R is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, where each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; or a pharmacologically acceptable salt thereof,

including the step of coupling a compound (a) using a palladium reagent to form compound (b) which can then be converted into a compound of XVII as defined above;




wherein in compound (a)

V is halogen selected from Cl, Br or I and Y₁, Y₂ are as defined above in this

claim;

and wherein in compound (b) Y_1 , Y_2 are as defined above in this claim, W is selected from an optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, and optionally substituted C_{2-12} alkynyl group, wherein the optional substituents is selected from halo, OH, OR^6 , NO_2 , NH_2 , NHR^6 , NR^6R^6 , SH, SR^6 , imidazolyl, R^6 -piperazinyl, morpholino, SO_2R^6 , CF_3 , CN, CO_2H , CO_2R^6 , CHO, COR^6 , $CONH_2$, $CONHR^6$, $CONR^6R^6$, wherein each R^6 is independently selected from an optionally substituted C_{1-4} alkyl or an optionally substituted C_{2-4} alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH_2 , NHR^7 , NR^7_2 or $N(OH)R^7$ wherein each R^7 is independently selected from C_{1-4} alkyl, C_{2-4} alkenyl, OH, NO_2 , NH_2 , CF_3 , CN, CO_2H or SH.

63. A method of making a compound of formula XVII'



wherein Y_1 represents at one or more of the available carbons 5-8 on the benzo ring the following groups: halo, H, R, OH, OR, NO_2 , NH_2 , NHR , NR_2 , SH, SR, SO_2R , CF_3 , CN, CO_2H , CO_2R , CHO, COR, $CONH_2$, $CONHR$ or $CONRR$, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino; Y_3 is selected from the following groups H, R, OR, NH_2 , NHR , NR_2 , SO_2R , CF_3 , CN, CO_2H , CO_2R , CHO, COR, $CONH_2$, $CONHR$ or $CONRR$, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;


wherein each R of groups Y_1 and Y_3 is independently selected from an optionally substituted C_{1-6} alicyclic or an optionally substituted C_{3-6} cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR^1 , NO_2 , NH_2 , NHR^1 , NR^1R^1 , SH, SR^1 , imidazolyl, R^1 -piperazinyl, morpholino, SO_2R^1 , CF_3 , CN, CO_2H , CO_2R^1 , CHO, COR^1 , $CONH_2$, $CONHR^1$, $CONR^1R^1$;

R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR² or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³ or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, wherein each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and or a pharmacologically acceptable salt thereof;

including the steps of coupling a compound (a) using a palladium reagent to form compound (b) which is then converted into a compound of XVII' as defined above in this claim;



wherein in compound (a) V is halogen which is selected from Cl, Br or I; Y₁, X and A is as defined above in this claim;


and wherein in compound (b) Y₁, X and A are as defined above in this claim,

W is selected from an optionally substituted C₁₋₁₂alkyl, optionally substituted C₂₋₁₂alkenyl, and optionally substituted C₂₋₁₂alkynyl group, wherein the optional

substituents is selected from halo, OH, OR⁶, NO₂, NH₂, NHR⁶, NR⁶R⁶, SH, SR⁶, imidazolyl, R⁶-piperazinyl, morpholino, SO₂R⁶, CF₃, CN, CO₂H, CO₂R⁶, CHO, COR⁶, CONH₂, CONHR⁶, CONR⁶R⁶, wherein each R⁶ is independently selected

from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR⁷, NR⁷₂ or N(OH)R⁷ wherein each R⁷ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH.

64. A compound of formula XVIII



wherein

Y₁ and Y₂ at one or more of the available carbons 5-8 on the benzo ring: are each independently selected from the following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

wherein each R is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;


R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents

are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O,

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂, or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, wherein each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; or a pharmacologically acceptable salt thereof.

65. A compound of formula XVII'



wherein

Y₁ represents at one or more of the available carbons 5-8 on the benzo ring the

following groups: halo, H, R, OH, OR, NO₂, NH₂, NHR, NR₂, SH, SR, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino;

5 Y₃ is selected from the following groups H, R, OR, NH₂, NHR, NR₂, SO₂R, CF₃, CN, CO₂H, CO₂R, CHO, COR, CONH₂, CONHR or CONRR, cyclic alkylamino, imidazolyl, alkylpiperazinyl and morpholino

wherein each R of groups Y₁ and Y₃ is independently selected from an optionally substituted C₁₋₆ alicyclic or an optionally substituted C₃₋₆ cyclic alkyl group, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NO₂, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹;

10 R can also represent an optionally substituted aryl or an optionally substituted heteroaryl group having up to 12 carbon atoms, and wherein the optional substituents are each independently selected from; halo, OH, OR¹, NH₂, NHR¹, NR¹R¹, SH, SR¹, imidazolyl, R¹-piperazinyl, morpholino, SO₂R¹, CF₃, CN, CO₂H, CO₂R¹, CHO, COR¹, CONH₂, CONHR¹, CONR¹R¹, and each heteroaryl group contains one or more heteroatoms in its ring system which are each independently selected from O, N or S;

wherein each R¹ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR², NR²₂ or N(OH)R² wherein each R² is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH, and

wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

25 A represents an optionally substituted C₁₋₁₂ alkyl group wherein the optional substituents are each independently selected from OH, OR, NH₂, NHR³, NR³₂ or N(OH)R³ wherein each R³ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN, CO₂H or SH; and wherein the optionally substituted C₁₋₁₂ alkyl chain is optionally interrupted by one or more heteroatom containing linkage

30 moieties selected from O, NH, NR⁴, CONH, CONR⁴, NHCO, NR⁴CO, wherein each R⁴ is independently selected from an optionally substituted C₁₋₄ alkyl or an optionally substituted C₂₋₄ alkenyl group and wherein the optional R⁴ substituents are each independently selected from OH, OR, NH₂, NHR⁵, NR⁵₂ or N(OH)R⁵ wherein each R⁵ is independently selected from C₁₋₄ alkyl, C₂₋₄ alkenyl, OH, NO₂, NH₂, CF₃, CN,

CO₂H or SH; and

wherein X represents NH, NMe, CH₂, SO, SO₂, or O;

or a pharmacologically acceptable salt thereof.

5 66. A method of making a compound of Formula I defined above in any one of claims 1 to 29 including the steps of

1 preparing a compound of Formula XVIII as defined above in claim 64;
and

10 2 coupling the compound of Formula XVIII with a DNA targeting agent as defined in claim 2 to provide a compound of Formula I.

67. A method of making a compound of Formula I' defined in any one of claims 30 to 53 including the steps of

15 1 preparing a compound of Formula XVII' as defined above in claim 65;
and

2 coupling the compound of Formula XVII' with a DNA targeting agent as defined above in claim 31 to provide a compound of Formula I'.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ03/00210

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl. ⁷ : C07D 253/10, 401/12, 403/12; A61K 31/53; A61P 35/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) STN Substructure Search based on compounds of Formulae I, I', XVIII and XVII'		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Biochemical Pharmacology, Vol. 65 (11), 2003, Delahoussaye et al, "Improved potency of the hypoxic cytotoxin tirapazamine by DNA-targeting", pages 1807-1815 See especially compound SN 26955, page 1808	1-67
X	Journal of Heterocyclic Chemistry, Vol. 30(2), 1993, Parrick et al, "The Synthesis of a Potential Anti-Cancer Agent Containing the Caffeine and 1,2,4-Benzotriazine Moieties", pages 323-327 See especially Compound 6, page 325	30-53, 61, 63, 65,67
X	Anti-Cancer Drug Design, Vol. 10(3), 1995, Mehta et al, "Potential bioreductively activated hypoxia probes and post-irradiation radiosensitizers related to NITP", pages 227-241 See especially Compound 3, page 228	30-53, 61,63,65,67
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 10 December 2003		Date of mailing of the international search report 17 DEC 2003
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustralia.gov.au Facsimile No. (02) 6285 3929		Authorized officer R.L. POOLEY Telephone No : (02) 6283 2242

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ03/00210

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages (Remove spaces when completed if the page is too long)	Relevant to claim No.
X	WO 91/04028 A (SRI INTERNATIONAL) 4 April 1991 See especially Examples 6, 10, 11, pages 14-21	64, 65
X	EP 972517 A2 (THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY) 19 January 2000 See especially Examples 6, 10, 11, 17, 18	64, 65
X	US 5827850 A (BROWN et al) 27 October 1998 See Column 4, lines 55-60	64, 65
X	DD 272591 A (NICLAS et al) 18 October 1989 See especially Examples 5,6	64, 65
P,X	Journal of Medicinal Chemistry, Vol. 46(1), 2003, Hay et al, "Structure-Activity Relationships of 1,2,4-Benzotriazine 1,4-Dioxides as Hypoxia-Selective Analogues of Tirapazamine", pages 169-182 See table 1, page 172	64, 65
X	Chemical Abstracts, Volume 129, Abstract 339530 (& Anti-Cancer Drug Design, Vol. 13 (6), 1998, Kelson et al, "1,2,4-Benzotriazine 1,4-dioxides. An important class of hypoxic cytotoxins with antitumour activity", pages 575-592) See for example RN 166182-17-8, 166182-18-9, 215034-31-4, 215535-59-4	64, 65
X	Chemical Abstracts, Volume 116, Abstract 187502 (& International Journal of Radiation Oncology, Biology, Physics, Vol. 22(4), 1992, Minchinton et al, "Second generation 1,2,4-benzotriazine 1,4-di-N-oxide bioreductive antitumour agents: pharmacology and activity in vitro and in vivo", pages 701-705)	64, 65
X	Chemical Abstracts, Volume 114, Abstract 164101 (& Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry, Vol 12, 1990, Argyropoulos et al, "Cycloadditions of nitril oxides with benzofuran n-oxides", pages 3277-3287) See RN 132922-33-9, 132922-34-0, 132922-35-1, 132922-36-2	65
X	Chemical Abstracts, Volume 112, Abstract 171760 (& Biochemical Pharmacology, Vol. 39(4), 1990, Tocher et al, "Electrochemical studies and DNA damaging effects of the benzotriazine-N-oxides", pages 781-786) See RN 121135-28-2, 121140-01-0	65
X	Chemical abstracts Volume 111, Abstract 3393 (& International Journal of Radiation Oncology, Biology, Physics, Vol. 16 (4), 1989, Zeman et al, "Structure-activity relationships for benzotriazine di-N-oxides", pages 977-981) See RN 121135-27-1, 121135-28-2, 121135-30-6, 121140-01-0	64, 65

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/NZ03/00210

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member					
WO	91/04028	AU	646794	DE	68929050	EP	478545
		JP	3034541	NO	920747		
EP	972517	AU	74117/94	CA	2132578	DE	69424915
		EP	649658	JP	7215882	RU	2148406
		US	5484612	US	5670502	US	6121263
		US	6277835				
US	5827850	AU	68548/96	AU	69690/98	CA	2232989
		CN	1202827	EP	866709	EP	1044005
		JP	11511479	JP	2001523248	RU	2166946
		US	6153610	WO	9711699	WO	9847512
DD	272591						
							END OF ANNEX

(To add more lines press TAB at end of last row, remove paragraph marker to join up 'END OF ANNEX' box)